


Aloha,

This cover letter is intended to provide an overview of the Laupāhoehoe Draft Management Plan and Environmental Assessment process including process status. It will be periodically updated as needed.

- Together the Department of Land and Natural Resources - Division of Forestry and Wildlife (DOFAW), the United States Department of Agriculture – United States Forest Service (USFS), and the Laupāhoehoe Advisory Council (LAC) have held 11 public meetings between 2012-2014, during which the draft management plan was developed and discussed.
- At the April 8, 2015 LAC meeting, after reviewing the latest internal version of the draft plan, the council voted to release the draft plan to the public.
- An informational poster session to introduce the draft plan is planned for June 17, 2015 from 6-8 pm at the Laupāhoehoe Community Public Charter School (LCPCS) cafeteria. Hard copies of the draft plan will be available at the LCPCS school office, Hilo and Laupāhoehoe public libraries and by request no later than May 15.
- A joint state-federal Environmental Assessment (EA) will be completed for the management plan and this formal process is expected to begin in several months. Ultimately, the management plan will be finalized and approved by the Board of Land and Natural Resources.
 - o Formal comments and questions related to the draft plan will be taken in conjunction with the EA process. If you would like to be involved in this process please contact Mel Dean, mkdean@hawaii.edu or 808-854-2651 to provide your contact information. We will inform you as to when the EA process has started with instructions on insuring your contributions are captured in the EA process.



Mahalo

Cover letter updated on 4/15/2015

LAUPĀHOEHOE FOREST

DRAFT MANAGEMENT PLAN



2015

DEPARTMENT OF LAND AND NATURAL RESOURCES
DIVISION OF FORESTRY AND WILDLIFE

UNITED STATES DEPARTMENT OF AGRICULTURE
UNITED STATES FOREST SERVICE



The Forest Service of the U.S. Department of Agriculture is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives -- as directed by Congress --to provide increasingly greater service to a growing Nation.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call (800)795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.





April 2015

**State of Hawai'i
Department of Land and Natural Resources
Division of Forestry and Wildlife
1151 Punchbowl Street, Room 325
Honolulu, HI 96813**

**USDA Forest Service
Pacific Southwest Research Station
Institute of Pacific Islands Forestry
60 Nowelo St.
Hilo, HI 96720**



TABLE OF CONTENTS

TABLE OF CONTENTS	i
ACRONYMS	iii
ACKNOWLEDGEMENTS	1
EXECUTIVE SUMMARY	2
INTRODUCTION	3
Background	3
Guiding Principles	3
General Information	4
<i>Location</i>	4
Management Focal Areas	7
<i>Natural Resources</i>	7
<i>Research</i>	8
<i>Education and Outreach</i>	8
<i>Public Access and Recreation</i>	9
<i>Infrastructure</i>	9
DESCRIPTION AND CURRENT CONDITION	10
Land Use	10
<i>Land Designation and Management</i>	10
<i>Land Use History</i>	13
<i>Public Access and Recreation</i>	15
Vehicle Access	15
Trails	17
Mountain Biking	17
Hunting	17
Camping	19
Gathering	19
<i>Existing Infrastructure and Facilities</i>	19
Roads	19
Facilities/Structures	19
<i>Regional Partnerships</i>	20
Site Description (Physical and Biological Resources)	20
<i>Topography, Climate, Geology, and Soils</i>	20
<i>Water Resources</i>	24
<i>Forest Ecosystems</i>	26
<i>Wildlife</i>	32
Cultural Resources and Traditional Practices	35
THREATS TO NATURAL AND CULTURAL RESOURCES	37
Non-Native Plants and Animals	37
Disease	40
Climate Change and Natural Disturbances	40
Illegal Human Activity	40
Wildfire	41



Threats to Cultural Resources	42
OVERVIEW OF EXISTING MANAGEMENT	43
Natural Resources	43
<i>Forest Protection and Management</i>	43
<i>Invasive Non-Native Plant Control</i>	43
<i>Rare Species Restoration</i>	44
<i>Monitoring</i>	44
Research.....	44
Education and Outreach.....	46
Public Access and Recreation.....	46
Infrastructure.....	46
PROPOSED MANAGEMENT PROGRAM	47
Natural Resources	47
<i>Forest Protection and Management</i>	47
<i>Invasive Non-Native Plant Control</i>	50
<i>Rare Species Restoration</i>	51
<i>Monitoring</i>	52
<i>Wildfire Prevention and Response</i>	53
Research.....	53
Education and Outreach.....	56
Public Access and Recreation.....	58
Infrastructure.....	63
ADMINISTRATION	64
Coordination	64
Permitting.....	64
Cultural Resources.....	64
Biosecurity	64
Safety	65
Budget and Staffing	65
REFERENCES	71
Appendix A – Other Available Resources.....	77
Appendix B – Biosecurity Protocols.....	78
Appendix C - List of Research Projects to Date.....	80



ACRONYMS

BLNR	Board of Land and Natural Resources
DAR	Division of Aquatic Resources
DLNR	Hawai‘i Department of Land and Natural Resources
DOCARE	Division of Conservation and Resource Enforcement
DOFAW	Division of Forestry and Wildlife
EA	Environmental Assessment
FIA	Forest Inventory and Analysis Program
FR	Forest Reserve
HIPPNET	Hawai‘i Permanent Plot Network
IPIF	Institute of Pacific Islands Forestry
LAC	Laupāhoehoe Advisory Committee
NAGPRA	Native American Graves Protection and Repatriation Act
NARS	Natural Area Reserves System
OCCL	Office of Conservation and Coastal Lands
PEPP	Plant Extinction Prevention Program
PIPES	Pacific Internship Programs for Exploring Science
PSW	Pacific Southwest Research Station
SHPD	State Historic Preservation Division
UH	University of Hawai‘i
USFS	USDA Forest Service
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey
VRPF	Volcano Rare Plant Facility
YCC	Youth Conservation Corps



ACKNOWLEDGEMENTS

The DOFAW and the USFS would like to thank the numerous stakeholders and community members that provided input and guidance during the writing of this plan. Special thanks in particular to the LAC members who participated in many meetings and field trips during plan writing and contributed a great deal of time and energy to the process.



Aerial view of Laupāhoehoe Forest



EXECUTIVE SUMMARY

The Laupāhoehoe Forest Management Plan seeks to comprehensively protect and preserve Laupāhoehoe Forest while enhancing public use and benefits through education, recreation, outreach, demonstration, and research activities. This plan documents the history of the forest, describes its current condition, provides an overview of current management activities and agency missions, recognizes the role intact forests play in providing clean freshwater for downstream human and wildlife populations and in supporting healthy coastal marine resources, and recommends management actions. The plan is the management vision for approximately fifteen years and provides:

- Guidance and recommendations to the Hawai‘i Department of Land and Natural Resources, Division of Forestry and Wildlife (DOFAW) and the US Department of Agriculture, US Forest Service (USFS) from stakeholders including the Laupāhoehoe Advisory Council (LAC)
- Prioritized recommendations on how to protect and preserve the area, as well as continue and enhance human use
- A compilation of natural and cultural history, resources, and research
- Documentation of current forest conditions and threats
- A planning and management tool for the DOFAW and the USFS to use to determine priorities, work plans, staffing requirements, budget requests, and more
- Funding guidance and a document that enables the DOFAW and the USFS to ask for resources necessary to protect Laupāhoehoe Forest

The plan was jointly developed by DOFAW, the USFS, and the Laupāhoehoe Advisory Council (LAC) through a collaborative planning process. Formed in December 2010, the LAC is a community-based advisory council that provides guidance and consultation to DOFAW and USFS on issues of management, research, and education in Laupāhoehoe Forest.

This draft plan includes proposed management actions to protect natural and cultural resources within Laupāhoehoe Forest while also enhancing compatible human uses. Protecting resources includes addressing the threats of invasive non-native species as well as climate change. The plan will be a guiding document for DOFAW and the USFS for management actions, including background information for why those actions were chosen. The management plan identifies objectives and strategic actions related to Natural Resources, Research, Education and Outreach, Public Access and Recreation, and Infrastructure. Objectives Include:

- Natural Resources - Protect, manage and restore native ecosystems and species
- Research - Provide lands for conducting research that serves as a basis for the restoration, conservation and management of tropical forest ecosystems
- Education and Outreach - Serve as a center for forest education, training, demonstration, and outreach on tropical forests, conservation biology, and natural resource management for groups ranging from school children to land managers, scientists and the general public
- Public Access and Recreation - Improve appropriate public access and recreational opportunities consistent with maintaining native natural resources, cultural resources and the wilderness character of these lands
- Infrastructure - Provide and maintain infrastructure and facilities to meet forest goals



INTRODUCTION

Background

The 12,343 acre (ac) (5,134 hectare (ha)) Laupāhoehoe Forest area consists of two state-managed parcels of land: 4,449 ac of state land designated as Forest Reserve (FR), and 7,894 ac of land designated as a Natural Area Reserve (NAR) (Figure 1). Both of these programs are under the state of Hawai‘i Department of Land and Natural Resources (DLNR), Division of Forestry and Wildlife (DOFAW). In addition, the Laupāhoehoe Forest is designated as part of the Hawai‘i Experimental Tropical Forest (HETF).

In 1992, the Hawai‘i Tropical Forest Recovery Act authorized the establishment of the HETF to serve as a center for long-term research and a focal point for developing and transferring knowledge and expertise for the management of tropical forests. In 2007, the HETF was formally established. The HETF consists of two units, one of which is the Laupāhoehoe Forest; the other unit is the Pu‘u Wa‘awa‘a Forest located on the leeward side of Hawai‘i Island (Figure 2).

USDA Forest Service (USFS), Pacific Southwest (PSW) Research Station’s Institute of Pacific Islands Forestry (IPIF), based in Hilo, works cooperatively with the state of Hawai‘i to coordinate research, management, and educational activities and to jointly develop and implement research and education, and management plans for the experimental forest. Land management and protection responsibilities remain with DLNR-DOFAW and these lands are managed under relevant state laws and regulations.

Guiding Principles

The management plan of Laupāhoehoe Forest is based on the overall vision for the area as part of the HETF, as well as the purpose of the land designations for the two parcels of state lands that are part of the Laupāhoehoe Forest. The plan also incorporates the values and interests of community members and stakeholder groups and attempts to balance the need for increased forest protection and management while enhancing compatible human uses.

The HETF vision for Laupāhoehoe Forest is as an important research, education, and demonstration forest where globally relevant activities are conducted to benefit the people and ecosystems of Hawai‘i, all Pacific islands and the tropical world. The forest will provide research, demonstration, training and education opportunities for scientists, post-doctoral fellows, graduate and undergraduate students, K-12 school children who are the future generations of land managers/stewards, forest users, landowners, and scientists in Hawai‘i.

DLNR lands within Laupāhoehoe Forest designated as NAR are managed by DOFAW under the state’s Natural Area Reserves System (NARS). The NARS seeks to protect the best remaining examples of the state’s unique ecosystems and strives to actively manage these reserves in order to preserve the unique characteristics that make these areas an integral part of the natural heritage of Hawai‘i. Reflecting this, the mission of the NARS program is: “The NARS exists to ensure the highest level of stewardship for Hawai‘i’s natural resources through acquisition, active management, and other strategies.”



DLNR lands within Laupāhoehoe Forest designated as FR are managed by DOFAW under the state's Forest Reserve System. The Forest Reserve System was created by the Territorial Government of Hawai'i through Act 44 on April 25, 1903 to protect key forested watersheds. In addition, forest reserves are managed to provide recreational opportunities; aesthetic benefits; native; cultural resources; and threatened and endangered species habitat protection among many other things (cultural use and gathering, hunting).

Laupāhoehoe Advisory Council (LAC)

Formed in December 2010, the LAC is a community based advisory council that provides guidance and consultation to DOFAW and USFS relating to management, research, and education activities in the Laupāhoehoe Forest. The LAC's mission statement: Within our Hawaiian culture establish community support in the development of strategies for the long-term management, protection and utilization of existing and potential forest resources.



Laupāhoehoe Advisory Council field trip

Management Plan Development Process

The management plan for Laupāhoehoe Forest was jointly developed by DOFAW, USFS and the LAC through a collaborative planning process. The Management Plan is a long-term management vision and covers a time-frame of fifteen years.

This plan serves as a DOFAW site-specific plan for the two state-managed parcels of land within Laupāhoehoe Forest. The plan provides a brief history of the FR and NAR, a description of cultural and natural resources, and proposed management actions for the area.

The plan also serves as the USFS plan for Laupāhoehoe Forest Unit of the experimental forest. The specific management, research, and education objectives and activities for the HETF are described in an overarching USFS Master Plan document and in individual plans for each of the two units: the Laupāhoehoe Forest plan (this document) and the Management Plan for the Ahupua'a of Pu'u Wa'awa'a and the Makai Lands of Pu'u Anahulu (state of Hawai'i 2003).

General Information

Location

Laupāhoehoe Forest is situated on the eastern, windward flanks of Mauna Kea in the North Hilo District on the island of Hawai'i (TMK #'s (3) 3-7-001:002, (3) 3-7-001:012). Laupāhoehoe Forest stretches from from about 1,700 to 6,100 feet (ft) (518–1860 meters (m)) elevation and includes several stream drainages (Figure 1). The forest is located on the Hamakua Coast above the town of Laupāhoehoe, which has an estimated total population of 614 (U.S. Census Bureau 2012).



Figure 1





Figure 2



Management Focal Areas

Natural Resources

The protection and effective management of natural resources particularly forested watersheds, unique native Hawaiian ecosystems and threatened and endangered species are a priority for the two parcels of state lands within Laupāhoehoe Forest, and the original purpose for their designations as state NAR and FR. These natural resources require active management in order to persist for the benefit of current and future generations. Key aspects of natural resources management addressed in this plan include:

- Protection and management of watershed and water resources for downstream human and wildlife populations
- Protection of unimpeded surface and/or groundwater flows to the ocean
- Overview of existing natural resources and description of known threats to those resources
- Description of past and current management actions
- Management actions proposed to effectively address threats to natural resources
- Protection and restoration of native Hawaiian ecosystems and species at Laupāhoehoe Forest, including rare and endangered plants and animals
- Linking management to research to develop more effective management techniques
- Monitoring results to assess the effectiveness of management actions, and adjusting management, if necessary, in an adaptive management approach



Natural resources in Laupāhoehoe Forest include numerous streams



Research

Laupāhoehoe Forest is part of a national network of experimental forests and provides researchers with a globally unique opportunity to study wet tropical ecosystems within a larger landscape that stretches across an environmental gradient ranging from sea level to alpine ecosystems. Information from research is critically needed by land managers to help effectively manage the area, particularly the threats posed by invasive species and changing conditions due to climate change impacts. Research can also help us to better understand basic ecological and evolutionary processes, and develop adaptable approaches and effective ways to sustain, enhance and restore the capacity of ecosystems to provide goods and services under changing environmental conditions. Primary areas of research to be promoted and emphasized at Laupāhoehoe Forest include:

- Long-term trends in native montane tropical forest ecosystem conditions and dynamics
- Montane tropical forest ecosystem structure, dynamics, and threats
- Impacts of climate change on wet tropical ecosystems and the services they provide
- Impacts of invasive, non-native plants and animals on montane tropical forest species, communities and ecosystems
- Methods of restoring and maintaining ecosystem function and services in the face of global change
- Methods of restoration that integrate cultural and scientific knowledge
- Methods and approaches that enhance community based collaborative stewardship of natural resources

Education and Outreach

Laupāhoehoe Forest will serve as a center for demonstration, education, training, and outreach on tropical forestry, conservation biology, and natural resources research and management. Education and outreach goals will be accomplished through a strong reliance on partnerships. Education and outreach goals span six focal areas:

- *Formal training for professionals:* Provide work experience and professional development in ecology, conservation, and restoration.
- *Community outreach:* Communicate research findings and management goals, and foster a connection to nature and forest stewardship through engagement activities that involve the public through collaboration with partners including K-12 education program partners.
- *Cultural training:* Exposing researchers, students and managers



Educational programs include hands-on activities such as growing trees for forest restoration projects



to cultural knowledge directed at managing forest and coastal resources, including native Hawaiian perspectives and approaches.

- *Demonstration for managers*: Delivery of information, tools and techniques to managers through demonstration research, conservation, and restoration projects.
- *Student research*: Foster and support undergraduate and graduate level research opportunities and research internships.
- *Academic education*: Collaborate with universities to integrate college courses and support courses via state and federal facilities.

Public Access and Recreation


Laupāhoehoe Forest is protected and managed by the state for the benefit of the people of Hawai‘i, and is open to the public for various recreational and cultural uses. The use of Laupāhoehoe Forest for activities such as hiking, hunting and traditional and cultural practices, are high priorities for the local community. Management to encourage and enhance public access and recreation includes:

- Encouraging low-impact recreational activities and improvements such as pedestrian trails that are consistent with the remote, wilderness nature of Laupāhoehoe Forest to minimize recreational impacts on natural and cultural resources.
- Improving recreational opportunities by maintaining existing trails and creating new trails to establish a connected trail system.
- Securing new public access routes and appropriate areas for additional forest protection through easements, land acquisition and/or public access agreements with adjacent landowners.
- Continuing to facilitate public hunting in Laupāhoehoe Forest by securing and improving access and trails.
- Providing trails that offer educational and outreach experiences for the public to highlight conservation, restoration and management demonstration sites.

Infrastructure

Facilities and infrastructure such as roads and trails are essential to facilitate and support the use of HETF for research and education, for natural resources management activities and for public recreation. Goals for the infrastructure and facilities at Laupāhoehoe Forest include:

- Develop and maintain roads, trails, cabins/shelters and campsites for the functions of the HETF and for resources management actions as well as for public recreational use.
- Ensure facilities and infrastructure has minimal impacts on the environment and natural and cultural resources.



Laupāhoehoe Forest access trail



DESCRIPTION AND CURRENT CONDITION

Land Use

Land Designation and Management

State lands in the Laupāhoehoe Forest are under overall administration of DLNR. The Board of Land and Natural Resources (BLNR) sets policies for the Department. DOFAW is the Division of DLNR charged with management of the lands which are designated as both NAR and FR. Other relevant planning documents associated with Laupāhoehoe Forest are provided in Table 1.

Both DOFAW and IPIF have responsibilities in the management of the experimental forest and IPIF activities are authorized under a 2006 Cooperative Agreement between the BLNR and the USDA Forest Service as well as a permit for use of state lands. Land management and protection responsibilities remain with the state and are managed under relevant state laws and regulations. The IPIF participates in the administration of research and education activities; authority for signing of all permits lies with DOFAW. The IPIF administers the research/education infrastructure.

The Hawai‘i State Constitution Article 11 states: “For the benefit of present and future generations, the state . . . shall conserve and protect Hawaii's natural beauty and all natural resources, including land, water, air, minerals and energy sources, and shall promote the development and utilization of these resources in a manner consistent with their conservation and in furtherance of the self-sufficiency of the state. All public natural resources are held in trust by the state for the benefit of the people.”

DOFAW has management responsibility for the 4,449 ac (1,800 ha) Laupāhoehoe section of Hilo FR, which is part of the state Forest Reserve System (Figure 1). Hilo FR was originally established in 1905 for the purpose of watershed protection, and various parcels have been added and withdrawn from the FR since it was established. The FR portion of the Laupāhoehoe Forest is approximately 1,700 to 6,100 ft (518–1,860 m) elevation.

The Forest Reserve System was created by the Territorial Government of Hawai'i through Act 44 on April 25, 1903. With Hawaii's increase in population, expanding ranching industry, and extensive agricultural production of sugarcane and later pineapple, early territorial foresters recognized the need to protect mauka (upland) forests to provide the necessary water for agriculture and surrounding communities. The Forest Reserve System is managed under the guidance of the Hawai‘i Revised Statutes ([Chapter 183](#)) and associated Hawai‘i Administrative Rules ([Chapter 104](#)). Through these directives, DOFAW focuses its resources to protect, manage, restore, and monitor the natural resources of the Forest Reserve System. The Forest Reserve System accounts for over 448,000 acres of state managed land on the island of Hawai‘i.

Laupāhoehoe Forest includes approximately 7,894 ac (3,195 ha) withdrawn from the FR and designated as a NAR in 1983 by Executive Order 3168 (Figure 1). The NAR was designated to protect wet forests of koa (*Acacia koa*) and ‘ōhia (*Metrosideros polymorpha*), wet grasslands and streams, all of which provide important habitat for plants and animals, including rare species. The NAR portion of Laupāhoehoe Forest includes lands from approximately 1,700 to 4,600 ft (518–1,402 m) on the slopes of windward Mauna Kea. The Natural Area Reserves System (NARS) was



created in 1971 by the Hawai‘i State Legislature to “preserve in perpetuity specific land and water areas which support communities, as relatively unmodified as possible, of the natural flora and fauna, as well as geological sites, of Hawai‘i (HRS § 195-1)”. The legislature further found that these unique natural assets should be protected and preserved, both for the enjoyment of future generations and to provide baselines against which changes to Hawaii’s environment can be measured. The NARS is administered by DLNR-DOFAW. NARS Commission members act in an advisory capacity for the Board of Land and Natural Resources, which sets policies for the Department. Hawai‘i Administrative Rules 13-209 relate to the management of the NARS.

The NARS presently consists of 20 reserves on five islands, encompassing more than 123,000 ac (49,776 ha) of the state’s most unique ecosystems. The diverse areas found in the NARS range from marine and coastal environments to alpine desert, and from fresh lava flows to wet forests. These areas often serve as habitat for rare native plants and animals, many of which are on the verge of extinction. The NARS also include important watersheds, contributing to Hawai‘i’s sources of drinking water. Finally, the NARS forms an important part of the scenic landscape and contributes to the natural beauty of Hawai‘i, contributing to the islands’ overall appeal to visitors. Some of the most recognizable and visited NARS include Mauna Kea Ice Age NAR (Hawai‘i), Ka‘ena Point NAR (O‘ahu), and ‘Āhihi-Kīna‘u NAR (Maui).

The DOFAW website located at <http://dlnr.hawaii.gov/2013dofaw/> provides general information on both NAR and FR programs and policies across the state.

Laupāhoehoe Forest is also designated as part of the HETF, which was formally established in 2007. The HETF currently includes two units: the Laupāhoehoe Forest Wet Forest Unit and the Pu‘u Wa‘awa‘a Dry Forest Unit (Figure 2). The USFS works with the state in the management of the experimental forest.

The purposes of the HETF are to: (1) understand how to restore, preserve, and sustainably manage native tropical forests, streams, and watersheds of the Pacific, and to provide information to those managing these landscapes; (2) provide a center for demonstration, education, training, and outreach on tropical forestry, conservation biology, and natural resources research and management; (3) provide sites dedicated to long-term research on tropical forestry, ecology, hydrology, conservation biology, and natural resource management; and (4) foster research cooperation and collaboration between state and federal agencies, and among agencies and other institutions in tropical forestry research.



Laupāhoehoe Forest provides a site for long-term research on tropical forests



Other DLNR Divisions and DOFAW Programs also have various responsibilities related to Laupāhoehoe Forest:

- Division of Conservation and Resource Enforcement (DOCARE) - DOCARE has full police powers and is responsible for the enforcement of state laws and rules at Laupāhoehoe Forest, including laws regulating hunting and protection of resources (e.g. illegal harvesting, vandalism etc.).
- Nā Ala Hele - The Hawai‘i Statewide Trail and Access System is a DOFAW Program that has responsibility for trails and access. This program regulates and manages specific trails and accesses; conducts trail and access inventory; investigates potential and needed trails and accesses; examines legal issues; acts as point of contact for trail and access information and issues; and conducts trail and access advisory council meetings.
- Division of Aquatic Resources (DAR) - Manages marine and freshwater resources through programs in commercial fisheries and aquaculture; aquatic resources protection, enhancement and education; and recreational fisheries.
- State Historic Preservation Division (SHPD) - Works to preserve and sustain reminders of earlier times which link the past to the present. SHPD has three branches - History and Culture, Archaeology, and Architecture.
- Office of Conservation and Coastal Lands (OCCL) - Responsible for overseeing private and public lands that lie within the State Conservation District, including designated Conservation District lands in Laupāhoehoe Forest. Conservation District subzones define identified land uses which may be allowed by discretionary permit or some sort of approval from the DLNR or BLNR. NAR portions of Laupāhoehoe Forest are in the Protective Subzone and lands within the FR are in the Resource Subzone.



Table 1. *Related Federal, State and County Planning Documents*

Planning Document	Comment
Hamakua Community Development Plan (plan under development)	County of Hawai‘i plan http://www.hawaiicountycdp.info/hamakua-cdp
The Rain Follows the Forest - A Plan to Replenish Hawaii’s Source of Water (DLNR 2011)	Laupāhoehoe Forest is identified as a priority watershed area on the island of Hawai‘i
DOFAW Statewide Assessment and Resource Strategy (SWARS) (DLNR 2010)	Identifies areas of greatest need and opportunity for forests in Hawai‘i and develops a long-term management strategy. Objectives include: 1.1. Identify and conserve high-priority forest ecosystems and landscapes; 2.2. Identify, manage and reduce threats to forest and ecosystem health; 3.3. Enhance public benefits from trees and forests; 3.1. Protect and enhance water quality and quantity; 3.5. Protect and enhance wildlife and fish habitat; 3.7. Manage and restore trees/forests to mitigate and adapt to global climate change.
Draft Mauna Kea Watershed Alliance Management Plan (Stewart 2010)	DOFAW and USFS are members of the Mauna Kea Watershed Alliance and Laupāhoehoe Forest is included in this partnership area. The Mauna Kea Watershed alliance includes major landowners on Mauna Kea with a shared interest to protect the ‘aina by working together to manage threats that occur across common land ownership



Planning Document	Comment
	boundaries, pooling limited resources to achieve conservation goals, and promoting collaboration in protecting vital resources across large landscapes.
U.S. Fish and Wildlife Revised Recovery Plan for Hawaiian Forest Birds (USFWS 2006)	Supports recovery actions 1 and 2: protect and manage ecosystems for the benefit and recovery of native forest birds.
Hawai'i Comprehensive Wildlife Conservation Strategy (DLNR 2005)	Statewide strategy for the conservation of native wildlife and plants. Identifies species of greatest conservation need.
County of Hawai'i General Plan (County of Hawai'i 2005)	8.2(c) Protect and promote the prudent use of Hawaii's unique, fragile, and significant environmental and natural resources. 8.2 (d) Protect rare or endangered species and habitats native to Hawai'i. 8.3 (b) Encourage a program of collection and dissemination of basic data concerning natural resources. 8.3 (e) Encourage an overall conservation ethic in the use of Hawai'i resources by protecting, preserving, and conserving the critical and significant natural resources of the County. 8.3 (o) Encourage the continued identification and inclusion of unique wildlife habitat areas of native Hawaiian flora and fauna with the NARS.
U.S. Fish and Wildlife Final Designation and Non-designation of Critical Habitat for 46 Plant Species From the Island of Hawai'i, HI (UFWS 2003)	Provides recommendations for habitat management for <i>Cyrtandra giffardii</i> , <i>Cyanea platyphylla</i> , <i>Clermontia peleana</i> , <i>Clermontia pyrularia</i> , <i>Cyrtandra tintinnabula</i> , and <i>Phyllostegia warshaueri</i>
U.S. Fish and Wildlife Service. Big Island II: Addendum to the Recovery Plan for the Big Island Plant Cluster (USFWS 1998a)	Describes recovery actions needed for endangered plant species: <i>Cyanea platyphylla</i> , <i>Phyllostegia racemosa</i> , and <i>Phyllostegia warshaueri</i>
U.S. Fish and Wildlife Recovery Plan for the Hawaiian Hoary Bat (USFWS 1998b)	Describes recovery actions needed for the Hawaiian Hoary Bat including protecting and managing current populations.
U.S. Fish and Wildlife Recovery Plan for the Big Island Plant Cluster (USFWS 1996)	Describes recovery actions needed for <i>Clermontia lindseyana</i> , <i>Clermontia peleana</i> , <i>Cyrtandra giffardii</i> , and <i>Cyrtandra tintinnabula</i> .
Hawai'i Tropical Forest Recovery Action Plan Hawai'i Tropical Forest Recovery Task Force (1994)	Plan to develop consensus actions needed to recover, manage, and enhance Hawaii's tropical forests
Laupāhoehoe Natural Area Reserve Draft Management Plan (DLNR 1989)	Previous management plan for the NAR

Land Use History

Laupāhoehoe Forest and the surrounding areas have a rich history that has shaped the way the landscape looks today. In 2006, Kumu Pono Associates prepared *Hilo Palikū - Hilo of the Upright Cliffs: A Study of Cultural-Historical Resources of Lands in the Laupāhoehoe Forest Section, Ahupua'a of the Waipunalei-Mauluanui Region, North Hilo District, Island of Hawai'i* (Maly and Maly 2006). This detailed study provides extensive background of the history and cultural resources of the area from ancient Hawaiian uses to the present day and is also discussed further in the Social and Cultural Background portion of this plan. The complete study is available online (see REFERENCES).

In the collection of native and historical accounts, Kumu Pono Associates notes that the lands of the Laupāhoehoe forest region were frequently mentioned in several prominent traditions. Significantly,



the importance of the Laupāhoehoe region koa forests, mountain bird habitats, and the traditional trails which connected the lowlands with the mountain lands and neighboring districts, are frequently referenced in traditions and historical accounts. Also, battles fought on the Laupāhoehoe lands were among those which established the kingdoms of chiefs between the early 1500s to the late 1700s.

In the mid Nineteenth Century the lower elevation portions of fourteen ahupua‘a (traditional land subdivisions encompassing swaths of land running from the coast to the interior mountains) were sought for sugar plantations. By 1876, the Laupāhoehoe Sugar Company and Mill was developed and lowland forests up to about the 2,000-foot elevation were cleared for sugar cultivation, the development of flumes and water resources and homestead lots (Maly and Maly 2006). By the late Twentieth Century sugar production on Hawai‘i was no longer economically competitive with production in other tropical locations and the sugar plantations closed (Maly and Maly 2006).

During the early historic period, the upland section of the Laupāhoehoe forests was impacted by herds of wild sheep and bullocks. By 1825, foreign bullock hunters had established camps on the outer edges of the forest, in the region where Laupāhoehoe and neighboring lands are cut off by the ahupua‘a of Humu‘ula. By the 1850s, the bullock and sheep hunting activities were giving way to formal ranching operations, with the land of Humu‘ula taking in sections of the Laupāhoehoe forests (Maly and Maly 2006).

Timber harvesting has also impacted the landscape. Forests in the area were used by traditional canoe makers and others over many centuries, but the pace of harvesting increased by the middle 1800s when sawmills were established in windward Hawai‘i Island and lumber was harvested and milled for growing island communities and businesses. The negative impacts of logging and grazing on the land were noted by the middle to late 1800s, and government leases began to incorporate conditions meant to conserve forest resources. After establishment of the Hilo FR in 1905, almost no collection of lumber occurred in the Laupāhoehoe section, except for that in direct association with management of the ranch lands. Timber harvest again impacted the area in 1969, when Blair, Inc. received a right of entry permit from DLNR and built a road within Laupāhoehoe FR. A license to harvest timber, primarily koa was awarded to Blair, Inc. in 1971. The Blair logging operation impacted approximately 1,000 acres and was terminated by 1979, when Blair went out of business (Maly and Maly 2006). Research after the logging operation indicated that disturbance from logging stimulated koa regeneration but invasion of banana poka (*Passiflora tarminiana*) was also noted (Scowcroft and Nelson 1976). A later study found impacts of the former logging operation including an increase in invasive tropical ash (*Fraxinus uhdei*), which outcompetes native tree species in disturbed areas (Friday et al. 2008).

In 2006, DOFAW prepared a plan and environmental assessment for reforestation of the timber harvest area (state of Hawai‘i 2006) that included scarification to increase koa (*Acacia koa*) regeneration. DOFAW staff never implemented the project because other management actions including removal of feral cattle and introduction of a biocontrol agent to control the invasive non-native banana poka in the area enhanced natural regeneration of the forest. The biocontrol agent proved successful in reducing banana poka (Trujillo et al. 2001). Native species, particularly koa and ‘ākala (*Rubus hawaiiensis*) are recovering in the former timber harvest area although other invasive non-native plants are still a problem.



Public Access and Recreation

Laupāhoehoe Forest is open to the public for various recreational and cultural uses. While the public is permitted to access and hike or hunt in any portion of the forest, there are limited legal access points and only a few minimally maintained and marked trails. This area is rough and remote rainforest wilderness and there are currently no amenities for recreational users.

Some uses in the NAR, including hiking or nature study with groups larger than ten, research, scientific collecting, gathering (including Native Hawaiian religious and customary gathering rights) and commercial uses require an HETF permit. Gathering within the FR also requires a permit (see Gathering).

Vehicle Access


Mauka and makai public access to Laupāhoehoe Forest is currently via two main access points off Spencer Road and Mana Road respectively (Figure 3). These roads provide vehicle access to approximately the forest boundary:

- Spencer Road Access - The state has an easement through private pasture lands at the top of Spencer Road to allow for public pedestrian access to the lower boundary of Laupāhoehoe Forest. Spencer Rd. is a paved County road passable in a two-wheel drive vehicle. There is a small grassy area where vehicles may park mauka of where Spencer Rd. terminates. A primitive, minimally user-maintained trail provides pedestrian access from the parking area to the forest (see other primitive trails below).
- Mana Road Access - Mana Rd. is a 4-wheel drive County road approximately 40 miles long that roughly transverses a contour along Mauna Kea. Mana Rd. is accessed via Mauna Kea Access Rd. (off Saddle Road) on the Hilo side or from the town of Waimea. The first six miles of the road are regularly maintained at both entrances. The road intersects the top of the Laupāhoehoe Forest. A Nā Ala Hele designated trail, Kaluakauka Trail, is accessed via this route (see description under Trails section below).



Figure 3

Laupāhoehoe Forest Existing Infrastructure



Trails

Trails within Laupāhoehoe Forest include the following:

- Kaluakauka Trail – The trailhead for this Nā Ala Hele designated trail is on the makai side of the Keanakolu-Mana Rd., 17.7 miles from the junction with Mauna Kea access road. The trail is considered moderately difficult and is unmarked and rarely maintained (as described per the official website, <http://hawaii Trails.ehawaii.gov/>). The trail goes downhill across forested pasture land to a foot gate in the FR boundary fence, then continues to the Dr. David Douglas monument erected in 1934 (approximate death site of Dr. David Douglas, the Scottish botanist for whom the Douglas Fir is named).
- Maulua Trail – A portion of this historic ranching-era trail goes across the upper section of Laupāhoehoe Forest from the boundary near Shack Camp to Waipunalei. Access to this portion of the trail is via Blair Rd.
- Other Trails – Additional trails can be found within Laupāhoehoe Forest, notably Peneki and Spencer trails (Figure 3). These trails are not formally recognized as public access trails and are not marked or maintained. These primitive trails were created by the hunting community from the Spencer Road access point. Trail conditions are hazardous, steep and muddy, and lower elevation portions of the trail within the strawberry guava belt may frequently be ‘tunneled’ in by guava tree windfall.

Mountain Biking

Mountain biking is legal on FR roads unless otherwise posted. There are no legal public access trails suitable for mountain biking that access Blair Rd. within the FR. Accessing roads or trails across private lands to reach Blair Rd. without landowner permission is illegal.


Hunting

DOFAW manages public hunting on all state lands and hunting in the Laupāhoehoe Forest is regulated by Chapter 13-122,123, Hawai‘i Administrative Rules (Rules Regulating Game Bird Hunting, Game Mammal Hunting). The Laupāhoehoe Forest includes hunting units B and C in the FR and hunting unit K in the NAR (Figure 4). There is a hunter check station at the Spencer Rd. access. DLNR’s Division of Conservation and Resource Enforcement (DOCARE) carries out enforcement of hunting regulations. Current information regarding hunting rules, seasons and bag limits for all game species can be obtained by contacting the DOFAW Hilo office at 19 East Kawili St. Hilo, Hawai‘i, (808) 974-4221.

All persons are required to have a valid Hawai‘i hunting license on their person to hunt or have a bagged game mammal in their possession. Hunting licenses may be purchased online from <http://www.ehawaii.gov.org/DLNR/hunting/>, from any DOFAW office or from any registered hunting license vendor. All hunting license applicants must show proof of having successfully completed a hunter education course that is recognized by the National Hunter Education Association.



Figure 4



Camping

There are no designated camping areas and no camping is currently allowed in the Laupāhoehoe Forest.

Gathering

Small-scale non-commercial harvesting or salvage is allowed in the FR section of Laupāhoehoe Forest, such as materials for cultural uses. Non-timber forest products such as ferns, maile (*Alyxia stellata*), flowers, fruits, and lei-making materials etc. for cultural or personal use may be collected from within the FR. Gathering of forest products is permitted and regulated by DOFAW through Forest Reserve System permit procedures. Permit applications for gathering plant material can be obtained from the DOFAW Hilo office at 19 East Kawili Ave. Hilo, Hawai‘i, (808) 974-4221. These permits are available, upon approval, free of charge (for common, personal use items) or for a fee, depending on the purpose. Gathering of materials from listed endangered species is not permitted. Gathering (including Native Hawaiian religious and customary gathering rights) within the NAR portion of the forest requires an HETF permit.

Existing Infrastructure and Facilities

Roads

Spencer and Mana roads provide public access to the Laupāhoehoe Forest boundary (see Public Access and Recreation). There is an existing road located within Laupāhoehoe Forest used for management, research and educational purposes, but this road is not currently available for public vehicular access as private lands/roads must be traversed in order to reach the road, as well as liability issues and maintenance costs.

- Blair Road - Blair Rd. requires 4-wheel drive and is contained within Laupāhoehoe Forest (Figure 3). This road is used for management, research and education purposes by the state and the USFS. Public pedestrian use of the road on state lands is allowed for people who access the road by hiking through the forest from public access points (Spencer Rd. or Mana Rd. access). The state has an easement trade with Parker Ranch for use of Blair Rd in exchange for use of Parker Ranch roads to access other portions of the Hilo FR (Humu‘ula section). Historically, Blair Road was built and extended between 1969-1973, and used for logging portions of Laupāhoehoe Forest.

Facilities/Structures

- USFS Facilities - The Laupāhoehoe Science and Education Center, located approximately 4 miles from the Laupāhoehoe Forest boundary, enhances the ability of the HETF to meet its goals for research, education, and demonstration (Figure 3). This facility serves as the primary support facility and includes a classroom, workshop, bunkhouse, and laundry facilities that can accommodate around 30 visiting scientists, educators or students for day-use and 15 visitors overnight. The facilities on this site are under construction and due to be completed in 2015. In addition the Forest Pavilion (Field Education Site) will be located on a 3-acre (1.2 ha) parcel located adjacent to the Laupāhoehoe Forest (Figure 3). It will consist of a covered pavilion, toilet/comfort station, and parking area, and it will be used primarily as a staging area for research and education trips into Laupāhoehoe Forest. The Forest Pavilion facilities are still in planning phase with completion dates TBD.



- Shack Camp – Shack Camp is located at approximately 5,200 ft (1,585 m) elevation near the intersection of the Laupāhoehoe Forest boundary and the historic Maulua Trail. This site contains the ruins of a historic structure associated with Kukaiau Ranch as well as an opening in the forest due to past cattle grazing.



USFS Laupāhoehoe Science and Education Center

Regional Partnerships

DOFAW and USFS are both members of the Mauna Kea Watershed Alliance and the Laupāhoehoe Forest is included in this partnership area. This partnership includes approximately 484,000 ac (195,868 ha) on the mountain of Mauna Kea. The Mauna Kea Watershed Alliance seeks to manage critical watersheds on a landscape-level by initiating planning for priority areas with the goal of implementing management actions for threats such as feral ungulates, fire, and invasive non-native plants. Coordinated management of these watershed lands is critical to sustain adequate quality and quantity of water and provide important habitat for a wide diversity of native plants and animals, including many that are endangered. The Mauna Kea Watershed Alliance is currently working on several projects related to the HETF including sharing of cultural awareness and protocols for IPIF staff, providing review of the HETF Master plan as it is developed, and planning and implementing restoration of the Laupāhoehoe Science and Education Center grounds.

Site Description (Physical and Biological Resources)

Topography, Climate, Geology, and Soils

Laupāhoehoe Forest is located on the eastern, windward flanks of Mauna Kea from about 1,700 to 6,100 ft (518–1860 m) elevation. As the trade winds off the Pacific Ocean strike the mountain, moist air is elevated and cooled, resulting in cloudy weather, high rainfall rates and afternoon fog and mist in the area. Condensation from ground-level clouds (fog drip) contributes additional




moisture at higher elevations. Average annual rainfall in the lower elevations is about 160 inches (in) (418 centimeters (cm)) and ranges from 60 to 100 in (157 to 261 cm) in the upper elevations (HETF Establishment Record 2007).

Temperatures decrease with elevation. At sea level the average monthly day time temperatures range from 79 to 82 °F (26-28 °C) and the night time temperatures range from 62 to 70 °F (17-21 °C). At highest elevations, the temperature could be more than 20 °F (13 °C) colder than in the lowlands.

There is a climate station (maintained by the USFS) at Laupāhoehoe Forest within the FR recording a variety of information including: air temperature, relative humidity, solar radiation, soil moisture and temperature (Figure 3).

Laupāhoehoe Forest is located on Mauna Kea, a dormant volcano and the second oldest volcano on the island. Figure 5 depicts substrate age, which ranges from 5,000 years before the present to 300,000 years before the present (Sherrod et al.



Climate station at Laupāhoehoe Forest

2007). The terrain and soils varies with the age and type of surface lava flows and the depth of volcanic ash deposited over these flows (Wolf and Morris 1996). The terrain in the highest elevation areas is the youngest and the roughest. Surface flows in this area are grouped with the youngest of Mauna Kea's post-shield formation flow series and are characterized as predominantly a'a or blocky a'a flows which are generally free of the wind-blown volcanic ash deposits that cover the older Mauna Kea flows (Wolf and Morris 1996:13; Sheet 2). These younger a'a flows form a series of pronounced ridges that give the upper areas of Laupāhoehoe Forest a distinct ridge and swale topography. Soils on these flows are described as very stony loam (Sato et al. 1973: 15, Sheet 40).

In the upper mid-elevation of Laupāhoehoe Forest, the surface lava flows are older but are still grouped with those erupted during the younger, post-shield phase of Mauna Kea's development (Wolf and Morris 1996: 13; Sheet 2). These flows are also predominantly a'a or blocky a'a flows but are partially mantled by volcanic ash deposits. Soils on these flows are described as silt loam formed from volcanic ash (Sato et. al. 1973: 14 and 50, Sheets 33 and 40). These ash-derived soils are more weathered in the lower elevations where rainfall is slightly greater. Some areas can also be rocky where volcanic ash deposits are discontinuous. Natural Resources Conservation Service Soil Classifications are depicted in Figure 6.



Figure 5






Figure 6

Laupāhoehoe Forest Soil Types



Legend

- | | | | |
|---|---|--|-------------------------|
|  | 'Akaka Soils |  | Streams |
|  | Apakuie Very Fine Sandy Loam, 12-20% slopes |  | Laupāhoehoe Forest/HETF |
|  | Hanipoe Silt Loam, 12-20% slopes | | |
|  | Hanipoe Very Stony Loam, 12-20% slopes | | |
|  | Honoka'a Silty Clay Loam, 10-20% slopes | | |
|  | Kaiwiki Silty Clay Loam, 10-20% slopes | | |
|  | Kaiwiki Silty Clay Loam, 20-35% slopes | | |
|  | Kiloa Extremely Stony Muck, 6-20% slopes | | |
|  | Maile Silt Loam, 6-20% slopes | | |
|  | Pi'ihonua Extremely Stony Silty Clay Loam, 6-20% slopes | | |
|  | Pi'ihonua Silty Clay Loam, 6-20% slopes | | |
|  | Pu'u O'o Silt Loam, 6-12% slopes | | |
|  | Rough Broken Land | | |



Water Resources

The Hilo FR (Laupāhoehoe section) was originally established in 1905 to protect the water supply of the district, and Laupāhoehoe Forest continues to provide important watershed services for the community. Native Hawaiians recognized the importance of forests in water production and water quality, as reflected in the Hawaiian proverb, “Haihai ka ua i ka ulu la au” (The rain follows after the forests). Early foresters also recognized the importance of Hawaiian forests as watersheds. Ralph Hosmer, the first Territorial Forester stated "In Hawai‘i, the most valuable product of the forest is water, rather than wood".

Laupāhoehoe Forest is an important source of fresh surface and ground water that supports downstream populations of humans and wildlife as well as supports healthy nearshore resources. Other watershed services provided by Laupāhoehoe Forest include: provision stream habitat for native waterbirds, fish, and invertebrates, forest habitat for native plants, birds, and bats, flood control, mitigation of climate change impacts, and economic, social, recreational and educational opportunities for the human communities in the area.

Numerous streams are found in the Laupāhoehoe Forest, including Ka‘awali‘i Stream, Laupāhoehoe Stream, Kīlau Stream, Kaiwilahilahi Stream, Ha‘akoa Stream, and Pāhale Stream (Figure 7, Table 2). The Atlas of Hawaiian Watersheds and Aquatic Resources (Parham et al. 2008) notes all these streams as perennial. However, the upper portions of these streams within Laupāhoehoe Forest are often intermittent. While the lack of surface water in these upper reaches makes it appear some of these streams within the forest may not necessarily be flowing year-round, subsurface groundwater flows from the forest maintain freshwater inputs to streams below Laupāhoehoe Forest. Stream gauges, used to measure natural stream flows, water quality and sediment in a non-destructive manner, are located in Manowai‘ōpai, Kaiwilahilahi, and Ka‘awali‘i streams below Laupāhoehoe Forest and are maintained by the USFS.




Trees capture moisture from mist which adds to watershed recharge

Table 2. Streams and Watershed Basins of Laupāhoehoe Forest (Parham et al. 2008)

Watershed Basin Name	Streams	Watershed Basin Name	Streams
Ka‘awali‘i Gulch	Ka‘awali‘i	Ha‘akoa	Ha‘akoa
Laupāhoehoe	Laupāhoehoe	Ka‘alau	Pāhale
Kīlau	Kīlau	Kapehu	Kapehu
Manowai‘ōpai	Manowai‘ōpai	Paeohe	Paeohe
Kaiwilahilahi	Kaiwilahilahi	Maulua	Maulua



Figure 7



Forest Ecosystems

Laupāhoehoe Forest contains native-dominated forested landscapes from lowland forest at 2,300 ft (701 m) above sea level extending to almost 6,500 ft (1,981 m) in elevation. It is part of the largest remaining native dominated forest in Hawai‘i and largely dominated by ‘ōhi‘a (*Metrosideros polymorpha*) and koa (*Acacia koa*), the two most widespread tree species in native forest remaining in Hawai‘i. Laupāhoehoe Forest contains five primary native communities, as well as significant areas between 1,700 ft (518 m) elevation and



‘Ōhi‘a (*Metrosideros polymorpha*) and koa (*Acacia koa*)

~3,000 ft (914 m) of highly altered, non-native dominated vegetation cover (Figure 8). Forestry plantings along the lower boundary and in the lower east corner include non-native trees such as toon (*Toona ciliata*) and *Ficus rubiginosa*, and in the upper north corner, tropical white ash (*Fraxinus uhdei*). The tropical ash has invaded significant portions of higher elevation areas of the Laupāhoehoe Forest. Other non-native species occupy large areas. Banana poka (*Passiflora tarminiana*), an introduced vine, occurs throughout mid to high elevation areas and forms thickets in the swales. At the lower edge of this community type, below 3,000 ft (914 m) elevation, the understory is heavily invaded by several non-native plants including strawberry guava (*Psidium cattleianum*), thimbleberry (*Rubus rosifolius*), Koster’s curse (*Clidemia hirta*), Himalayan ginger (*Hedychium gardnerianum*), various grasses, and three species of parasitic strangler fig. A number of these species occur into mid or even high elevation areas of the Laupāhoehoe Forest. Non-native grasses and herbs are primarily pasture species (e.g., kikuyu grass, *Holcus lanatus*, and *Ehrharta stipoides*) and the vine, German ivy (*Delairea odorata*).

Native Plant Communities

a. Koa/‘Ōhi‘a Lowland Wet Forest

This forest type dominates in the lowest elevation area up to about 3,000 ft (914 m) elevation, where the Montane Wet Forest takes dominance. Lower elevation portions of this forest type are badly invaded by invasive, non-native species. In addition, some mixed non-native tree plantings occur at the lower boundary of the area in the east corner. Under the 80 ft (24 m) tall closed to open canopy of koa and ‘ōhi‘a, is a secondary tree layer in which olomea (*Perrottetia sandwicensis*), mehame (*Antidesma platyphyllum*), ālani (*Melicope clusiifolia*), and kōpiko



Olomea, kanawao, and mehame (from left to right)



(*Psychotria hawaiiensis*) are common. Other trees, such as ‘ōlapa (*Cheirodendron trigynum*) and kāwa‘u (*Ilex anomala*) are present, but not as common.

Hāpu‘u (*Cibotium glaucum*) is present, but of lower stature than in the Koa/‘Ōhi‘a Montane Wet Forest, and forms a discontinuous layer. Common shrubs include manono (*Hedyotis terminalis*), kanawao (*Broussaisia arguta*), ‘ōhelo (*Vaccinium calycinum*), and saplings of kāwa‘u and ‘ōlapa. The vines ‘ie‘ie (*Freycinetia arborea*) and maile (*Alyxia stellata*) are present, and ‘ie‘ie is sometimes abundant. Native ferns include wahine noho mauna (*Adenophorus* spp.), *Lycopodium cernuum*, *Athyrium* spp., *Elaphoglossum* spp., *Sphenomeris chinensis*, and others.

Rare plants observed in this forest type in surveys in the 1980’s include *Cyrtandra giffardii*, *Cyanea tritomantha*, *Gardenia remyi* and *Platydesma remyi*.

b. Koa/‘Ōhi‘a Montane Wet Forest

This montane wet forest distributes in areas from about 3,000 ft (914 m) elevation up to 4,500 ft (1372 m) elevation. It differs from the Koa/‘Ōhi‘a Lowland Wet Forest on its subcanopy species composition. Koa and ‘ōhi‘a form an open to closed canopy (about 100 feet (30 m) in height) with a very well-developed subcanopy of tree ferns (*Cibotium glaucum*, *C. chamissoi*, and *C. hawaiiense*). Trees in the secondary tree layer include ‘ōlapa, kāwa‘u, kōlea (*Myrsine lessertiana*), and pilo (*Coprosma rhynchocarpa* and *C. pubens*).

In the understory, native shrubs include ‘ōhelo, ‘ākala (*Rubus hawaiiensis*), *Cyrtandra* spp., *Clermontia parviflora*, māmaki (*Pipturus albidus*), manono, and saplings of ‘ōlapa, ‘ōhi‘a, pilo, and kāwa‘u. Ferns are often the prevalent ground cover, including *Asplenium* spp., *Dryopteris wallichiana*, ‘ākōlea (*Athyrium microphyllum*), *Ophioglossum pendulum* subsp. *falcatum*, and *Lepisorus thunbergianus*. The rare mint, *Stenogyne macrantha*, is known from the area between Kaiwilahilahi Stream and the NAR's western boundary.



‘Ōhelo, māmaki and ‘ākolea (from left to right)

c. ‘Ōhi‘a/Hāpu‘u/Uluhe Montane Wet Forest

This vegetation cover type occurs on the east side between 3,500 and 4,500 ft (1067-1372 m) elevation, almost bisecting the upper area of Koa/‘Ōhi‘a Montane Wet Forest. A tall (approximately 80 ft (24 m)) open to scattered canopy of ‘ōhi‘a with a secondary layer of native trees such as olomea, mehame, ‘ōlapa, and pilo and hāpu‘u grow over a layer composed largely of uluhe fern (*Dicranopteris linearis*). Under the hāpu‘u, there is a mix of native shrubs, such as manono, young ‘ōlapa, pilo, *Cyrtandra* spp., and *Clermontia parviflora*. Hō‘i‘o (*Athyrium*



sandwichianum) is the most abundant native fern, although *Asplenium* spp., *Vandenboschia davallioides*, wahine noho mauna, *Elaphoglossum* spp., and *Lepisorus thunbergianus* are also present.

d. *Carex alligata* Montane Wet Grassland

Much of the mid elevation area in Laupāhoehoe Forest, between about 4,000 and 4,500 ft (1220-1370 m), is poorly drained, and several low-lying, very wet sections are dominated by *Carex alligata*. Species from the surrounding natural communities, such as scattered ‘ōhi‘a, ‘ōlapa, and ‘ōhelo, are also found in this community type.

e. Koa/‘Ōhi‘a Montane Forest

This forest type has been significantly altered by past land uses, including ranching and logging, and has also been heavily impacted by feral cattle. Compared to the Montane Wet Forest, the Montane Forest receives less rainfall; the annual rainfall is about 39.3-74.7 in (100-190 cm). The forest consists of scattered-to-open uneven canopy of 115 ft (35 m) tall koa emergent above 82 ft (25 m) tall ‘ōhi‘a . The tall-stature trees tend to grow along the ridge formations. Swales between the ridges and open areas are dominated primarily by thick patches of ‘ākala (*Rubus hawaiiensis*).


The understory has many species in common with Koa/‘Ōhi‘a Montane Wet Forest, but the distinct hāpu‘u tree fern layer of the latter is absent. Species more characteristic of drier areas may also be components here. Ground cover is often dominated by native ferns, especially laukahi (*Dryopteris wallichiana*). Species found in this forest type include: ‘ōlapa, pilo, manono, kāwa‘u, *Myoporum sandwicense*, kōlea, alani, *Ranunculus hawaiiensis*, *Sophora chrysophylla*, *Styphelia tameiameia* and ‘ōhelo.



‘Ākala fruits and flower



Figure 8



Rare Plants

A diversity of native plants, including rare species are found within Laupāhoehoe Forest, and there is critical habitat for six species (Table 3). Table 3 also includes species that may not have been found within Laupāhoehoe Forest, but are known from similar habitat in nearby areas. The U.S. Endangered Species Act defines Critical Habitat as areas that may or may not be occupied by a threatened or endangered species, but are essential to the conservation of the species. These areas may require special management considerations or protection (16 U.S.C. § 1532 (5)).



Rare plants include 'ohe, jewel orchid and 'oha wai (clockwise from top left)



Table 3. Rare Plants with habitat in or near Laupāhoehoe Forest

<i>Species</i>	<i>Common Name</i>	<i>Federal Status*</i>	<i>Critical Habitat</i>	<i>Known From Laupāhoehoe Forest**</i>	<i>Known From Adjacent Areas</i>	<i>Outplanted in Laupāhoehoe Forest</i>
<i>Anoectochilus sandvicensis</i>	<i>jewel orchid</i>	<i>SOC</i>		<i>X</i>		
<i>Asplenium schizophyllum</i>	-	<i>SOC</i>		<i>X</i>		
<i>Clermontia lindseyana</i>	<i>'oha wai</i>	<i>LE</i>		<i>X</i>	<i>X (Piha/Hakalau)</i>	<i>X</i>
<i>Clermontia peleana</i>	<i>'oha wai</i>	<i>LE</i>	<i>X</i>		<i>X (Upper Hilo FR)</i>	<i>X</i>
<i>Clermontia pyrularia</i>	<i>'oha wai</i>	<i>LE</i>	<i>X</i>	<i>X (Historic)</i>		<i>X</i>
<i>Cyanea fernaldii</i>		<i>No Status***</i>		<i>X</i>		
<i>Cyanea platyphylla</i>	<i>'akū 'akū</i>	<i>LE</i>	<i>X</i>	<i>X</i>		
<i>Cyanea shipmanii</i>	<i>hāhā</i>	<i>LE</i>			<i>X</i>	<i>X</i>
<i>Cyanea tritomantha</i>	<i>'akū</i>	<i>LE</i>		<i>X</i>		<i>X</i>
<i>Cyrtandra giffardii</i>	<i>ha 'iwale</i>	<i>LE</i>	<i>X</i>	<i>X</i>		
<i>Cyrtandra tintinnabula</i>	<i>ha 'iwale</i>	<i>LE</i>	<i>X</i>	<i>X</i>	<i>X (Hakalau)</i>	
<i>Cyrtandra wagneri</i>	<i>ha 'iwale</i>	<i>LE</i>		<i>X</i>		
<i>Deparia kaalaana</i>	-	<i>SOC</i>		<i>?</i>		
<i>Gardenia remyi</i>	<i>nānū</i>	<i>C</i>		<i>X</i>		
<i>Huperzia manni</i>	-	<i>LE</i>		<i>X</i>		
<i>Huperzia stemmermannii</i>	-	<i>C</i>		<i>X</i>		
<i>Joinvillea ascendens</i>	<i>'ohe</i>	<i>C</i>		<i>X</i>		
<i>Melicope zahlbruckneri</i>	<i>alani</i>	<i>LE</i>		<i>X (Historic)</i>		<i>X</i>
<i>Ochrosia haleakalae</i>	<i>hōlei</i>	<i>C</i>		<i>X</i>		
<i>Phyllostegia brevidens</i>	-	<i>SOC</i>		<i>X (Historic)</i>		
<i>Phyllostegia floribunda</i>	-	<i>LE</i>		<i>X (Historic)</i>		<i>X</i>
<i>Phyllostegia macrophyllum</i>	-	<i>SOC</i>		<i>X</i>		
<i>Phyllostegia racemosa</i>	<i>kīponapona</i>	<i>LE</i>			<i>X (Hakalau)</i>	
<i>Phyllostegia warshaueri</i>	-	<i>LE</i>	<i>X</i>	<i>X</i>		<i>X</i>
<i>Platydesma remyi</i>	-	<i>LE</i>		<i>X</i>		<i>X</i>
<i>Pritchardia lanigera</i>	<i>loulu</i>	<i>LE</i>		<i>X</i>		
<i>Ranunculus hawaiiensis</i>	<i>makou</i>	<i>C</i>			<i>X (Mauna Kea FR)</i>	
<i>Stenogyne macrantha</i>	<i>mā 'ohi 'ohi</i>	<i>SOC</i>		<i>X</i>		
<i>Strongylodon ruber</i>	<i>nuku 'i 'iwi</i>	<i>C</i>		<i>X historic</i>	<i>X</i>	
<i>Trematolobelia grandifolia</i>	<i>koli 'i</i>	<i>SOC</i>		<i>X</i>		<i>X</i>

* Key to Federal Status: Listed Endangered (LE) = Taxa listed as endangered. Candidate (C) = Taxa for which substantial information on biological vulnerability and threat(s) support proposal to LE. Species of Concern (SOC) = Taxa for which available information meets the criteria for concern and the possibility to recommend C.

** Current (wild plants currently present), Historic (historic records describe presence in area); ***newly described or resurrected species



Wildlife

Birds


Laupāhoehoe Forest was surveyed for forest birds as part of the Hamakua Study Area during the Hawai‘i Forest Bird Survey (1976-1983). This survey and several subsequent surveys of the area have provided information on the bird species present and their distribution.

The forest provides habitat for six honeycreepers (Subfamily Drepanidinae) endemic to the Hawaiian Islands. These include three endangered species: Hawai‘i ‘ākepa (*Loxops coccineus*), Hawai‘i creeper (*Oreomystis mana*) and ‘akiapōlā‘au (*Hemignathus munroi*). The non-endangered honeycreepers found in the project area include: ‘apapane (*Himatione sanguinea*), Hawai‘i ‘amakihi (*Hemignathus virens*), and ‘i‘iwi (*Vestiaria coccinea*). U.S. Fish and Wildlife Service (USFWS) is currently reviewing the status of ‘i‘iwi to determine whether it should be listed as endangered or threatened. Other native forest birds reported from the project area include, ‘elepaio (*Chasiempis sandwichensis*), and ‘ōma‘o or Hawaiian thrush (*Myadestes obscurus*). Native forest birds are primarily found in the upper elevations (above 4,000 ft (1,219 m)) where lower numbers of mosquitoes and the effects of cooler temperatures on plasmodium parasite reduce the incidence of diseases such as avian malaria and pox. It is important to note that Hakalau Forest National Wildlife Refuge is adjacent to Laupāhoehoe Forest where these species are seen regularly along with many other native species.

Other native bird species listed as endangered by the USFWS, have been reported from the Laupāhoehoe Forest area including the Hawaiian duck or koloa maoli (*Anas wyvilliana*), and the Hawaiian hawk or ‘io (*Buteo solitarius*). Koloa maoli are generally found in a wide variety of natural and artificial wetland habitats including freshwater marshes, flooded grasslands, streams, montane pools, irrigation ditches, reservoirs, etc.

Laupāhoehoe Forest is considered a recovery area for Hawai‘i creeper, Hawai‘i ‘ākepa, and ‘akiapōlā‘au in the USFWS forest bird recovery plan and in the State Comprehensive Wildlife Strategy. Recovery areas are habitat that will allow for the long-term survival and recovery of endangered Hawaiian forest birds.

The most widespread non-native birds observed at Laupāhoehoe Forest include hwamei (*Garrulax canorus*), Japanese white-eye (*Zosterops japonicus*), red-billed leiothrix (*Leiothrix lutea*), northern cardinal (*Cardinalis cardinalis*) and kalij pheasant (*Lophura leucomelana*). Other non-native birds observed in the area are summarized in Table 4 below.



‘Akiapōlā‘au and ‘i‘iwi (courtesy of Jack Jeffrey Photography)



Table 4. Laupāhoehoe Forest bird species

Species	Common Name	Status
<i>Acridotheres tristis</i>	Common myna	Non-native
<i>Alauda arvensis</i>	Eurasian skylark	Non-native
<i>Anas wyvilliana</i>	Hawaiian duck, koloa maoli	Endemic (Endangered)
<i>Asio flammeus sanwicensis</i>	Short-eared owl, pueo	Endemic
<i>Buteo solitarius</i>	Hawaiian hawk, 'io	Endemic (Endangered)
<i>Cardinalis cardinalis</i>	Northern cardinal	Non-native
<i>Carpodacus mexicanus</i>	House finch	Non-native
<i>Cettia diphone</i>	Japanese bush warbler	Non-native
<i>Chasiempis sanwicensis sandwichensis</i>	'Elepaio	Endemic
<i>Francolinus erckelli</i>	Erkel's francolin	Non-native
<i>Garrulax canorus</i>	Hwamei	Non-native
<i>Hemignathus munroi</i>	'Akiapola'au	Endemic (Endangered)
<i>Hemignathus virens virens</i>	'Amakihi	Endemic
<i>Himatione sanguinea sanguinea</i>	'Apapane	Endemic
<i>Leiothrix lutea</i>	Red-billed leiothrix	Non-native
<i>Lonchura punctulata</i>	Nutmeg mannikin	Non-native
<i>Lophura leucomelana</i>	Kalij pheasant	Non-native
<i>Loxops coccineus coccineus</i>	Hawai'i 'ākepa	Endemic (Endangered)
<i>Meleagris gallopavo</i>	Wild turkey	Non-native
<i>Myadestes obscurus</i>	Hawai'i thrush, 'oma'o	Endemic
<i>Nycticorax nycticorax hoactli</i>	Auku'u or black crowned night heron	Indigenous
<i>Oreomystis mana</i>	Hawai'i creeper	Endemic (Endangered)
<i>Passer domesticus</i>	House sparrow	Non-native
<i>Phasianus colchicus</i>	Ring-necked pheasant	Non-native
<i>Pluvialis fulva</i>	Kolea or pacific golden plover	Indigenous
<i>Serinus mozambicus</i>	Yellow fronted canary	Non-native
<i>Streptopelia chinensis</i>	Spotted dove	Non-native
<i>Tyto alba</i>	Barn owl	Non-native
<i>Vestiaria coccinea</i>	'I'iwi	Endemic
<i>Zosterops japonicus</i>	Japanese white-eye	Non-native

Mammals

Laupāhoehoe Forest is considered very important habitat for Hawai'i's only native land mammal, the 'ōpe'ape'a – the endemic and endangered Hawaiian hoary bat (*Lasiurus cinereus semotus*), which uses the area for roosting, reproduction and foraging. U.S. Geological Survey (USGS) Biological Resources Division Hawaiian Hoary Bat Project has monitored bats for five years and has found high levels of bat activity and occupancy. The Hawaiian hoary bat is the



only native terrestrial mammal from the Hawaiian archipelago (USFWS 1998). It is a medium-sized, nocturnal, insectivorous bat with short, thick, rounded ears and a furry tail. "Hoary" refers to the white-tinged, frosty appearance of the bat's grayish brown or reddish brown fur.

A variety of non-native mammals such as feral pigs (*Sus scrofa*), rats (*Rattus* spp.), mice (*Mus musculus*), cats (*Felis catus*), wild dogs (*Canis lupus familiaris*) and mongoose (*Herpestes auropunctatus*) are present in Laupāhoehoe Forest.



Photo by: Corinna A. Pinzari

'Ōpe'ape'a.

Invertebrates

Native invertebrates known from Laupāhoehoe Forest include numerous species of *Drosophila*. Species including *D. sproati*, *D. murphyi*, *D. tanythrix*, and *D. yooni* are relatively common. One notable collection was of a female specimen believed to be *D. papala*, taken at 4,800 ft (1,463 m) elevation. In 2012, a researcher cataloging *Drosophila* species noted a small patch of *Pisonia brunoniana* at about 4,000 ft (1,219 m) elevation within the FR as the most diverse site sampled. Although too small to support any picture-wing species, this disjunct grove of mesic trees has a community of smaller *Drosophila* species not found elsewhere in Laupāhoehoe Forest. Some of these are associated with *Pisonia* in particular (*D. kambysellisi*, *D. nr. dissita*), while others are associated with other plants but seem to be attracted to the site. This spot is worthy of greater conservation attention, especially since *Pisonia* is relatively rare in the Hamakua area. Table 5 provides a list of *Drosophila* species found during the 2012 survey. Several picture-wing species that breed in *Charpentiera*, *Pisonia*, and *Urera* were formerly known from lower elevations (~2,500 ft (762 m)), but this area now appears to be largely non-native, and no flies were found there.

Laupāhoehoe Forest also contains habitat for four endemic species of pinao or Hawaiian damselfly. *Megalagrion calliphya* and *Megalagrion hawaiiense* breed in small pools or seeps in the forest, whereas *Megalagrion blackburni* breeds in streams. *Megalagrion xanthomelas* is a candidate for listing as an endangered species and is known from Kaiwilahilahi Stream (Parham et al. 2008), below the lower boundary of Laupāhoehoe Forest.

Table 5. Laupāhoehoe Forest *Drosophila* species found during 2012 Survey

<i>Drosophila basisetae</i>	<i>Drosophila murphyi</i>	<i>Drosophila sordidapex</i>
<i>Drosophila brunneicrus</i>	<i>Drosophila neutralis</i>	<i>Drosophila sproati</i>
<i>Drosophila canipolita</i>	<i>Drosophila nr. ancyla</i>	<i>Drosophila tanythrix</i>
<i>Drosophila cnecopleura</i>	<i>Drosophila nr. dissita</i>	<i>Drosophila tendomentum</i>
<i>Drosophila cracens</i>	<i>Drosophila nr. medialis</i> #2	<i>Drosophila trichaetosa</i>
<i>Drosophila dasyncemia</i>	<i>Drosophila nr. medialis</i> #3	<i>Drosophila yooni</i>
<i>Drosophila hawaiiensis</i>	<i>Drosophila papala</i>	<i>Scaptomyza (Elmomyza) tumidula</i>
<i>Drosophila imparisetae</i>	<i>Drosophila percnosoma</i>	
<i>Drosophila kambysellisi</i>	<i>Drosophila propiofacies</i>	
<i>Drosophila kikalaeleele</i>	<i>Drosophila seclusa</i>	
<i>Drosophila latigena</i>	<i>Drosophila setosimentum</i>	
<i>Drosophila medialis</i>	<i>Drosophila silvestris</i>	



Aquatic Species

Streams provide habitat for endemic waterbirds, four gobies, two crustaceans, one snail, and several aquatic insects (e.g., damselflies, chironomids) that are noted in the Hawai‘i Stream Atlas (Parham et al. 2008). There are also two species of invasive amphibians that have been observed in or near streams in Laupāhoehoe, *Rana catesbeiana* (American bullfrog) and *Rhinella marina* (Cane toad). Both of these species lay eggs in water and have a tadpole stage to their lifecycle.

Surface and groundwater that originate from the Laupāhoehoe Forest also support healthy populations of native nearshore fish assemblages. Many of these fish are an important component of the subsistence-based economy in the region.

Table 6. Native aquatic species known from at least a portion of the 10 Laupāhoehoe streams (Parham et al. 2008)

<u>Native Fish</u> <i>Awaous guamensis</i> <i>Lentipes concolor</i> <i>Sicyopterus stimpsoni</i> <i>Eleotris sandwicensis</i> <i>Kuhlia xenura</i>	<u>Native Aquatic Insects</u> <i>Megalagrion blackburni</i> <i>Megalagrion xanthomelas</i> <i>Telmatogeton sp.</i>	<u>Native waterbirds</u> <i>Anas wyvilliana</i>
<u>Native Crustaceans</u> <i>Atyoida bisulcata</i>	<u>Native Snails</u> <i>Neritina granosa</i>	

Cultural Resources and Traditional Practices

The 2006 cultural historical study prepared by Kumu Pono Associates is an important reference for cultural resources management in the Laupāhoehoe Forest (Maly and Maly 2006). It references the ethnographical and historic uses of the region, and identifies several historic, ethnographic, and archaeological site types and features that may be found in the area. Excerpts from this study are included below, and the complete study is available online (see REFERENCES).

“The lands of the Laupāhoehoe Forest are part of an ancient region, traditionally known to the Hawaiians as the wao akua (region of the gods), wao ma‘ukele (wet forest zone) and wao nahele (forest zone). In traditional times—pre-western contact in 1778, and in subsequent years through the early 1800s—these forested regions, particularly the wao akua, were considered sacred, the abode of the gods. Travel through the forest lands, undertaking collection of resources—gathering woods and other plant materials, collection of feathers and catching birds, and even travel through the forests, simply to reach another destination beyond the forest—was undertaken with prayer, caution, and respect. Damage to the living forests was often punished by acts of nature—heavy rains might wash the careless traveler from the path; dense mists or sudden growth of such plants as uluhe or ‘ōpiko, might cause the trail to be lost from view, and the traveler to wander aimlessly through the forests.

In this collection of native and historical accounts we also find that the lands of the Laupāhoehoe forest region are frequently mentioned in several prominent traditions.



Significantly, the importance of the Laupāhoehoe region koa forests, mountain bird habitats, and the traditional trails which connected the lowlands with the mountain lands and neighboring districts, are frequently referenced in traditions and historical accounts. Also, battles fought on the Laupāhoehoe lands were among those which established the kingdoms of chiefs between the early 1500s to the late 1700s. While many of the accounts cited in the study relate to the lower lands of the Laupāhoehoe vicinity—those lands situated below the 2,000 foot elevation—there are occasional references to travel through the upland forests to the mountain region. There are also specific references to the traditional significance of Laupāhoehoe, and the occurrence of numerous heiau (ceremonial sites) of local and regional significance, though the record is seemingly silent on the location of heiau that might have occurred in the forest region.”

Archaeological and Historic Sites

The 2006 cultural historical study prepared by Kumu Pono Associates also identifies types of archaeological sites that might be found in the Laupāhoehoe Forest (Maly and Maly 2006).

“These include, but are not limited to—trails extending from the shore to the mountain lands; shelters and resting places along trail sides; shrines used by travelers, bird catchers, canoe makers and other practitioners; battle sites and hiding places; and possible burial sites. Traditional features would include several forms, ranging from stone platforms, terraces, cairns, and walls; and shelter features—called pāpa‘i by the ancient Hawaiians—generally made of wood, leafy branches and ferns. Many of the features would naturally deteriorate and evidence of them would return to the earth. Other features of stone might still be visible in the understory, though only found upon careful search. Another feature of importance would be stone filled fractures or crevices, and caves. Such features were sometimes used for shelters over generations, or as burial sites, and as places in which to hide valued cultural artifacts.”

The historic resources study prepared for the aforementioned 2006 reforestation project sampled transects in the proposed project area of potential effect (and found no historic properties (e.g. no stone surface features, potential shelter caves or overhangs, subsurface cultural deposits) in the Laupāhoehoe Forest portion of the proposed project (Carpenter et al. 2006).

Historical sites identified in the region include the following (Carpenter et al. 2006, Maly and Maly 2006):

- The sheep ranch station at Keanakolu (in the original place of that name, near the Laupāhoehoe-Humu‘ula boundary). There remain on the land in the present-day, the ruins of stone shelters, pens, and foundations.
- Noted places such as Keanakolu (not the same location of the present-day cabin of that name), Lahohinu, and Keahua-ai (Douglas Pit), are considered significant features of the historical landscape.
- Laupāhoehoe-Waipunalei Trail
- Maulua Trail, established as an old pack trail.



THREATS TO NATURAL AND CULTURAL RESOURCES

Non-Native Plants and Animals

Invasive Non-Native Plants

Invasive non-native plants constitute a severe threat to the native ecosystems in Laupāhoehoe Forest. Certain non-native plants are considered invasive because they can establish and survive in undisturbed native forest, disperse long distances via wind or birds, affect large portions of land, displace native vegetation, grow and reproduce rapidly, and convert a diverse native forest plants to a monoculture of alien species. Invasive non-native plants can displace distinctive native flora, resulting in a loss of species diversity and eventually in changes to ecosystem function such as nutrient cycling. Many invasive non-native plants completely replace native vegetation by preventing any regeneration of native species or in the case of strangler figs – direct replacement of native trees, resulting in total loss of native habitats thereby negatively affecting native birds and invertebrates (Cuddihy and Stone 1990; Vitousek 1992). Invasive species can also encourage fire by increasing the amount of available fuels or decrease water input to streams and ground water. For example, forests severely invaded by invasive non-native plants such as strawberry guava show increased evaporation of water to the atmosphere, which reduces the amount of water available for human use (Giambelluca, unpublished research).

Only a small portion of Laupāhoehoe Forest has had systematic surveys for invasive non-native plants. In general, upper forested areas between 3,500 - 4,500 ft (1,067-1,372 m) elevation contain relatively low densities of invasive non-native plants described below. However, below 3,500 ft (1,067 m) forests become heavily invaded by strawberry guava, clidemia or Koster's curse, yellow Himalayan raspberry, and kahili ginger. Above 4,500 ft (1,372 m) forests are heavily invaded by grasses, banana poka, and tropical ash. Throughout the entire forest Australian tree fern (*Sphaeropteris cooperi*) can be found at low densities, although populations are increasing. *Ficus* spp. is concentrated in the northern portion of the forest, near Blair Rd., at about 2,800 ft (853 m) and is spreading into adjacent areas.

Invasive non-native plants with great potential for spreading and causing habitat modification are identified in this plan as high priority for control. Invasive non-native plant species were prioritized based on observed invasiveness and other criteria including growth form, dispersal mechanisms, ability to displace native vegetation and ability to alter ecosystem cycles (water, nutrients and succession). High priority invasive non-native plants currently present in Laupāhoehoe Forest include:

- Australian tree fern (*Sphaeropteris cooperi*)
- Banana poka (*Passiflora tarminiana*)
- Florida blackberry (*Rubus argutus*)
- Cane tibouchina (*Tibouchina herbacea*)
- Clidemia or Koster's curse (*Clidemia hirta*)
- *Ficus* spp.
- Himalayan ginger (*Hedychium gardnerianum*)
- Mules foot fern (*Angiopteris evecta*)
- Palm grass (*Setaria palmifolia*)
- Passion fruit (*Passiflora edulis*)



- *Polygonum chinensis*
- Strawberry guava (*Psidium cattleianum*)
- Tropical ash (*Fraxinus uhdei*)
- Yellow Himalayan raspberry (*Rubus ellipticus*)
- German ivy (*Delairea odorata*)

There are additional invasive non-native plants species of serious concern to land managers that are present in adjoining areas but have not yet been detected in Laupāhoehoe Forest. It is a high priority to prevent the establishment of these species, which include but are not limited to miconia (*Miconia calvescens*), faya (*Morella faya*), gorse (*Ulex europaeus*), firethorn (*Pyrocantha angustifolia*), *Cotoneaster pannosus*, and New Zealand flax (*Phormium tenax*). Other invasive non-native plant species may be added to the priority invasive non-native plant list if monitoring shows their range and abundance increasing in native ecosystems targeted for management.

Ungulates

Ungulates are hoofed animals such as pigs, sheep, goats and cattle. The primary ungulate in Laupāhoehoe Forest is the wild pig (*Sus scrofa*), which is found throughout the forest except in the small fenced areas (35 total acres) that currently exist. Feral ungulates can pose a threat to native ecosystems, species and watersheds because they eat and trample native plants (Cooray and Mueller-Dombois 1981), and disperse the seed of invasive non-native plants. These changes can cause increased erosion and soil runoff. Hawaiian plants evolved without such animals and so some have lost defenses such as thorns and chemical compounds in leaves. The rooting and wallowing behavior of pigs can increase the inputs of pollutants to streams (i.e. animal waste), stream water turbidity due to soil erosion (Stone 1985, Dunkel 2009, Bruland et al. 2010), and wallows can result in breeding areas for disease carrying mosquitoes (Baker 1979, USGS 2005, USGS 2006c). Pigs also eat some invasive plant fruits, such as strawberry guava, which they can transport and then defecate in new areas (Aplet et al. 1991). In addition, feral pigs have been shown to spread root-rot fungi (Baker 1979), and can carry parasites and diseases transmittable to humans and dogs, such as leptospirosis (Warner 1959-1969, Sasaki et al. 1993) and tuberculosis (Giffin 1978).

Pigs were originally brought to Hawai‘i by the first Polynesian settlers as a domesticated species (Tomich 1986). After the arrival of Captain Cook, the larger European wild boar was introduced as a game species and quickly became feral. Today, feral pigs in Hawai‘i are generally smaller in size than mainland varieties as a result of over 200 years of interbreeding between the smaller Polynesian pig and the larger European boar (Tomich 1986).

Feral cattle have been a problem in Laupāhoehoe Forest in the past; however, all feral cattle were removed from the area by 2003. Feral cattle are still a potential future threat as they are still present in adjoining areas (Waipunalei and Humu‘ula) and occasionally get into the forest.

Other Non-Native Animals

A variety of non-native mammalian predators are serious pests to the biodiversity found in Laupāhoehoe Forest. Mongoose, feral cats, rats, but also mice prey upon native species and have a severe impact on native birds. In addition, small mammals serve as vectors of diseases and can



affect the water quality and cause human and wildlife diseases. Leptospirosis and cryptosporidiosis are potentially fatal illnesses caused by water-borne microorganisms spread by non-native mammals (Sasaki et al. 1993).

Feral cats kill forest birds as well as native sea birds and other species that nest on the ground or in burrows (USGS 2006a). Cats are the host of a potentially fatal disease called toxoplasmosis. In Hawai‘i, toxoplasmosis has killed native Hawaiian birds such as the ‘alalā, the endangered nēnē and even seabirds such as the red-footed booby (*Sula sula*). Because the organism that causes toxoplasmosis (*Toxoplasma gondii*) can complete an important part of its life cycle in seawater, this disease also poses a threat to marine mammals such as the endangered Hawaiian monk seal (*Monachus schauinslandi*) and spinner dolphin (*Stenella longirostris*). In addition to threatening wildlife, toxoplasmosis poses a significant health risk to pregnant women (USGS 2006a).

Rats prey on native bird eggs, nestlings, native land snails and also eat the fruits/seeds and strip the bark of native plants. Similarly, mice consume the seeds of native plants; seed predation can be a major factor contributing to species decline (Atkinson 1985, Cole et al. 2000, Cuddihy and Stone 1990, Shiels and Drake 2011)

Laupāhoehoe Forest has been invaded by non-native forest birds; however their impacts on native species have not been determined. The non-native kalij pheasant occur in high densities through the forest and can disturb soils, as well as transport non-native seeds. Non-native birds may compete with native forest birds for food and other resources and act as vectors for avian diseases. Non-native birds also contribute to the spread of invasive non-native plants by eating the fruits and spreading seeds – especially strangler figs, clidemia and ginger.

Non-native invertebrates are present, but largely undocumented, and can consume native plants, interfere with plant reproduction, predate or act as parasites on native species, transmit disease, affect food availability for native birds, and disrupt ecosystem processes. The invasion of the yellow jacket wasp (*Vespula pennsylvanica*), voracious predators of numerous species of native invertebrates, is of concern. Other non-native parasitoids adversely impact native moth species, and ants are a significant mortality factor for native invertebrates. Slugs (*Milax gagates*, *Limax maximus*, and *Veronicella* spp.) consume fruit from native plants and prey on seedlings and mature plants. Black-twig borer (Coleoptera: Scolytidae) is a threat to koa. The two-spotted leafhopper (*Sophonia rufofascia*) is a major concern for the uluhe fern, which is particularly sensitive to leafhopper feeding. Mosquitoes (*Aedes albopictus* and *Culex quinquefasciatus*) transmit deadly diseases to native birds and humans. Little fire ant (*Wasmannia auropunctata*) is known from the community below Laupāhoehoe Forest and could pose a threat to forest resources if it spreads mauka.

Both Jackson’s chameleon (*Chamaeleo jacksonii*) and coqui frog (*Eleutherodactylus coqui*) have growing populations on the island, and these species can consume native invertebrates, such as insects, spiders, and small snails. Coqui frogs are known from pastures below Laupāhoehoe Forest as well as from streams within the town of Laupāhoehoe. Cane toads and American bullfrogs, have also been observed in or near streams in Laupāhoehoe Forest. However, it is unclear how these species might be impacting native ecosystems in the forest.



Disease

Introduced diseases and pathogens threaten native animals and plants. Given the lack of biosecurity in Hawai‘i, the introduction of new diseases and pathogens is highly likely. Avian pox and avian malaria are mosquito-transmitted diseases that currently kill or weaken many native Hawaiian birds and are thought to be responsible for the extinction of numerous forest bird species. In the extreme isolation of the Hawaiian Islands, birds evolved in the absence of these diseases and lost their natural immunity. Avian pox is caused by a virus (Avipoxvirus) and avian malaria by a single-celled parasite (*Plasmodium relictum*). For many native forest bird species, infection with these diseases is almost always fatal (USGS 2005, USGS 2006c).

Introduced plant diseases such as ‘ōhi‘a rust (*Puccinia psidii*) and koa wilt (caused by the fungus *Fusarium* sp.) could potentially impact the most common native trees within Laupāhoehoe Forest. ‘Ōhi‘a rust affects ‘ōhi‘a as well as other plants in the same family (Myrtaceae) (HEAR 2010). In severe infections, growing tips wither and die back. Koa wilt is a serious, often fatal disease of the native tree, koa. Trees affected with the disease rapidly lose their canopies and may die within a few months (UH-CTAR 2010).

Climate Change and Natural Disturbances

Climate change may affect the Laupāhoehoe Forest by altering rainfall patterns and amounts. Changing climate may affect the abundance and seasonality of precipitation, thereby altering forest composition, growth and structure (Iwashita et al. 2013). Rare ecosystems and species may be negatively affected by relatively rapid changes in precipitation, temperature, and humidity that result from a rapid and drastic change in regional or local climate patterns (e.g. prolonged drought, higher temperatures). Detrimental invasive species may change their distribution and abundance due to changes in the climate (e.g. mosquitoes may be more frequently found at higher elevations due to warming temperatures). Monitoring and relating climate and any climate change to the ecology of the Laupāhoehoe Forest is a major goal of the HETF.

Although natural disturbances such as hurricanes, droughts, flooding are regular occurrences in Hawai‘i, wide spread insect-driven defoliation can also impact the forest (koa moth (*Scotorythra paludicola*)). Similarly, ‘ōhi‘a undergoes periodic declines where entire stands of ‘ōhi‘a die off at the same time (Akashi and Mueller-Dombois 1995, Anderson et al. 2001, Mueller-Dombois 1980). Native species and ecosystems may have evolved under these disturbance regimes, but today, they may not be able to recover from such disturbances as readily due to small populations, changing climate, introduced diseases (‘ōhi‘a rust), and/or competition with non-native plant species. Further, these types of natural disturbances may increase as a result of climate change.

Illegal Human Activity

Illegal human activity occurs on a small scale, primarily in the form of illegal camping, off-road all-terrain vehicle use, dumping, unpermitted harvesting (koa, maile, hāpu‘u, and other native trees and plants), poaching, marijuana cultivation, and vandalizing signs and fences. These activities destroy infrastructure and native species. Some illegal activities create openings in the forest that can be invaded by invasive non-native plants.



Wildfire

Fire poses a threat to Laupāhoehoe Forest, particularly in the drier upper elevation during times of drought and in areas adjacent to human activity. Hawai‘i’s flora evolved with infrequent, naturally-occurring fire, so most native species are not fire-adapted and are unable to recover quickly after wildfires. Wildfires leave the landscape bare and vulnerable to erosion and non-native weed invasions (D'Antonio et al. 2000, Dunkell et al. 2011, Smith and Tunison 1992). Continued feral ungulate damage to native ecosystems can convert native forest to non-native grasses and shrubs, which provide more fuel for fire (Ainsworth Kauffman 2010, Cabin et al. 2000, Chynoweth et al. 2013, Cole et al. 2012, Nogueira-Filho et al. 2009, Scowcroft and Giffin 1983, Thaxton et al. 2010). Invasive non-native plants, particularly grasses, are often more fire-adapted than native species and will quickly exploit suitable habitat after a fire (D'Antonio et al. 2000, Mack and D'Antonio 1998). The principal human-caused ignition threats are from catalytic converters and other hot surfaces of vehicles or heavy equipment and illegal campfires. The principal natural ignition source is lightning.

There have been three fires recently in the vicinity of Laupāhoehoe Forest. The Piha Fire in 2008 started off Mana Rd. from a vehicle parked in dry grass. The fire burned 2,800 acres (1133 ha) of non-native grasses and koa canopy from Hopuwai Corral above Piha FR across to Laupāhoehoe FR and up to Mauna Kea FR. The Waipunalei Fire in 2012 was probably started from a lightning strike. The fire burned 22.2 acres (9 ha) near the koa mill in Waipunalei, between Laupāhoehoe Forest and the Humu‘ula section of Hilo FR. Fuels were mainly kikuyu grass with koa, ‘ōhi‘a and sugi pine canopy. The Keanakolu fire in 2013 started from an abandoned campfire below Keanakolu Cabins in the Humu‘ula Section of Hilo FR and burned approximately 3 acres (~1.2 ha).



2013 Keanakolu fire



Threats to Cultural Resources

Threats to cultural resources are in large part the same as the threats to biological and physical resources described below in excerpts from Maly and Maly 2006.

“In Hawaiian culture, natural and cultural resources are one and the same. Native traditions describe the formation (literally the birth) of the Hawaiian Islands and the presence of life on and around them, in the context of genealogical accounts. All forms of the natural environment, from the skies and mountain peaks, to the plateau lands, watered valleys and lava plains, and to the shoreline and ocean depths are believed to be embodiments of Hawaiian gods and deities.

The forest lands of this region represent significant native (endemic and indigenous) resources, and are part of a unique cultural landscape—in that the native flora, fauna, mist, rains, water, natural phenomena and resources, are all believed to be *kino lau* (the myriad body-forms) of gods, goddesses, and lesser nature spirits of Hawaiian antiquity. Knowledge of the environment and respect for the resources ensured a sustainable life upon the land. And in their evolving relationship with natural resources such as those of this region, Hawaiians came to consider everything about them as godly manifestations. Care for, and respect of the earth, meant that in-turn, the earth would care for the *kānaka* (people).”

Threats to resources such as stone features (walls, terraces, mounds, platforms, shelters, caves, trails or boundary *ahu*) and burials include vandalism and destruction during ground altering activities such as construction. The Hawai‘i State Historic Preservation Statute (Chapter 6E), affords legal protection to historic sites, including traditional cultural properties of ongoing cultural significance.



Cultural Resources in Laupāhoehoe Forest include maile and palapalai



OVERVIEW OF EXISTING MANAGEMENT

Numerous management actions have been completed and/or are ongoing in Laupāhoehoe Forest. The section below summarizes these actions and accomplishments.

Natural Resources

Forest Protection and Management

To date, DOFAW staff has built small fenced enclosures to protect approximately 35 acres (14 ha) of native forest habitat and rare and endangered plant species from feral ungulates (3 enclosures: Kilau Uka, Loulu, and Scowcroft units are each approximately 10 acres, and 10 enclosures are < 1/4 acre). The Kilau Uka enclosure near Blair Rd. was completed in 2008; the Loulu enclosure was completed in 2010, and the Scowcroft enclosure was installed in 2014. These enclosures are also used for restoration of rare plants through outplanting.

Invasive Non-Native Plant Control

DOFAW staff control priority non-native invasive plants in rare plant enclosures, along roadsides and in other priority areas. Staff spends approximately 30 person days/year working on weed control, with additional work scheduled in the summer when Youth Conservation Corps (YCC) crews are available.



DOFAW staff installs fencing to protect the forest



Rare Species Restoration

DOFAW staff work cooperatively with other organizations and agencies on rare plant recovery including the Hawai‘i State Plant Extinction Prevention Program (PEPP) and the Volcano Rare Plant Facility (VRPF) of the University of Hawai‘i. Management actions specific to rare plant recovery include rare plant surveys to locate wild individuals, protection of wild plants in fenced exclosures, collection of propagation and genetic storage materials and reintroduction through outplanting in fenced, protected exclosures. PEPP is focused on preventing the extinction of taxa with fewer than 50 individuals in the wild.

DOFAW staff follow rare plant collection and reintroduction guidelines recommended by the Hawai‘i Rare Plant Restoration Group (interagency group of rare plant experts)

<http://hear.org/hrprg/>. DOFAW staff tag and map the locations of all outplanted plants and monitor their survival and growth. Rare plants reintroduced into Laupāhoehoe Forest in fenced, protected exclosures through outplanting include: *Anoectochilus sandvicensis*, *Clermontia lindseyana*, *Clermontia pyralaria*, *Joinvillea ascendens*, *Ochrosia haleakalae*, *Phyllostegia macrophyllus*, *Phyllostegia warshaueri*, *Stenogyne macrantha* and *Trematolobelia grandifolia* (see Table 3 for species status and common (Hawaiian) names).



DOFAW staff monitor rare plants in Laupāhoehoe Forest

Monitoring

In 1982, a rare plant survey of the proposed NAR noted the presence or absence of certain priority invasive non-native plants (Cuddihy et al. 1982). More intensive invasive non-native plant monitoring was completed across 9 transects in the NAR portion of Laupāhoehoe Forest in 1988, to gather information for the 1989 management plan. These transects were re-monitored in 1998. In addition, in 2008 NAR staff monitored vegetation plots along new transects which included invasive non-native plant monitoring.

Forest birds in Laupāhoehoe Forest were surveyed as part of the Hawai‘i Forest Bird Survey (Hamakua Study Area) from 1976-1983 (Scott et al. 1986). These same transects were re-surveyed in 1993. In 2013, additional surveys were conducted to assess the status of forest birds as part of this management plan. Recent survey data has not yet been analyzed; however, four species of endangered bird species were detected: ‘akiapōla‘au, Hawai‘i ‘ākepa and Hawai‘i creeper was present above 4,500 ft (1,372 m), and ‘Io was also detected during the survey.

Research

Research conducted in the Laupāhoehoe Forest is a combination of long-term monitoring of environmental conditions and biotic responses, and directed research to address key questions about forest ecosystem function, health, and sustainability. Research in the Laupāhoehoe Forest is conducted by universities, government agencies including the USFS, and private organizations. Projects vary in focus, scope and length. Research projects proposed for the



Laupāhoehoe Forest are subject to review and permitting prior to access being granted for project implementation. A complete list of projects that have been conducted in Laupāhoehoe Forest to date can be found in on page 80 and is also available in HETF annual reports at www.hetf.us.

Long-Term Data Collection

Long-term monitoring is an integral component of research field sites such as the HETF. In the Laupāhoehoe Forest, long-term monitoring infrastructure has been established for vegetation, climate, and stream monitoring. The Hawai‘i Permanent Plot Network (HIPNET), a collaborative project between the University of Hawai‘i, the USFS, and University of California Los Angeles, has established a co-located vegetation plot and climate station in the Laupāhoehoe Forest that is part of a worldwide study of tropical forests with the Smithsonian Tropical Research Institute’s Center for Tropical Forest Science (www.ctfs.si.edu). Climate conditions in the forest are monitored by a weather station installed in 2009. The climate station extends 10 feet (3.3 m) above the forest canopy and collects data on rainfall, temperature, relative humidity, wind-speed, solar radiation (sunlight), soil moisture, soil temperature, and wind direction. The Laupāhoehoe climate station is part of the EPSCoR-ENDER (Experimental Program to Stimulate Competitive Research - Environmental Dynamics and Ecosystem Responses) Climate Network, an island-wide network of climate stations at locations across the island of Hawai‘i. Research conducted in the HIPNET will enable advancement in the studies of global change, ecohydrology, ecosystem services, remote sensing, restoration, community structure and organization, population genetics, comparative forest ecology and biogeochemical processes.

The Forest Inventory and Analysis Program (FIA) is a nationwide USFS program aimed at collecting, analyzing, and reporting information on the status and trends of America's forests. The Laupāhoehoe Forest includes 29 FIA plots (12 in the FR and 17 in the NAR) which represent a portion of the approximately 600-700 plots proposed for all of Hawai‘i. With the support and coordination of numerous entities in Hawai‘i, Hawai‘i’s FIA program collects additional information on the presence of invasive plants and other disturbances such as feral pigs to provide a baseline assessment of the current state of forests all over Hawai‘i. Plots are scheduled to be re-measured every 10 years to provide insights into changes in forest extent, composition, structure, and disturbances.

The hydrology of native forests and watersheds is an integral part of long-term sustainability of forest ecosystems, including aquatic biota. Freshwater availability for a variety of human uses and for its contribution to nearshore marine ecosystems is also critical. One facet of understanding forest hydrology is by monitoring flow in major streams through the use of stream gauges. One stream gauge has been established in Manowai‘ōpai Stream for monitoring Laupāhoehoe Forest and the gauge is currently maintained by the USFS.

Short Term Research Projects

In addition to long term data collection, the HETF supports a range of research projects that contribute to the greater ecological understanding of Hawai‘i’s forests and species. Research topics include species identification, monitoring, ecosystem services and life history studies, koa productivity, biodiversity and invasive species impacts and control. Representative examples of the diversity of topics include:

- Hawaiian hoary bat habitat occupancy, reproduction and diet



- Acoustic variability and loss of song complexity in Hawaiian honeycreepers
- Adaptive radiation in Hawaiian spiders
- *Drosophila* bar coding project as a method to determine species
- Native and non-native snail surveys
- ‘Ōhi‘a rust monitoring
- Assessing the *Scotorythra paludicola* (Lepidoptera: Geometridae) outbreak on koa: population abundance, rates of parasitism and patterns of spread
- Comparative nutritive values of traditional and exotic foraging substrates for upper elevation forest birds
- Quantifying the effects of ungulate and vegetation on the hydrology of Hawaiian tropical forests
- Sources and fates of nutrients on a substrate age gradient across the Hawaiian archipelago and their consequences for forest dynamics

Education and Outreach

Educational activities associated with Laupāhoehoe Forest currently include support for internships that focus on restoration and education (AmeriCorps, Youth Conservation Corps (YCC), and Pacific Internship Programs for Exploring Science (PIPES)), and securing national, regional, and local grants that fund educational programs. Further, IPIF staff collaborations with teachers at local middle and high schools have resulted in classroom field trips into the forest to learn about botany, ecology, natural resources management, traditional ecological knowledge and cultural geography. Many of the educational activities involve substantial contributions from additional partners including Mauna Kea Watershed Alliance, the USFWS, and the University of Hawai‘i (UH) at Hilo and UH-Mānoa.

Public Access and Recreation

Public access for recreational and cultural uses is ongoing in Laupāhoehoe Forest in accordance with existing rules and policies described earlier in this plan (see DESCRIPTION AND CURRENT CONDITION section). Current public access routes are shown Figure 3.

Infrastructure

Plans for development of two facilities for education and research outside the forest boundary were finalized in the Laupāhoehoe Construction Project EA. The Laupāhoehoe Science and Education Center is currently in the final stages of construction for building commissioning in 2015. The Forest Pavilion (Field Education Site) will be located on a 3-acre (1.2 ha) parcel located adjacent to the Laupāhoehoe Forest (Figure 3). It will consist of a covered pavilion, toilet/comfort station, and parking area, and it will be used primarily as a staging area for research and education trips into Laupāhoehoe Forest. See the Laupāhoehoe Construction Project EA (http://www.hetf.us/page/projects_plans/) for more detailed information regarding the project.



PROPOSED MANAGEMENT PROGRAM

The proposed management program outlined in this section includes new proposed actions as well as ongoing research, management and education projects. This section provides background information on projects within the main management focal areas, management objectives, and proposed actions to accomplish the objectives. In addition, some sections provide rationale for the proposed actions in cases where there were alternative actions considered.

Natural Resources

Forest Protection and Management

Background: The protection and management of forested watersheds and unique native Hawaiian ecosystems is a priority for Laupāhoehoe Forest. Effective management of forested watersheds provides fresh water for public use, improves water quality, reduces soil erosion, improves coastal water quality, and maintains native ecosystems. In addition, many native, threatened and endangered species rely on forested watersheds for their survival. These forested watersheds require active management to effectively address threats in order to persist for the benefit of current and future generations.

- Locations for conservation units were chosen in consultation with the LAC and high use hunting areas were avoided where possible.
- The areas planned for fencing have some of the highest quality, most intact native habitat in Laupāhoehoe Forest.
- Planned conservation units will protect existing populations of rare plants and animals and can also be used as restoration sites for rare species recovery.

Objective: Protect, manage and restore native ecosystems and species at Laupāhoehoe Forest by effectively managing conservation units and implementing forest restoration practices.

Proposed Actions:

1. Fence and remove feral pigs from two conservation units (Figure 9) to protect the biological and water resources and limit damage to native Hawaiian ecosystems. Without fencing, ungulate control requires ongoing effort, due to reproduction of existing populations and continued ingress from adjacent properties.
 - Conduct field surveys of final fence alignment to avoid any impacts to botanical, faunal, and archaeological resources.
 - Construct fencing of two conservation units (342 acres (138 ha) and 2,317 acres (938 ha) (subdivided by a cross fence (location to be determined))).
 - Install gates and walkovers for pedestrian access into fenced conservation units.
 - Implement feral pig control using approved methods following fencing to remove all pigs from within conservation units. Public hunting will be encouraged initially, but additional control methods including drives, trapping/release, staff control, and snaring, may be needed to remove all the ungulates not removed by active hunting within the conservation units.
 - Monitor fenced conservation units for pig ingress, and control pigs, if necessary.




2. Maintain all fences through regular inspection and maintenance and replace fences, when needed, including perimeter fencing to prevent cattle ingress from adjacent ranch lands.
3. Restore forested ecosystems in areas that have been disturbed (e.g. formerly logged areas, areas disturbed by cattle) through tree plants and invasive non-native plant control.
4. Plant koa and other native trees to restore native forest. Priority areas are depicted in Figure 9. Priorities include the following: 1) within the FR section along Mana Rd. and inward; 2) Shack Camp; and 3) section off Blair Rd. toward Waipunalei. Restoration is not limited to these areas and additional areas may also be restored. These priority sites were chosen because they have past damage from logging and feral cattle. They are also accessible for staff and volunteers, particularly with proposed infrastructure improvements at Shack camp (shelters and camping sites discussed in public recreation and access section below).
5. Control non-native pasture grasses and other non-native invasive weeds in restoration areas to enhance the natural regeneration of native trees and shrubs and prevent fire.
6. Construct emergency rare plant exclosures between 1-5 acres in size, when needed to protect individuals or populations of endangered plants.
7. Pursue potential land acquisitions of adjacent lands for protection and restoration of a large conservation landscape in cooperation with conservation partners (when applicable).

Rationale: Community and LAC member feedback related to conservation units in Laupāhoehoe Forest ranged widely from fencing the entire forest for protection to desires for no additional fences. As mentioned throughout this plan, the protection and management of forested watersheds and unique native Hawaiian ecosystems is a priority for the state within Laupāhoehoe Forest as only 35 acres are currently protected. Through discussion with the LAC and community members, the importance of Laupāhoehoe Forest to the community for hunting opportunities was also documented. Across Hawai‘i Island, state land managers are facing tough choices when tasked with protecting valuable native resources while still providing hunting opportunities. In particular with feral pigs, the negative impacts to the forest are well documented. The proposed conservation units in this plan attempt to meet conservation needs while considering hunting community desires. High quality native forest habitat areas that are less heavily used for public hunting were purposely selected. The proposed conservation units also provide suitable habitat for the recovery of rare and endangered species and are more feasible to manage because they are not dominated by invasive species, as is the case in lower elevation portions of the forest. One conservation unit with high quality native forest habitat includes an area dominated by tropical ash but this area has high restoration potential once fenced and the tropical ash controlled. When this plan is fully implemented, approximately 2,694 acres (1090 ha) or 22% of Laupāhoehoe Forest will be protected through conservation unit fencing.

In conjunction to Action 7 above, the state is also pursuing the development of additional access and/or acquisition of lands to expand hunting opportunities on appropriate lands.



Figure 9



Invasive Non-Native Plant Control

Background: Invasive non-native plants are a major threat to Laupāhoehoe Forest, and species with high potential for spreading and modifying habitat are a high priority for control. The overall approach includes preventing the establishment of new habitat modifying species that are either not currently present (e.g. miconia) or are still localized through biosecurity measures. For priority weeds already present, the goal is to identify control areas, eliminate all known occurrences within targeted control areas and/or to contain the further spread.

Objective: Protect intact native forest by preventing the establishment and/or removing high priority non-native, invasive plants and other invasive species.

Proposed Actions:

1. Regularly monitor and map the distribution of high priority invasive non-native plants, develop a comprehensive control strategy and revise control strategy, as needed, based on monitoring data.
 - Regularly compile transect monitoring data, incidental observations and reconnaissance surveys to map changes in invasive plant distribution and abundance and detect new species.
 - Cooperate with research on new mapping technologies (high resolution aerial imagery) to assist in monitoring and/or locating priority weeds for control.
2. Control priority non-native invasive plants in identified areas using approved methods.
 - Control invasive non-native plants in high quality native forest, particularly within fenced, ungulate-free management units. These areas are a high priority for more intensive management, and the recovery of native vegetation, reductions in ground disturbance and the spread of weeds by ungulates may increase the effectiveness of invasive plant control.
 - Target control of certain incipient invasive non-native plants (just beginning to invade) in unfenced areas to prevent their establishment and spread.
 - Focus control efforts in disturbed areas such as roads and trails as these often serve as corridors for weed establishment and spread.
 - High priority species present in Laupāhoehoe Forest include Australian tree fern, banana poka, Florida blackberry, cane tibouchina, *Clidemia*, *Ficus*, Himalayan ginger, mules foot fern, palm grass, passion fruit, *Polygonum chinensis*, strawberry guava, tropical ash, and yellow Himalayan raspberry.
 - A combination of control techniques including manual, mechanical and approved herbicides will be used to remove weeds. The technique selected will be based on the characteristics of the target species, the sensitivity of the area in which the species is found, and the effectiveness of the control technique.
 - Due to widespread and heavy infestations of certain weeds and limited resources, DOFAW will use approved biocontrol agents within the forest, when available, and if shown to be effective.
3. Monitor non-native invasive plants to determine whether weed control measures are effective and to detect changes in long term distribution and abundance.



4. Maintain procedures to prevent introduction of new weeds (see biosecurity section).
 - Avoid and/or reduce the inadvertent introduction and spread of weeds by staff, researchers and the general public working through education regarding biosecurity and implementation of biosecurity protocols for staff and research/education permittees.
 - Prevent the establishment of high priority invasive non-native plants species that are present in adjoining areas but not yet detected in Laupāhoehoe Forest. These species include miconia, faya, gorse, firethorn, *Cotoneaster pannosus*, and New Zealand flax.

Rare Species Restoration

Background: Landscape-scale habitat protection and management through management actions described in the habitat protection and management section are critical to the long-term integrity and recovery of native ecosystems including rare plants, forest birds and other native species. Such management actions, along with non-native invasive plant management and the prevention of new habitat-modifying weeds and harmful non-native species are the most critical actions needed to protect existing native habitat and rare species. However, in some instances, these actions are not enough to recover certain rare and endangered plants and animals. These species may have wild populations that are so small that the species cannot survive and recover without additional species-specific management.

Objective: Protect threatened and endangered plants and animals in Laupāhoehoe Forest and restore populations of these species in appropriate habitat to assist with the overall recovery of these species.

Proposed Actions:

1. Maintain the integrity of high quality forest ecosystems to the extent possible through fencing, feral ungulate control, non-native invasive plant control and preventing the introduction and establishment of other habitat-modifying species and new threats.
2. Map, monitor and protect existing wild populations of rare and endangered species to contribute to their population recovery and stabilization. Identify and remove threats to these species and ensure their long-term survival in secure and self-sustaining wild populations.
3. Re-introduce certain species of rare and endangered plants Table 3 in appropriate protected habitat through outplanting. Over the past decade, numerous species of rare plants have been propagated and reintroduced into fenced, ungulate-free areas to contribute to their overall recovery in the wild.
 - Coordinate rare plant management actions with the PEPP, VRPF and other agencies and organizations working on rare plant recovery.
 - Survey rare plants to locate wild individuals, collect propagation and genetic storage materials and reintroduce through outplanting.
 - Follow rare plant collection and reintroduction guidelines recommended by the Hawai'i Rare Plant Restoration Group.
 - Tag and map the locations of all outplanted plants and monitor survival and growth.



- Provide additional management of wild and/or reintroduced populations if needed (e.g. small fences around wild plants and populations that are not within fenced management units, control of damaging weeds, insects, slugs, plant disease and/or mammalian predators).
4. Determine additional actions needed to protect rare invertebrates. Previously discussed habitat management will also benefit rare native invertebrates, as they are generally dependent on native plants for food and as host plants.
 - Perform additional invertebrate surveys to inventory species, identify important habitat for rare species, and determine threats and needed recovery actions.
 5. Enhance habitats for native species through small mammalian predator removal and other habitat management actions.
 - Small mammalian predator removal (e.g., removal of rats, mongoose, cats) may provide significant benefits to endangered birds, plants, and endemic invertebrates, but is extremely difficult and costly to implement. DOFAW staff may implement predator removal in certain high priority areas (e.g. endangered bird nesting sites, rare plant restoration sites) using approved methods.
 - Other management may be implemented, as feasible. These include removal of larval habitats (e.g standing water providing mosquito breeding habitat) that may be responsible for seasonal epizootics of avian pox and malaria. Reducing or eliminating vespid wasps (yellow jackets) may also provide benefits to forest birds, as these wasps prey on insects that provide food for forest birds.

Monitoring

Background: DOFAW staff will continue to implement basic monitoring programs which are directly informing ongoing management. DOFAW staff regularly monitor ungulates, non-native invasive plants, rare plants and forest birds and are planning on continuing these monitoring programs. Additional monitoring is described under research, and will primarily be implemented by USFS staff and other researchers.

Objective: Monitor current status and trends of natural resources throughout Laupāhoehoe Forest as part of a long-term monitoring program.

Proposed Actions:

1. Continue ongoing monitoring programs for ungulates, non-native invasive plants and rare plants to measure the success of management and detect changes in abundance and distribution.
 - Monitor ungulates in fenced management units to detect the presence or absence of ungulates. Units that are free of ungulates will be regularly monitored to detect ingress animals. Units with active ungulate control programs will be monitored to assess the success of and/or direct control efforts.
 - Continue non-native invasive plant monitoring along transects to detect changes in distribution and abundance over time as well as detect incipient invaders. Control areas are monitored to determine the success of management efforts.



- Rare plant monitoring is conducted to assess the survival and growth of wild and re-introduced rare plants. DOFAW and PEPP program staff monitor rare plants to assess their survival and reproduction, collect propagation materials, search for additional wild individuals and determine if additional management is necessary.
2. Continue ongoing monitoring program for forest birds. Provide monitoring data to the Hawai'i Forest Bird Interagency Database Project for analysis of bird population densities and trends.
 3. Develop improved monitoring protocols, data management and analysis for existing monitoring programs and review and summarize past monitoring data and inventories.
 4. Develop and/or identify appropriate monitoring protocols and implement monitoring for key community indicators that are not currently being monitored (e.g., native vegetation communities, invertebrates, etc.).

Wildfire Prevention and Response

Background: Fire is a threat to the drier upper elevation portions of the Laupāhoehoe Forest. Many fires are caused by humans, so fire prevention measures will include increased educational efforts. DOFAW staff will respond to fires in Laupāhoehoe Forest using measures that result in the least amount of impact or disturbance to natural and archeological resources. The method of suppression will be determined by the on-site situation, with special regard to the potential expansion of fire damage to natural resources. Minimum impact methods of suppression will be applied whenever such methods are sufficient. Bulldozing is justified when a fire cannot be otherwise controlled and potential bulldozing damage is outweighed by a probable greater loss of natural and archeological resources.

Objective: Employ appropriate fire management strategies including pre-suppression, suppression, and post-suppression rehabilitation to reduce wildfire occurrence and minimize wildfire impacts.

Proposed Actions:

1. Implement fire prevention measures, including educational outreach to neighbors and signage along roads and road or area closures in the event of extreme fire danger.
2. Control invasive plants, particularly non-native grasses and plant common native species to restore certain disturbed areas to prevent fire and/or following damage from fire.
3. DOFAW staff to suppress fires safely and aggressively using appropriate means to minimize wildfire impacts.
4. Continue DOFAW staff training and certifications for effective and safe fire response.
5. Maintain access and fuelbreaks for fire pre-suppression and suppression.

Research

Background: The USFS and state of Hawai'i, along with the consortium of institutions and agencies involved with the HETF, will continue to encourage and facilitate research in Laupāhoehoe Forest. Research projects that contribute to the greater purpose of the HETF, that are relevant to land management issues and that are compatible with existing research and



management will be encouraged. USFS will support facilities to enhance the ability of the experimental forest to meet its goals for research and science. In addition, the USFS will facilitate access to basic biological, physical, and climatological data for the experimental forest through readily accessible web-based platforms and tools for researchers and the public to provide a foundation on which research projects can be built. All research within the HETF requires a valid permit.

Laupāhoehoe Forest provides many opportunities for research. Information on the basic natural history and abundance of the many endemic and often endangered plants and animals associated with Laupāhoehoe Forest is needed to understand how species may respond to changing environmental conditions (e.g., as a result of climate change) and how management and conservation measures can be used to help enhance adaptation. With its mixture of native and non-native dominated landscapes, Laupāhoehoe Forest provides an ideal site in which to test hypotheses about how invasive species impact ecosystems and determine the most effective methods of controlling or eliminating invasive species. Research aimed at effective ecosystem restoration is of great relevance not only in Hawai‘i and the Pacific Islands, and throughout the wet tropics.

Objective: Provide lands for conducting research that serves as a basis for the restoration, conservation and management of tropical forest ecosystems in Hawai‘i and across the tropics.

Proposed Actions:

1. Promote applied research with direct relevance to land management issues such as effective management of invasive species, forest restoration and climate change impacts on Laupāhoehoe Forest.
 - Host semi-annual meeting with land managers and community members to outline pressing information needs.
 - Host semi-annual meetings with research community to identify opportunities for collaboration and funding.
 - Bring together research institutions and conservation land stewards on Hawai‘i Island (e.g., US Fish and Wildlife Service, US Park Service, Nature Conservancy, Kamehameha Schools) to establish a network of field sites that together can be used to promote research on a broader suite of questions about tropical ecosystem conservation.
 - Offer stimulus funding in the form of modest awards to promote research in key areas.
 - Host theme-based research discussions and symposia at the Laupāhoehoe Science and Education Center, or elsewhere, that is sponsored by the experimental forest.
 - Promote and support dissemination of research conducted in association with the experimental forest (e.g., research highlights, presentations, acknowledgements in publications and presentations).
2. Effectively administer and coordinate the research application process including review of applications, issuance of research permits, research compliance with permit conditions and relevant land designation statutes and rules.
 - Provide administrative support to research permit review and approval process.



- Provide research expertise and management guidance to address potentially controversial or complex research proposals.
 - Promote timely and efficient processing of requests to encourage use.
3. Establish and maintain a system for archiving research data and reports to facilitate the exchange and transfer of information among agencies, scientists, and the community.
 - Maintain a publically available, web-based archive of baseline monitoring data, historical data and descriptions of all research activities. Historical data includes archive of maps, land-use histories, cultural practices and historical data relating to the HETF.
 - Require all permitted researchers to make appropriate contributions to the data archive, including project descriptions, project reports, sampling locations, and publications that have resulted from research.
 - Maintain an electronic library of publically available reports, research and publications that pertain to the HETF available on the Internet. Materials protected through copyright would be available by request.
 - Publish syntheses of research and monitoring activities and results from HETF to provide information and findings more readily accessible to research, conservation, and local community members.
 4. Improve dissemination of scientific research information and results to land managers and the local community.
 - Provide presentation series, workshops and seminars in association with the experimental forest to disseminate information to the local community.
 - Partner with University of Hawai‘i, Hawai‘i Community College, and local K-12 schools to jointly host field trips and field courses.
 - Host annual open houses at the experimental forest units to highlight research activities, associated knowledge gained, and opportunities for community involvement.
 - Develop a diversity of outlets for information in non-technical formats (e.g., calendars, posters, computer apps).
 5. Promote cultural research including information on traditional Hawaiian use/presence in the forest, oral histories, cultural impacts of management actions and archeological studies.
 - Collaborate with cultural researchers at University of Hawai‘i and other schools to use Laupāhoehoe Forest as a focus of cultural research.
 - Seek funding for cultural research including integration of culture into scientific research as well as archeological and ethnographic studies.
 6. Link ongoing research to education programs by encouraging researchers to work with local schools such as the Laupāhoehoe Community Public Charter School and universities.
 - Collaborate with University of Hawai‘i and Hawai‘i Community College to develop courses and seminars that incorporate field site visits to the experimental forest.
 - Collaborate with local educators to develop course curricula and materials that highlight or utilize research findings associated with the experimental forest.



7. Encourage basic research and monitoring to establish historical baselines of all natural resources.
 - Collect data on vegetation as part of a global tropical forest monitoring network, to monitor the status and condition of vegetation at Laupāhoehoe Forest and to develop vegetation maps.
 - Maintain a weather station as part of a larger island-wide network of climate stations across the island of Hawai‘i to collect data on rainfall, temperature, relative humidity, wind-speed, solar radiation (sunlight), soil moisture, soil temperature, and wind direction.
 - Establish and maintain stream gauges to monitor natural stream flows, water quality and sediment in a non-destructive manner. Hydrologic information can be used as a foundation for research on aquatic ecology, watershed dynamics, and climate change.
 - Perform regular surveys that systematically document and describe plant and animal species distribution and status to provide a baseline for research and assist with developing management actions and assessing their success.

Education and Outreach

Background: Education and outreach are key component of the overall vision for Laupāhoehoe Forest. Education and outreach goals span six focal areas: formal training for professionals; community outreach; cultural training; demonstration for natural resources managers; student research; and academic education. Educational goals will be accomplished through a strong reliance on partnerships and will be integrated with other aspects of research and natural resource management.

Objective: Serve as a center for forest education, training, demonstration and outreach on tropical forests, conservation biology, and natural resource management for groups ranging from school children to land managers, scientists and the general public.

Proposed Actions:

1. Encourage appropriate educational and cultural uses of Laupāhoehoe Forest through the development of general criteria, priorities and rules to effectively manage multiple educational uses.
2. Provide general orientation and training (e.g. on-line video) for all new research permittees and educational programs that includes recommendations on forest stewardship and invasive species prevention protocols as well as cultural components.
3. Collaborate with universities and relevant non-governmental organizations to integrate classes, student research/internships and provide support via Center facilities.
 - Host courses from local and visiting universities.
 - Co-develop courses with local and off-island universities that incorporate the HETF into the field component of classroom courses or where the HETF and perhaps other field sites are the focus of field courses.
 - Work with instructors from various universities to formulate courses that also contribute to our understanding of tropical ecosystems.



4. Foster and support undergraduate and graduate student research opportunities and research internships through partnerships with the University of Hawai‘i, other universities and local research agencies and organizations.
 - Identify research projects suitable for undergraduate and graduate students.
 - Sponsor students (e.g., projects, mentoring, equipment, funding) in conducting research pertinent to the HETF.
 - Invite researchers from other institutions to come to the HETF to conduct their research.
5. Provide a connection to nature and promote forest stewardship through engagement activities that involve the public and through collaboration with K-12 education program partners and other community partners including but not limited to the Laupāhoehoe Community Public Charter School. Educational activities include:
 - Educator workshops and programs
 - Field projects geared toward kids
 - Partnering with schools, educators, community and other non-governmental organizations to create and implement activities that facilitate forest stewardship
6. Communicate research findings, land use, and management goals to the community via:
 - Community field trips
 - Informational materials suitable for non-professionals of all ages
 - Participation in community events
 - Service learning opportunities
 - Interpretive trails and guided walks
 - Public participation in scientific research
7. Provide work experience and formal professional development training to land management professionals in ecology, conservation, and restoration of natural and cultural resources.
 - Provide opportunities for students in internship programs, such as PIPES to join research teams for short periods of time.
 - Provide opportunities for young professionals in YCC programs to understand research needs and management applications.
 - Enable emerging professionals in AmeriCorps programs to experience working alongside research scientists and natural and cultural resource managers.
 - Provide short-courses (0.5 to 5 days) designed for working professionals to help integrate new information and ideas into agency activities and approaches.
8. Serve as a demonstration site for land managers by providing information, tools and techniques through demonstration research, conservation, and restoration projects.
 - Engage managers in the design and implementation of research to understand effective conservation and restoration approaches.
 - Engage managers in the design and implementation of biophysical monitoring programs and projects.



- Engage managers in the design and implementation of science based restoration projects.
 - Engage managers in the role of traditional ecological knowledge in land management.
9. Provide readily accessible scientific information through web-based platforms and tools.
 10. Encourage researchers to share research results with the local community through informational presentations to schools and community groups and popular articles.
 11. Hire staff and/or establish volunteer positions to facilitate educational and outreach experiences (e.g. education staff, rangers).
 12. Serve as a site for alternative educational activities for non-profit groups and organizations (e.g. search and rescue training, orienteering, survival skills, back-country travel, hunter education programs, forest stewardship, Junior Youth Council, recreational and/or life skills, cultural immersion and traditional ecological knowledge training).

Public Access and Recreation

Background: Laupāhoehoe Forest is protected and managed by the state for the benefit of the people of Hawai‘i, and is open to the public for various recreational and cultural uses. While the public is allowed to access and hike or hunt in any portion of the forest, there are limited legal access points and only a few minimally maintained and marked trails. This area is a rough and remote rainforest wilderness and there are currently no amenities for recreational users. The access and recreational improvements proposed are intended primarily for local residents and to improve staff management access rather than for large-scale ecotourism. Large groups (over ten people) accessing the NAR require an HETF permit. The LAC is also supportive of improving public access elsewhere in the broader region, which has similar access issues.

Objective: Improve public access and recreational opportunities in Laupāhoehoe Forest consistent with maintaining natural resources and the wilderness character of these lands.

Proposed Actions:

1. Improve Public Access
 - Work with adjacent landowners to survey roads through private and state-leased lands depicted on some maps (“paper roads”) and/or work on alternative access to the forest.
 - Work with the County to ensure all future subdivision plans in the area include public access to the forest.
 - Pursue potential land acquisitions through fee-simple purchase (e.g. possible purchase of Waipunalei from Parker Ranch to connect forest reserve sections increasing access by eliminating private land barriers).
 - Consider DOFAW acquisition of long-term leases of state and/or private lands adjacent to the forest when current leases expire.
 - Consider adding appropriate trails and accesses (e.g. Maulua Trail) to the Nā Ala Hele Statewide Trail and Access System to improve overall management.



2. Trail Maintenance (Pedestrian) - Maintain existing trails (Spencer, Maulua & Peneki) as primitive trails (minimally maintained and marked) (Figure 10).
 - Peneki can serve as a connector trail from Spencer to Blair Rd. for both public recreational and research use.
 - Spencer can provide both public recreational and research use.
 - Maulua Trail can be used for education and outreach, public recreation and research and management.
 - Exclosure trail is used by DOFAW staff for management purposes and can also be used as a connector trail by the public.
 - Control spread of non-native invasive plants by providing boot brushes and informational signage at all trailheads and access points.
3. Trail Improvement (Pedestrian) - Improve existing and/or create and maintain new trails. Trail improvement is a lower priority than maintenance of existing trails; however the trails below would be minimally maintained and marked in a similar fashion to the maintained trails above. Specific alignments for new trails generally described below have not yet been determined on the ground and may vary depending on the vegetation and terrain (Figure 10).
 - Create trail along the north fence line (upper boundary) from Mana Rd. to Blair Rd. This would provide a public trail from Mana Rd. to Blair Rd. and would also be useful for management access. This trail would be relatively easy to create because it would be along an existing fence line.
 - Improve Spencer trail to Peneki and Peneki to Blair. These improvements would provide the public a legal connector within the forest boundary from Spencer to Peneki as well as fulfill the public interest in connecting Spencer-Peneki-Blair-Mana Rd.
 - Improve Maulua Trail to the south boundary/fence line. This improvement would provide an education/outreach trail for community forest restoration projects and would also be useful for management access. This trail can easily be improved given terrain and existing vegetation.
 - Create trail from Mana Rd. to Maulua roughly parallel to the southern forest boundary (specific location to be determined). This would create a loop trail for the upper area for the public and education/outreach opportunities as well as being useful for management access.
 - Create a trail at the FR/NAR boundary (specific location to be determined). This would provide a public trail through high quality forest and would help with management and research site access for management, monitoring and research, particularly for invasive non-native plants.
4. Hunting - Maintain and improve public hunting opportunities in Laupāhoehoe Forest.
 - Secure and improve/create access and trails as outlined above.
 - Facilitate additional hunter education classes in the Laupāhoehoe region.
 - Consider changing permitted hunting method for Unit C (currently rifle only) if desired by the hunting community.
 - Work with DOCARE to address hunting community concerns about illegal activities at Laupāhoehoe Forest and elsewhere on the island.



5. Camping and Shelters - Establish designated camping area at Shack Camp with primitive camp sites (Figure 10).
 - Construct a shelter or cabin at Shack Camp for both public recreational and management staff use. Cabin would be open to the public using a permitted reservation system.
 - Establish other infrastructure needed for camping and shelters at Shack Camp (composting toilet, fire pits, helicopter landing zone).

Rationale: Community and LAC member feedback related to public access in Laupāhoehoe Forest ranged widely from comments that current public access is adequate and no additional management actions are needed, to recommendations for expansion and enhancement of vehicular, mountain biking and pedestrian access.


- **Pedestrian access within the forest** – Feedback ranged from current pedestrian opportunities are adequate and no additional management actions are needed, to recommendations to consider the development of highly developed and maintained trails. A majority of comments recommended improving pedestrian access to allow a pedestrian to follow a recognized trail through the forest that would connect to Blair Rd. Due to the rugged terrain of Laupāhoehoe Forest and financial resources needed to create and maintain trails, the proposed actions primarily focus on improving existing primitive trails to allow pedestrians to traverse the forest and enter and exit at existing legal access points. There are numerous primitive trails throughout Laupāhoehoe Forest created and used by the hunting community, but only three of these existing trails would be identified as recognized public trails.
- **Vehicular access to and within the forest** – Feedback ranged from recommendations to dismantle Blair Rd., to opening Blair Rd. to public vehicular access. Blair Rd. is a valuable resource for management, research and education/outreach opportunities in Laupāhoehoe Forest and for these reasons dismantling Blair Rd. was not considered as an action. Opening Blair Rd. to public vehicular access was not considered as an action considering entry is through private lands, road maintenance considerations, and increased risks associated with transport of invasive species. Feedback from LAC members included holding an annual open house where this could be accommodated or agencies offering chaperoned trips for the public. While an annual open house is not feasible or realistic due to public vehicular access restrictions across neighboring private lands, both USFS and DOFAW currently offer agency chaperoned service learning/outreach opportunities and these types of activities are also proposed for expansion.
- **Mountain biking** – Feedback ranged from mountain biking should not be allowed, to trails should be maintained/created to accommodate mountain biking. Biking is legal on FR roads including Blair Rd. but currently the only way to reach Blair Rd. on a bike is through private property, which is not legal. An alternative considered was multi-use trails that would allow for legal entry; however this was not selected as a plan action because there were only a small number of trails proposed for pedestrian improvement, concerns about user conflicts, and bicycle damage to trails.



- **Hunting** – Feedback from LAC hunter working group members included a desire for the state to pursue game management in the unfenced conservation units within Laupāhoehoe Forest primarily to allow for closures to create recovery periods coupled with DOCARE enforcement. Game management to increase game mammal populations for hunting conflicts with DOFAW management priorities for Laupāhoehoe Forest, as well as with federally designed critical habitat for endangered species. As mentioned elsewhere in this document, the proposed conservation units in this plan attempt to meet conservation needs while also considering hunting community desires. High quality habitat that is less heavily used for public hunting due to remoteness was purposely selected for the conservation units. This plan seeks to increase public hunting opportunities in more accessible areas outside the fenced conservation units through improvements in access. Once this plan is fully implemented, approximately 9,649 acres (3905 ha), or 78% of Laupāhoehoe Forest will be available for public hunting.



Figure 10



Infrastructure

Background: Infrastructure is needed to improve management, research and education and public recreational uses in Laupāhoehoe Forest. Infrastructure includes roads, facilities, helicopter landing zones and structures, cabins/shelters and equipment.

Objective: Provide and maintain infrastructure and facilities to enhance the ability of the Laupāhoehoe Forest to meet its goals for management, research, education, and demonstration.

Proposed Actions:

1. Develop and maintain roads, cabins/shelters and campsites and helicopter landing zones for resources management actions, the functions of the HETF and for public recreational use and safety (Figure 10).
 - Develop facilities at Shack Camp (described above in public access and recreation section).
 - Establish a forest management shelter and landing zone at 3,500 feet (1067 m) elevation on the south east side of the NAR (specific location to be determined). This shelter will also be available for the public.
 - Establish approximately three other forest management shelters, as needed - locations to be determined.
 - Establish additional helicopter landing zones to be used for management and search and rescue operations - locations to be determined. Helicopter landing zones will use existing natural openings to avoid any damage to natural resources.
2. Ensure the development and maintenance of facilities and infrastructure has minimal impacts on the environment and natural and cultural resources.



ADMINISTRATION

Coordination

The HETF Cooperative Agreement states that "owing to the many values and benefits that arise from research, education and demonstration on the HETF and elsewhere, the Parties (the USFS and the state of Hawai‘i) further agree they will consult and reach agreements with each other to coordinate research, management, and education activities." Coordination of research, management and education is managed by the HETF Planning Group, which includes IPIF and DOFAW staff and two to three external partners. The HETF Planning Group is facilitated by the USFS and meets quarterly. The USFS produces an annual report for the BLNR and NARS Commission summarizing research, management and monitoring actions in the HETF.

Permitting

Permit applications for research and education activities Laupāhoehoe Forest are reviewed by agency staff in the HETF Planning Group as well as the LAC (which provides comments and/or recommendations). Permit processing and tracking is administered by the USFS. Signing authority for all permits lies with the Hawai‘i Island DOFAW Branch Manager (permit approval authority for the NAR was delegated to the Branch Manager by the NARSC on May 21, 2007). All research permits are valid for one year and require an annual report.

Cultural Resources

All state and federal employees, permittees and the public are required to comply with state and federal laws relating to the protection of cultural resources. All cultural and historical sites should be left alone and artifacts should not be collected. Burial sites and archeological sites are often accidentally disturbed either by nature (erosion) or by human activity through projects that involve excavation. Chapter 13-300, Hawai‘i Administrative Rules addresses rules of practice and procedure relating to burial sites and human remains. If a burial site is discovered, activity in the immediate area must be stopped and remains left in place. Reporting a burial site disturbance is required by law (Hawai‘i Revised Statutes, Chapter 6E) and reports of burial sites or other cultural resources should be made to the state Department of Land and Natural Resources, [Historic Preservation Division](#) Kakuhihewa Building, 601 Kamokila Blvd., Suite 555, Kapolei, HI 96707; Ph: (808) 692-8015.

All Federal agencies are subject to The [Native American Graves Protection and Repatriation Act](#) (NAGPRA), a Federal law passed in 1990. NAGPRA provides a process for museums and Federal agencies to return certain Native American cultural items -- human remains, funerary objects, sacred objects, or objects of cultural patrimony -- to lineal descendants, and culturally affiliated Indian tribes and Native Hawaiian organizations. NAGPRA includes provisions for unclaimed and culturally unidentifiable Native American cultural items and penalties for noncompliance and illegal trafficking.

Biosecurity

Biosecurity is a set of precautions that aim to prevent the introduction and spread of harmful organisms (pests, pathogens or invasive species) into Laupāhoehoe Forest. New plants and animals arrive in the islands on a continual basis. On average, 100 new plants, 20 species of insects, plus the occasional disease are introduced to Hawai‘i each year (Nature Conservancy of



Hawai‘i 2003). The source population of an invasive species can be the mainland, another island in the Hawaiian archipelago or even another area from the same island. Preventing the introduction of new invasive species into Laupāhoehoe Forest by staff, researchers and the general public is a high priority as these introductions only serve to increase the funding needed to control these species and further put Hawai‘i’s native forests at risk.

Organism introduction in Laupāhoehoe Forest can occur via transportation by animals or humans, the wind and/or through species nearby expanding their range (e.g., coqui frog and little red fire ant). There is also the risk of introductions from management work such as outplanting native plants grown in a nursery or road maintenance with heavy equipment that is not clean. Staff and individuals approved for research or education permits are provided information and protocols to help minimize or eliminate the introduction and spread of alien organisms into Laupāhoehoe Forest (Appendix B – Biosecurity Protocols). Staff and permittees are responsible for making sure that clothing, equipment, and vehicles are free of seeds, dirt, or organisms to lessen the chance of introducing any non-native plants or animals. Boots soles should be brushed and all equipment thoroughly cleaned prior to entry into Laupāhoehoe Forest.

Sightings of new alien organisms within Laupāhoehoe Forest or existing high risk organisms seen in previously un-infested areas should be reported to DOFAW or the HETF.

Safety

Overall Laupāhoehoe Forest terrain is very rugged and existing trails are primitive, uneven and muddy. The weather can be variable and conditions for heat stroke and hypothermia are possible given the elevation and weather patterns in this forest. Cell phone service is intermittent. Hunting with rifle is permitted in the upper portion and hunting with dogs is permitted in the lower portion. Forest visitors/users should dress appropriately for variable weather conditions, wear highly visible attire in consideration of hunting, and travel with adequate food and water. Other hazards include but are not limited to flash floods and falling trees and/or branches.

Vehicle access to the makai forest boundary currently includes traveling over a low water crossing. Heavy rains in the mauka areas of the forest can create flash floods and hazardous conditions at the low water crossing.

Forest visitors are advised to always identify a person to check in and out with that knows their planned route and can alert search and rescue teams should the forest visitor not check in when expected.

Budget and Staffing

NAR staff for the island of Hawai‘i work on all eight NAR on the island, including Laupāhoehoe. Currently, NAR staff include six DOFAW staff, six University of Hawai‘i contractors (Pacific Cooperative Studies Unit) and 2 year-round interns.

Forestry staff works on all nineteen forest reserves on the island. Currently there are eight staff (1 Botanist, 4 Foresters and 3 Technicians).



USFS staff works across the Hawaiian Islands and the Pacific. Staff dedicated to the HETF currently includes two full time administrative positions and the IPIF Director. Education and outreach goals are currently met via contributions to partner programs such as YCC, PIPES and watershed partners as well as through USFS volunteers.

Table 7 outlines estimated expenses related to specific projects and activities such as equipment, supplies, additional staff or contractor time. Costs are very broad estimates and will vary considerably over the time-frame of the plan. Existing staff costs are considered fixed and outlined above, although these may change due to state and federal budget fluctuations. Costs described as ‘to be determined’ or ‘TBD’ are unknown at this time. Funding to implement the estimated budget will be sought from various state, federal and private entities through annual budget requests as well as grants and/or other funding sources.



Youth Conservation Corps (YCC) interns assist staff with management actions while getting education and training in natural resources management



Table 7. Estimated costs

ACTION	DESCRIPTION	Periodic estimated costs		Annual estimated costs	ESTIMATED TOTAL COST (15 years)
		Amount	Time scale		
Natural Resources					
<u>Forest Protection and Management</u>					
1. Fence Construction and pig removal from conservation units	Fence construction (materials and contractual labor)	\$5,000,000.00	one time	\$0.00	\$5,000,000.00
	DOFAW staff feral pig removal	\$350,000.00	one time	\$0.00	\$350,000.00
2. Inspect/Maintain/Replace all fences	DOFAW staff, supplies/materials	\$0.00	n/a	\$15,000.00	\$225,000.00
3. Restore forested ecosystems	DOFAW staff, supplies/materials	\$0.00	n/a	\$20,000.00	\$300,000.00
4. Construct emergency rare plant exclosures	DOFAW staff, supplies/materials	\$0.00	n/a	\$5,000.00	\$75,000.00
5. Pursue potential land acquisitions for protection and restoration of a large landscape	DOFAW and USDA-FS Staff – No additional Costs	\$0.00	n/a	\$0.00	\$0.00
<u>Invasive Non-native Plant Control</u>					
1. Monitor/ map high priority invasive non-native plants and develop a control strategy	DOFAW staff, supplies/materials	\$0.00	n/a	\$5,000.00	\$75,000.00
2. Control priority non-native invasive plants in identified areas using approved methods	DOFAW staff, supplies/materials	\$0.00	n/a	\$35,000.00	\$525,000.00
3. Monitor to determine whether control is effective and detect change in distribution/ abundance.	Costs included in #1 (invasive non-native plant control)	\$0.00	n/a	\$0.00	\$0.00
4. Maintain procedures to prevent introduction of new weeds (biosecurity)	DOFAW and USDA-FS Staff – No additional Costs	\$0.00	n/a	\$0.00	\$0.00
<u>Rare Species Restoration</u>					
1. Maintain the integrity of high quality forest ecosystems	Costs included in #1 (forest protection and management and invasive non-native plant control)	\$0.00	n/a	\$0.00	\$0
2. Map, monitor and protect existing wild populations of rare and endangered species	DOFAW staff, supplies/materials - \$5,000/year	\$0.00	n/a	\$5,000.00	\$75,000
3. Re-introduce rare and endangered plants in appropriate protected habitat through outplanting.	DOFAW staff, supplies/materials - \$5,000/year	\$0.00	n/a	\$5,000.00	\$75,000
	Contractor Costs (VRPF and PEPP) - \$5,000/year	\$0.00	n/a	\$5,000.00	\$75,000
4. Determine addition actions needed to protect rare invertebrates	Costs to be determined	TBD	TBD	TBD	TBD



ACTION	DESCRIPTION	Periodic estimated costs		Annual estimated costs	ESTIMATED TOTAL COST (15 years)
		Amount	Time scale		
5. Enhance habitats for native species through small mammalian predator removal and other management	Costs to be determined	TBD	TBD	TBD	TBD
Monitoring					
1. Continue ongoing monitoring programs for ungulates, non-native invasive plants and rare plants	DOFAW staff, supplies/materials; Additional Costs included in #1 (invasive non-native plant control), #2 (rare species restoration)	\$0.00	n/a	\$5,000.00	\$75,000
2. Continue monitoring program for forest birds.	DOFAW staff and contractor monitoring/ data analysis costs - every five years (years 2019, 2024 and 2029)	\$35,000.00	\$3.00	\$0.00	\$105,000
3. Develop improved monitoring protocols, data management and analysis	Contractor staff costs - \$10,000/year	\$0.00	\$0.00	\$10,000.00	\$150,000
4. Develop appropriate monitoring protocols and implement monitoring for key community indicators	Costs included in #3 (Monitoring)	\$0.00	\$0.00	\$0.00	TBD
	Other costs TBD	TBD	TBD	TBD	
Wildfire Prevention and Response					
1. Implement fire prevention measures	DOFAW Staff – No additional Costs	0	n/a	0	0
2. Control invasive plants to restore disturbed areas to prevent fire and/or following damage from fire.	Costs included in #2 (invasive non-native plants)	0	n/a	0	0
3. DOFAW staff to suppress fires to minimize wildfire impacts.	TBD – dependent on size, location and intensity of fire	TBD	TBD	TBD	TBD
4. Continue DOFAW staff training and certifications for fire response.	DOFAW Staff – No additional Costs	0	n/a	0	0
5. Maintain access and fuelbreaks for fire pre-suppression and suppression.	DOFAW Staff – No additional Costs	0	n/a	0	0
Research					
1. Promote applied research with direct relevance to land management issues	Potential monetary incentives to promote work through competitive proposals and/or grants as well as contract work to target specific goals.	\$0.00	n/a	\$20,000.00	\$300,000.00
2. Effectively administer and coordinate the research application process	USDA-FS and DOFAW staff – no additional costs	\$0.00	n/a	\$0.00	\$0.00
3. Establish and maintain system for archiving research data and reports to facilitate exchange and transfer of information	USDA-FS – no additional costs; possible tie in with education/outreach staff see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
4. Improve dissemination of scientific results to land managers and the local community.	USDA-FS – no additional costs; tie in with education/outreach costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00



ACTION	DESCRIPTION	Periodic estimated costs		Annual estimated costs	ESTIMATED TOTAL COST (15 years)
		Amount	Time scale		
5. Promote cultural research including information on traditional Hawaiian use/presence, oral histories, cultural impacts of management actions and archeological studies.	Promote work through competitive proposals and/or grants as well as contract work to target specific goals.	\$0.00	n/a	\$20,000.00	\$300,000.00
6. Link ongoing research to education programs	USDA-FS staff – no additional costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
7. Encourage basic research and monitoring to establish historical baselines of all natural resources.	USDA-FS staff – no expected additional staff costs; potential monetary incentives to promote work through competitive proposals and/or grants	\$0.00	n/a	\$0.00	\$0.00
Education and Outreach					
1. Encourage appropriate educational and cultural uses through the development of general criteria, priorities and rules.	DOFAW and USDA-FS Staff – No additional Costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
2. Provide general orientation and training for all new research permittees and educational programs.	USDA-FS staff – no additional costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
3. Collaborate with universities and non-governmental organizations to integrate classes, student research/internships and provide support.	USDA-FS staff– no additional costs; possible tie in with education/outreach staff see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
4. Foster and support undergraduate and graduate student research opportunities and research internships.	USDA-FS staff – no additional costs; apply for funding to support training positions	\$0.00	n/a	\$20,000.00	\$300,000.00
5. Provide a connection to nature through educational activities that involve the public and K-12 partners.	USDA-FS and DOFAW staff – no additional costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
6. Communicate research findings, land use, and management goals to the community	DOFAW and USDA-FS staff– no additional costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
7. Provide work experience/professional development training to land management professionals	DOFAW and USDA-FS Staff – No additional Costs; tie in with EDU/OUTREACH #11; apply for funding to support training positions such as YCC and 21 CSC \$50K+ annually	\$0.00	n/a	\$50,000.00	\$750,000.00
8. Serve as a demonstration site for land managers by providing information, tools and techniques	DOFAW and USDA-FS staff– no additional costs; possible tie in with education/outreach staff costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00



ACTION	DESCRIPTION	Periodic estimated costs		Annual estimated costs	ESTIMATED TOTAL COST (15 years)
		Amount	Time scale		
9. Provide readily accessible scientific information through web-based platforms and tools.	USDA-FS staff– no additional costs; tie in with education/outreach costs see EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
10. Encourage researchers to share research results with the local community	DOFAW and USDA-FS Staff – No additional Costs	\$0.00	n/a	\$0.00	\$0.00
11. Hire staff and/or establish volunteer positions to facilitate educational and outreach experiences	\$75,000/year for contract staff salary/benefits, materials/supplies and volunteer program	\$0.00	n/a	\$75,000.00	\$1,125,000
12. Serve as a site for alternative educational activities for non-profit groups and organizations	Costs included under EDU/OUTREACH #11	\$0.00	n/a	\$0.00	\$0.00
Public Access and Recreation					
1. Improve Public Access	DOFAW and USDA-FS staff – no additional costs; TBD funds to purchase or lease adjacent lands	TBD	TBD	TBD	TBD
2. Trail Maintenance (Pedestrian) - Maintain existing trails (Spencer, Maulua & Peneki)	DOFAW and USDA-FS staff – no additional costs; apply for funding to support training positions such as YCC and 21 CSC – see EDU/OUTREACH #7	\$0.00	n/a	\$5,000.00	\$75,000.00
3. Trail Improvement (Pedestrian) - Improve existing and/or create new trails and maintain	DOFAW and USDA-FS staff – no additional staff costs; apply for funding to support training – see EDU/OUTREACH #7	\$0.00	n/a	\$5,000.00	\$75,000.00
4. Hunting - Maintain and improve public hunting opportunities in Laupāhoehoe Forest	TBD funds to purchase or lease adjacent lands	TBD	TBD	TBD	TBD
5. Camping and Shelters - Establish designated camping area at Shack Camp with primitive camp sites	\$50K/shelter; \$8K camping pavilion; 3 campsites @ \$1500/each; \$2,500/year to maintain shelter(s)/campsites	\$62,500.00	one time	\$0.00	\$62,500.00
Infrastructure					
1. Develop and maintain roads, cabins/shelters and campsites and helicopter landing zones	USDA-FS staff– no additional costs; materials and labor associated with project completion – see PUBLIC ACCESS/REC #5; \$5,000/year to maintain shelter(s), camping sites and LZ's	\$0.00	n/a	\$5,000.00	\$75,000.00
2. Ensure the development/maintenance of facilities and infrastructure has minimal impacts	DOFAW, USDA-FS, and MKWA staff – no additional costs; apply for funding to support cabins/shelters and campsites; road maintenance costs \$3K/year	\$0.00	n/a	\$3,000.00	\$45,000.00

TOTAL COST OVER 15 YEARS

\$10,212,500.00

REFERENCES

- Ainsworth, A. and J. B. Kauffman. 2010. Interactions of Fire and Nonnative Species Across an Elevation/Plant Community Gradient in Hawaii Volcanoes National Park. *Biotropica* 42(6): 647-655.
- Akashi, Y. and D. Mueller-Dombois. 1995. A landscape perspective of the Hawaiian rain forest dieback. *Journal of Vegetation Science* 6: 449-464.
- Anderson, R. C., et al. 2001. Dieback of *Acacia koa* in Hawaii: ecological and pathological characteristics of affected stands. *Forest Ecology and Management* 5576: 1-14.
- Aplet, G. H., S.J. Anderson and C.P. Stone. 1991. Association between feral pig disturbance and the composition of some alien plant assemblages in Hawai'i Volcanoes National Park. *Vegetation* 95: 55-62.
- Atkinson IAE. 1985. The spread of commensal species of *Rattus* to oceanic islands and their effects on avifaunas. In: Moors PJ (ed) *Conservation of Island birds*. ICBP Technical Publication, Cambridge, pp 35–81.
- Baker, J. K. 1979. The feral pig in Hawai'i Volcanoes National Park, *Proceedings: First Conference on Scientific research in the National Parks*, pp. 365-367.
- Bruland, G. L., Browning, C. A., & Evensen, C. I. (2010). Effects of feral pigs (*Sus scrofa*) on watershed health in Hawai'i: A literature review and preliminary results on runoff and erosion. In J. Roumassett, K. Burnett, & A. Balisacan (Eds.), *Sustainability science for watershed landscapes* (pp. 251–277). Singapore: Institute of Southeast Asian Studies.
- Cabin, R. J., et al. 2000. Effects of long-term ungulate exclusion and recent alien species control on the preservation and restoration of a Hawaiian tropical dry forest. *Conservation Biology* 14(2): 439-453.
- Carpenter, A., McEldowney, H. and Yent, M. 2006. *Historic Properties Identification and Field Report Upper Laupahoehoe and Humu'ula Sections Hilo Forest Reserve Reforestation Project*. Report prepared for Division of Forestry and Wildlife, Department of Land and Natural Resources.
- Chynoweth, M. W., et al. 2013. Biology and Impacts of Pacific Island Invasive Species. 9. *Capra hircus*, the Feral Goat (Mammalia: Bovidae). *Pacific Science* 67(2): 141-156
- Cole, R. J., et al. 2012. Vegetation Recovery 16 Years after Feral Pig Removal from a Wet Hawaiian Forest. *Biotropica* 44(4): 463-471.
- Cole, F.R., Loope L.L., Medeiros A.C., Howe C.E., and L.J. Anderson. 2000. Food habits of introduced rodents in high-elevation shrubland of Haleakala National Park, Maui, Hawaii. *Pacific Science* 54:313–329.



Cooray, R. G. and D. Mueller-Dombois. 1981. Island Ecosystems: Biological Organization in Selected Hawaiian Communities, Hutchinson Ross Publishing Co., Stroudsburg, PA, pp. 309-317.

County of Hawai'i. 2014 (draft plan under development). Hamakua Community Development Plan. Available at: <http://www.hawaiicountycdp.info/hamakua-cdp>

County of Hawai'i. 2005. General plan. Available at: <http://www.cohplanningdept.com/community-planning/general-plan/>

Cuddihy, L. W., J. A. Davis, and S. J. Anderson. 1982. A Botanical Survey of the Proposed Laupāhoehoe Natural Area Reserve, Hawai'i. Endangered Plant Species Program. Division of Forestry and Wildlife, Department of Land and Natural Resources, Hilo, Hawai'i.

Cuddihy, L.W., and C.P. Stone. 1990. Alteration of native Hawaiian vegetation: effects of humans, their activities and introductions. University of Hawaii Press, Honolulu, Hawaii

D'Antonio, C. M., et al. 2000. Variation in the impact of exotic grasses on native plant composition in relation to fire across an elevation gradient in Hawaii. *Austral Ecology* 25(5): 507-522.

Dunkell, D. O., et al. 2011. Runoff, Sediment Transport, and Effects of Feral Pig (*Sus scrofa*) Exclusion in a Forested Hawaiian Watershed. *Pacific Science* 65(2): 175-194.

Friday, J. B., Scowcroft, P.G., and A. Ares. 2008. Responses of native and invasive plant species to selective logging in an *Acacia koa*-*Metrosideros polymorpha* forest in Hawai'i. *Applied Vegetation Science* 11: 471-482.

Giambelluca, T.W., Nullet, M.A., and Schroeder, T.A. 1986. Hawai'i Rainfall Atlas, Report R76, Hawai'i Division of Water and Land Development, Department of Land and Natural Resources, Honolulu. vi + 267 p.

Giffin, Jon. 1978. Ecology of the feral pig on the island of Hawai'i. *Elepaio* 37 (12): 140-142.

Gillespie, Rosemary G. 1992. Distribution and diversity of an endemic lineage of invertebrate predators in the Natural Area Reserves of Hawai'i. Report to the Natural Area Reserves System.

Hawai'i Department of Land and Natural Resources (DLNR) Division of Forestry and Wildlife. 1983. (1) Abundance and Distribution of Koloa on Island of Hawai'i. (2) Movements, Survival, Reproductive Success and Habitat of Koloa on the Island of Hawai'i. Project # W-18-R-8. Job # R-III-H. July 1, 1981 to July 31, 1983.

Hawai'i Department of Land and Natural Resources Division of Forestry and Wildlife Natural Area Reserves System. 1989. Laupāhoehoe Draft Management Plan. Available at: <http://dlnr.hawaii.gov/ecosystems/files/2013/07/Laupahoehoe-Management-Plan.pdf>



Hawai‘i Department of Land and Natural Resources Division of Forestry and Wildlife. 2003. Management Plan for the Ahupua‘a of Pu‘u Wa‘awa‘a and the Makai Lands of Pu‘u Anahulu.

Hawai‘i Department of Land and Natural Resources Division of Forestry and Wildlife. 2005. Hawaii’s comprehensive wildlife conservation strategy. Available at: <http://dlnr.hawaii.gov/wildlife/cwcs/hawaii/>

Hawai‘i Department of Land and Natural Resources Division of Forestry and Wildlife. 2006. Final Environmental Assessment for the Hilo Forest Reserve Reforestation Project. Available at: http://oeqc.doh.hawaii.gov/Shared%20Documents/EA_and_EIS_Online_Library/Hawaii/2000s/2006-10-08-HA-FEA-HILO-FOREST-RESERVE-REFORESTATION.pdf

Hawai‘i Department of Land and Natural Resources Division of Forestry and Wildlife. 2007. Technical Report No. 07-01. Review of methods and approach for control of non-native ungulates in Hawai‘i.

Hawai‘i Department of Land and Natural Resources. 2012. The rain follows the forest - a plan to replenish Hawaii’s source of water. Available at: <http://dlnr.hawaii.gov/rain/plan/>

Hawai‘i Tropical Forest Recovery Task Force. 1994. Hawai‘i Tropical Forest Recovery Action Plan. Available at: <http://www.fs.fed.us/psw/ef/hawaii/documents/HITropicalForestRecoveryPlan.pdf>

Iwashita, D. K., et al. 2013. Coarse woody debris carbon storage across a mean annual temperature gradient in tropical montane wet forest. *Forest Ecology and Management* 291: 336-343.

Jacobi , J.D. 1989. Vegetation maps of the upland plant communities on the islands of Hawai‘i, Maui, Molokai, and Lanai. Cooperative National Park Resources Studies Unit Technical Report 68. University of Hawai‘i at Manoa and National Park Service.

Mack, M. C. and C. M. D'Antonio. 1998. Impacts of biological invasions on disturbance regimes. *Tree* 13(5): 195-198.

Maly, K. and O. Maly. 2006. Hilo Palikū - Hilo of the Upright Cliffs: A Study of Cultural-Historical Resources of Lands in the Laupāhoehoe Forest Section, Ahupua‘a of the Waipunalei-Mauluanui Region, North Hilo District, Island of Hawai‘i. Kumu Pono Associates, LLC.

Mueller-Dombois, D. 1980. The ‘ōhia dieback phenomenon in the Hawaiian rain forest. The recovery process in damaged ecosystems. J. Cairns, Jr. Ann Arbor, Ann Arbor Science Publishers Inc.

Nature Conservancy of Hawai‘i and member agencies of the Coordinating Group on Alien Pest Species (CGAPS). 2003. Silent Invasion.



- Nogueira-Filho, S. L., et al. 2009. Ecological impacts of feral pigs in the Hawaiian Islands. *Biodiversity and Conservation* 18(14): 3677-3683.
- Palmer, D. D. 2003. *Hawai'i's ferns and fern allies*. University of Hawai'i Press. Honolulu.
- Parham, J.E., G.R. Higashi, E.K. Lapp, D.K. Kuamo'o, R.T. Nishimoto, S. Hau, J.M. Fitzsimons, D.A. Polhemus, and W.S. Devick. 2008. *Atlas of Hawaiian Watersheds and Their Aquatic Resources, Island of Hawai'i*. Bishop Museum and Division of Aquatic Resources. 1262 p.
- Paton, P. W. C. 1981. The Koloa (Hawaiian Duck) on the Island of Hawai'i. *'Elepaio* 41(12):131-133.
- Price, J. S. M. Gon III, J. D. Jacobi, and D. Matsuwaki, 2007 Mapping Plant Species Ranges in the Hawaiian Islands: Developing a Methodology and Associated GIS layers. Hawai'i Cooperative Studies Unit Technical Report HCSU-008. University of Hawai'i at Hilo. 58 pp., incl. 16 Figures and 6 Tables.
- Pukui, Mary Kawena and Samuel H. Elbert. 1986. *Hawaiian Dictionary*. University of Hawai'i Press: Honolulu.
- Sasaki, D.M., L. Pang, H.P Minette, C.K Wakida, W.J. Fujimoto, S.J. Manea, R. Kunioka, C.R. Middleton. 1993. Active surveillance and risk factors for *Leptospirosis* in Hawai'i. *The American Journal of Tropical Medicine and Hygenics*. January 1993, 48(1):35-43.
- Scott, M.J., S. Mountainspring, F.L Ramsey, and C.B. Kepler. 1986. *Forest bird communities of the Hawaiian Islands: their dynamics, ecology, and conservation*. Studies in Avian Biology No. 9, Cooper Ornithological Society. Allen Press, Lawrence, Kansas.
- Scowcroft, P. G. and J. G. Giffin. 1983. Feral herbivores suppress mamane and other browse species on Mauna Kea, Hawaii. *Journal of Range Management*: 638-645.
- Scowcroft, P.G. and R.E. Nelson. 1976. Disturbance during logging stimulated regeneration of koa. USDA-FS Pacific Southwest Forest and Range Experiment Station. Research Note PSW-306.
- Shiels A.B. and D.R. Drake. 2011. Are introduced rats (*Rattus rattus*) both seed predators and dispersers in Hawaii? *Biological Invasions* 13:883–894
- Sherrod, David R., Sinton, John M., Watkins, Sarah E., and Brunt, Kelly M., 2007, Geologic map of the State of Hawai'i: U.S. Geological Survey Open-File Report 2007-1089. Available at: <http://pubs.usgs.gov/of/2007/1089/>
- Smith, C.W. and T. J. Tunison. 1992. Fire and alien plants in Hawaii: research and management implications for native ecosystems. *Alien plant invasions in native ecosystems of Hawaii: management and research*. Stone, C. P., C. W. Smith, and J. T. Tunison (eds.). Honolulu, Cooperative National Park Resources Studies Unit: 394–408



State of Hawai‘i. 2013. Office of Planning. State GIS database.

Stewart, Carolyn. 2010. Draft Mauna Kea Watershed Management Plan. Plan prepared for the Mauna Kea Watershed Alliance.

Stone, C.P. 1985. Feral ungulates in Hawai‘i: Toward controlling the adverse effects of introduced vertebrates. In Stone, C.P. and J.M. Scott (eds.). *Hawaii’s Terrestrial Ecosystems: Preservation and Management*.

Stone, C.P., Cuddihy, L.W., and J.T. Tunison. 1992. Responses of Hawaiian ecosystems to the removal of feral pigs and goats. Pp. 666-704 in C.P. Stone, C.W. Smith, and J.T. Tunison (eds.). *Alien Plant Invasions in Native Ecosystems of Hawai‘i: Management and Research*. University of Hawai‘i Cooperative National Park Resources Studies Unit, University of Hawai‘i Press, Honolulu.

Thaxton, J. M., et al. 2010. Native Species Regeneration Following Ungulate Exclusion and Nonnative Grass Removal in a Remnant Hawaiian Dry Forest. *Pacific Science* 64(4): 533-544.

Tomich, P.Q. 1986. *Mammals in Hawai‘i*. Second edition. Bishop Museum Press. Honolulu, HI. 375 pp.

Trujillo, E.E., Kadooka, C., Tanimoto, V., Bergfeld, S., Shishido, G. and Kawakami, G. 2001. Effective biomass reduction of the invasive weed species banana poka by *Septoria* leaf spot. *Plant Disease* 85: 357-361.

Tunison, J.T. and C.P. Stone. 1992. Special ecological areas: an approach to alien plant control in Hawai‘i Volcanoes National Park. Pp. 781-798 In C.P. Stone, C.W. Smith, and J.T. Tunison (eds.). *Alien Plant Invasions in Native Ecosystems of Hawai‘i: Management and Research*. University of Hawai‘i Cooperative National Park Resources Studies Unit, University of Hawai‘i Press, Honolulu.

United States Department of Agriculture Forest Service (USFS). 2007. Establishment Record for the Hawai‘i Experimental Tropical Forest within Hawai‘i County, Hawai‘i
Available at: <http://www.fs.fed.us/psw/ef/hawaii/documents/HETFEstablishmentRecord.pdf>

USFS. 2011. Final Environmental Assessment HETF Laupāhoehoe Construction Project.
Available at: http://www.hetf.us/page/projects_plans/pdf/HETF_Final_EA%20_08312011.pdf

United States Department of the Interior Geological Survey (USGS). 2005. Biocomplexity of introduced avian diseases in Hawai‘i: Threats to biodiversity of native forest ecosystems. Prepared by B.L. Woodworth, C. T. Atkinson, M. D. Samuel, D. A. LaPointe, P. C. Banko, and J. A. Ahumada. USGS Fact Sheet 2005-3139.
Available at: <http://biology.usgs.gov/pierc/Biocomplexity/Biocomplexity.pdf>.



USGS. 2006a. Feral cats: Too long a threat to Hawaiian wildlife. Prepared by S.C. Hess and P.C. Banko. USGS Fact Sheet 2006-3006. Available at:
http://biology.usgs.gov/pierc/Fact_Sheets/Feral_cats.pdf.

USGS. 2006b. Hawai'i forest bird interagency database project: Collecting, understanding, and sharing population data on Hawaiian forest birds. Prepared by T.K. Pratt, B.L. Woodworth, R.J. Camp, P.M. Gorresen. USGS Fact Sheet 2006-3013. Available at:
http://biology.usgs.gov/pierc/Native_Birds/Forest_birds.pdf.

USGS. 2006c. Feral pigs, introduced mosquitoes, and the decline of Hawai'i's native birds. Prepared by D.A. LaPointe. USGS Fact Sheet 2006-3029. Available at:
http://biology.usgs.gov/pierc/Fact_Sheets/Pigs_and_mosquitoes.pdf.

U.S. Fish and Wildlife Service (USFWS). 1978. Hawaiian Waterbirds Recovery Plan. USFWS, Portland, Oregon.

USFWS. 1984. Hawaiian Hawk Recovery Plan. Prepared under contract with C. R. Griffin. USFWS, Portland, Oregon.

USFWS. 1996. Big Island Plant Cluster Recovery Plan. USFWS, Portland, OR. 202+ pp. Available at: http://ecos.fws.gov/docs/recovery_plans/1996/960926a.pdf.

USFWS. 1998a. Big Island II: Addendum to the Recovery Plan for the Big Island Plant Cluster. USFWS, Portland, OR. 115 pp. Available at:
http://ecos.fws.gov/docs/recovery_plans/1998/980511a.pdf.

USFWS. 1998b. Recovery Plan for the Hawaiian Hoary Bat (*Lasiurus cinereus semotus*). USFWS, Portland, OR. 59 pp. Available at:
http://ecos.fws.gov/docs/recovery_plans/1998/980511b.pdf.

USFWS. 2003. Final designation and nondesignation of critical habitat for 46 plant species from the island of Hawai'i, HI. Federal Register Final Rule (68 FR 39623 39672). Available at:
http://ecos.fws.gov/docs/federal_register/fr4117.pdf

USFWS. 2006. Revised Recovery Plan for Hawaiian Forest Birds. Region 1, Portland, OR. 622 pp. Available at:
http://ecos.fws.gov/docs/recovery_plan/060922a.pdf

Wagner, W. L., Herbst, D. R., and Sohmer, S.H. 1999. Manual of the flowering plants of Hawai'i. University of Hawai'i Press, Bishop Museum, Honolulu.

Warner, Richard E., 1956 – 1969. Ecological investigation on the Hawaiian pig (1959); Ecology of the Wild Pig (1963, 1963); and Feral Game Mammal Survey (1969). State of Hawai'i, Division of Fish and Game, Project # W-r-R10; Project # W-5-R13; Project # W-5-R14; and Project # W-5-R-20.



Appendix A – Other Available Resources

The following resources related to Laupāhoehoe Forest are available on-line at:

<http://dlnr.hawaii.gov/ecosystems/nars/reserves/hawaii-island/laupahoehoe/>

- 1) *Hilo Palikū - Hilo of the Upright Cliffs: A Study of Cultural-Historical Resources of Lands in the Laupāhoehoe Forest Section, Ahupua‘a of the Waipunalei-Mauluanui Region, North Hilo District, Island of Hawai‘i* (Maly and Maly 2006)
- 2) Laupāhoehoe Plant Species List



Appendix B – Biosecurity Protocols



HETF BIOSECURITY



What is Biosecurity and why is it important?

Biosecurity is a set of precautions that aim to prevent the introduction and spread of harmful organisms. These may be pests, pathogens or invasive species. Biosecurity measures are the practical steps designed to minimize the risk of introducing or spreading pests and diseases. Control of invasive plants and animals is a high priority for all resource managers in Hawai'i. A large amount of personnel time and resources are expended each year to battle invasive species. Biosecurity is important because invasive species have significant negative impacts to resources - including native species, ecosystem function (e.g. nutrient cycling), ecosystem services (e.g. ground-water recharge, prevention of erosion, reef sedimentation, etc.), ecosystem structure (distribution and abundance of species), etc. New plants and animals arrive in the islands on a continual basis and natural resource managers must focus on preventing their spread into natural areas. Visitors, volunteers, contractors, researchers and staff are potential vectors of invasive species. The origin of these pests can be the mainland, another island in the Hawaiian archipelago or even a weedy or urban area of the same island.

Described below are likely routes of invasive species introduction and procedures that should be followed to minimize or eliminate introduction and spread of invasive plants and animals in the Hawai'i Experimental Tropical Forest.

Required HETF Biosecurity Measures:

1. Inspect field gear and equipment BEFORE going into the field. Three categories of pests that merit attention are weeds/seeds, insects (especially ants and cockroaches), and slugs. Clods of dirt or mud can hide these hitchhikers. If materials have been in storage, be aware that arthropods can lay eggs within equipment. It is necessary to inspect and clean all: footwear (treads, laces, boot tongues); socks; pant legs; pockets, jackets, raingear (pockets and cuffs); food-stuffs; tools; packs (folds, pockets, Velcro); tents; hammocks; tarps; helicopter sling nets; wood and building materials; fencing materials; bottoms of plastic buckets; tool bags; and other containers or equipment.

Soil, cinder and potted plants are also major invasive species transporters. Ensure soil and cinder are sterile and free of insects. Potted plants should be visibly weed-free, pest-free, and pathogen-free, anything suspicious should not be taken into the field.

Preferably a different set of field gear should be used for each field site. If this is not possible, gear should be thoroughly cleaned before going into the field. All gear (personal or field) must be free of any plant, animal, or earthen materials. Appropriate methods for cleaning include: water and hose, brush, clean rag, knife edge, bleach rinse and/or insecticide. Cleaning should be done away from the forest in a designated area with a nearby receptacle for disposal. Laundering of washable items should be done after each trip. This is particularly important when you have come from working on different islands or different parts of the island, as each area may have its own assemblage of pests that must be prevented from spreading to new locations. If items cannot be cleaned, they should not be taken into the forest.

2. Prepare a checklist of items to be inspected before any extended field operations or camping trips. Note all materials that will accompany staff during these operations. On prep day, an assigned person should inspect those supplies, concentrating on those most susceptible in aiding accidental introductions. Items having the potential for carrying unwanted species: helicopter sling nets, wood, building materials, fencing material, plastic buckets, cardboard boxes (roaches and ants!), open food and water containers, tool bags, backpacks, raingear and tents. Methods for cleaning articles range from using brushes, tweezers, bleached rags, or insecticides. Have these items on hand. If contaminated supplies are found, they must be disposed of away from the HETF.

3. Avoid carrying weed seeds from an infested part of the forest to the pristine areas. If routes of transit go through invasive weed belts, be conscious of the potential for spread. A safeguard could be having a brush on hand to clean shoe soles at the boundary of such a site. This is especially a concern when performing priority weed work where there is a large seed bank. When working in a different area, be very careful to avoid spreading pests by inspecting and cleaning gear before and after each excursion into the field. It is important to use completely different gear on each island or area.

4. Keep field vehicles clean. Clean field gear can easily be compromised by using a dirty truck. The inside of field vehicles should be vacuumed and pickup truck beds swept out regularly, especially if used off-road. When cleaning vehicles it is important to pay particular attention to the inside lip of bidders, seats, floors, dashboards, door jambs, tires (especially treads) and the undercarriage. All mud should be hosed off to avoid transporting weed seeds.

Continued on next page



L-R: Faya (*Morella faya*), Miconia (*Miconia calvescens*), Smoke bush (*Buddleja madagascariensis*), Cotoneaster (*Cotoneaster pannosus*) and Pampas grass (*Cortaderia jubata*). ALL are invasive.

Photo credits: Forest and Kim Starr



5. **Pack out your trash and unused foodstuffs.** Something as harmless as an apple core or banana peel could result in dire consequences out in the field. Discarded foods can become food for invasive insects, like ants; seeded fruits (apples, pear, avocado, peach, etc.) could germinate, as certain seeds have a strong potential to naturalize. Many foods carry unseen insects, fungus, bacteria and parasites that could affect native flora and fauna, Hawai'i has many closely related species that could be negatively affected by leaving behind foodstuffs.

6. **Become acquainted with invasive species in Hawai'i** and their varying status regarding particular localities. Learn which of these species is localized to your area and which to be on the alert for that are established in other areas. Take inventory of weeds that occur along trails and byways, and pay attention to any unusual changes or additions to these. There is plenty of literature on the identification of weedy plants that will provide current status throughout the various islands. Start with www.hear.org.

7. **Educate visitors to these protocols.** This includes other researchers, visitors, or volunteer groups. A very high probability exists that much of these visitors' gear (primarily packs and footwear) has been to other areas in Hawai'i that have infestations of alien species, and they could act as dispersal agents. People who share our appreciation of Hawai'i's native forests will not be insulted by these procedures if related firmly yet politely, but rather will be impressed with their conscientious implementation.

New Invasive Species Detection

Sightings of new invasive species within the HETF or existing high risk species seen in previously un-infested areas should be reported to the:

Hawai'i Experimental Tropical Forest
(808) 443-5931
hawaii_experimental_tropical_forest@fs.fed.us

Please provide the following information in your report: 1) Your name and contact information, 2) date you observed the species, 3) species name and/or detailed description, 4) approximate location (latitude/longitude, UTM, or other approximate estimation) of the organism, and 5) a rough estimate of the overall size of the area infested (if stationary).

There are additional invasive species of serious concern to land managers that are present in adjoining areas but have not yet been detected in HETF Units. It is a high priority to prevent the establishment of these species, which include but are not limited to:



Fire Thorn (*Pyracantha angustifolia*)

Iaupāhoehoe

- Miconia (*Miconia calvescens*)
- Faya (*Morella faya*)
- Gorse (*Ulex europaeus*)
- Firethorn (*Pyracantha angustifolia*)
- Cotoneaster (*Cotoneaster pannosus*)
- New Zealand flax (*Phormium tenax*)
- Indonesian cinnamon (*Cinnamomum burmanii*)
- Red Robin (*Photinia davidiana*)
- Ants (especially Little Red Fire Ants)

Pu'u Wa'awa'a

- Cotoneaster (*Cotoneaster pannosus*)
- Pampas Grass (*Cortaderia jubata* and *selloana*)
- Pereskia (*Pereskia aculeata*)
- Rubbervine (*Cryptostegia madagascariensis*)
- Smokebush (*Buddleja madagascariensis*)
- Wax Myrtle (*Morella cerifera*)
- Barbados Gooseberry (*Pereskia aculeata*)
- Jerusalem Thorn (*Parkinsonia aculeata*)
- Axis Deer (*Axis axis*)
- Ants (especially Little Red Fire Ants)

KNOWN AND POTENTIAL RISKS

FOR INFORMATION ON CURRENT PEST SPECIES THAT SHOULD BE REPORTED IF SEEN VISIT:

<http://www.reportapest.org/pestlist.htm>



L-R: Gorse (*Ulex europaeus*), Pereskia (*Pereskia aculeata*), Axis Deer (*Axis axis*), Rubbervine (*Cryptostegia madagascariensis*) and Jerusalem Thorn (*Parkinsonia aculeata*). ALL are invasive. Plant photo credits: Forest and Kim Starr. Axis deer photo credit: [daemonsandanalyses.tumblr.com](https://www.tumblr.com/daemonsandanalyses)

Other invasive species may be added to the priority invasive species list if monitoring shows their range and abundance increasing in native ecosystems targeted for management.



Appendix C - List of Research Projects to Date

Studies are grouped by total project years (one, two, or 3 and greater), then by project start date, and last name.

Project Description	Name	Affiliation	Project Start Date	Project End Date	Total Years of Project
Comparative nutritive values of traditional and exotic foraging substrates for upper elevation forest birds.	Kapono, M.	UH-Hilo	2007	2008	1
An assessment of tropical ecosystem dynamics in response to climate variability using long-term satellite data records.	Park, S.	UH-Hilo	2007	2008	1
Sprouting vs. seeding in Hawaiian wet, mesic, and dry forests.	Busby, P.	Stanford University	2008	2009	1
Objective quantification of plumage coloration of a Hawaiian honeycreeper (<i>Hemignathus virens</i>) along environmental gradations of biogeography: does variation exist between sub-populations?	Gaudioso, J.	UH-Hilo; USGS-PIERC	2008	2008	1
Study of the molecular evolution of arthropods - phylogenetic study.	O'Grady, P.	UC-Berkeley	2008	2009	1
Study of alien snails, survey of native snails	Yeung, N	UH-Manoa	2008	2009	1
Examining the impacts of strawberry guava on Native biodiversity in Hawaiian forest	Giardina, C.	USDA Forest Service	2009	2010	1
Remote sensing across multiple spatial scales for use with calibration of satellite sensors.	Goodenough, D.G.	Canadian Forest Service Natural Resources	2009	2010	1
Soil Survey		USDA-NRCS	2009	2010	1
DNA sequencing of endemic Hawaiian Drosophila.	Price, D.	UH-Hilo	2012	2013	1
Climatic influences on Lycopsid and fern leaf physiognomy	Benca, J.	UC-Berkeley	2013	2014	1
Community assembly & diversification of Hawaiian Arthropods	Gillespie, R.	UC-Berkeley	2013	2014	1
Experimental cultivation of simple rare ferns in common garden conditions.	Husby, C.	Montgomery Botanical Center	2013	2014	1
Epiphytes as an indicator of climate change in Hawaii	Kettwich, S.	UH-Hilo	2013	2014	1
Assessment of mosquito-borne avian disease risk in non-breeding habitat for foraging iiwi adjacent to Hakalau Forest NWR	LaPointe, D.	USGS-PIERC	2013	2014	1
Assessing the <i>Scotorythra paludicola</i> outbreak on koa: population abundance, rates of parasitism and patterns of spread	Peck, R.	UH-Hilo	2013	2014	1
UH Hilo Geography and Environmental Studies Senior Seminar	Price, J.	UH-Hilo	2013	2014	1



Project Description	Name	Affiliation	Project Start Date	Project End Date	Total Years of Project
An assessment of Hawaiian tropical ecosystem dynamics in response to climate variability using long-term satellite data records.	Miura, T.	UH-Manoa	2007	2009	2
Passive study of vocalizations of Hawaiian hoary bat for evaluation of conservation status.	Bonaccorso, F.	USGS-PIERC	2008	2010	2
Developing a DNA barcoding method to be used as a system of species identification	Magnacca, K.	UH-Hilo	2009	2011	2
Do expected evolutionary trade-offs in enzyme activities manifest at the level of microbial community function?	Bradford, M	Yale University	2010	2012	2
Inventory of Hawaii's Forests	Hughes, F.	USDA-USFS	2011	2013	2
Understanding the genetic variation in <i>Acacia koa</i> across environmental gradients	Michler, C.	USDA-USFS; Purdue University	2012	2014	2
Quantifying the effects of ungulate and vegetation on the hydrology of Hawaiian tropical forests	MacKenzie, R.	USDA-USFS	2012	2014	2
Long-term monitoring of forest and climate inventory	Cordell, S.	USDA-USFS	2007	ongoing	3+
Forest architecture, Carbon dynamics, and climate change interactions: linking field and remote sensing along temporal and spatial gradients.	Broadbent, E.	Stanford University; Carnegie Institute of Washington	2008	2012	3+
Investigating productivity of koa forests on different substrates and climate zones. Long-term plots.	Friday, J.B.	UH-Manoa-CTAHR	2008	2012	3+
Sources and fates of nutrients on a substrate age gradient across the Hawaiian archipelago and their consequences for forest dynamics.	Vitousek, P.	Stanford University	2008	ongoing	3+
Assessing Forest Structure, community composition, diversity, carbon mass, and biomass on a landscape scale in the HETF	Hughes, F.	USDA-USFS	2009	2015	3+
Experimental Test of the impacts of rising temp on C input, allocation, and loss in model forests.	Litton, C.	UH-Manoa	2009	ongoing	3+
Hawaiian Hoary Bat habitat occupancy, reproduction and diet	Bonaccorso, F.	USGS-PIERC	2010	2014	3+
Adaptive Radiation in Hawaiian Spiders	Gillespie, R.	UC-Berkeley	2011	2014	3+
Forest Disease Monitoring for Rust Disease affecting Ohia Lehua	Yeh, A.	Contractor for DLNR/DOFAW Forest Health	2011	ongoing	3+
Impacts of strawberry guava management across a density gradient	Johnson, T.	USDA-USFS	2013	2028	3+



