



Terrestrial Invertebrates

Bees and wasps Order Hymenoptera

ORDER INCLUDES:

17 Native Families

60 Native Genera

650+ Native Species

650+ Endemic Species

GENERAL INFORMATION: The order Hymenoptera is large and diverse, and is best known because of the social behavior of ants, bees, and wasps. Hawai'i's native Hymenoptera fauna, however, comprises non-social bees and wasps and does not include any native ants. The most speciose genus is *Sierola* (Bethyridae) with 180 described and an estimated 400–500 total species. Several species in each of the genera *Hylaeus* (Colletidae), *Ectemnius* (Sphecidae), and *Nesodynerus* (Vespidae), including *H. chlorostictus*, *H. difficilis*, *H. pubescens*, *E. nesiotus*, *E. polynesianalis*, *N. peles*, and *N. scoriaceus*, are common and relatively abundant. However, the status of many other species in these genera, as well as species in the genera *Enicospilus* (Ichneumonidae), *Sclerodermus* (Bethyridae), *Sierola* (Bethyridae), is unknown. As elsewhere, Hawaiian bees and wasps have played an important role in keeping other insects, particularly herbaceous species, in check, and in pollinating plants. The native *Hylaeus*, or yellow-faced bees, are important pollinators for many native plants. Most of the native wasps are arthropod predators or parasites, often specific to particular taxa. For example, species in the family Mymaridae parasitize small insects and insect eggs; species in the family Encyrtidae prey on scale insects (Homoptera); species in the family Eucoilidae lay their eggs on fly (Diptera) pupae, and species in the families Vespidae and Bethyridae prey on Lepidoptera caterpillars. As an example of the diversity of Hawaii's hymenopterans, two genera are briefly outlined. The 63 species in the bee genus *Hylaeus* occur on all the Main Hawaiian Islands (MHI) and Nihoa. They nest in hollow stems, holes in trees, under bark, in crevices, or in burrows in soil. Potential threats include non-native bees (*Ceratina* spp.) found in the native coastal habitats used by *Hylaeus* species, and competition with the European honeybee (*Apis mellifera*) for nectar and pollen. Confirmed threats include introduced ants (Formicidae) which compete with *Hylaeus* for nesting sites, and the big-headed ant (*Pheidole megacephala*) and Argentine ant (*Linepithema humile*) which prey on the native bees. Since *Hylaeus* bees pollinate native plants, their loss would be detrimental to recovery of native plants. Wasps in the genus *Sierola* are found throughout the Indo-Pacific region, but over 90 percent of the known species are endemic to Hawai'i. They are small, black wasps found primarily in wet and mesic forest. Fullway (1920) described 171 species, with 119 from O'ahu and 44 from the island of Hawai'i. Possibly hundreds of species remain to be described, and it is not uncommon to find greater than ten morphospecies (i.e., species established solely on morphological characteristics) at a single site. These species are difficult to collect, and many morphospecies are found in very low numbers; this may be an artifact of collection methods or due to their rarity. Similar to other wasps, females find a Lepidoptera larva, sting and paralyze it, and lay an egg on it. Once the egg hatches the larva feeds on the caterpillar.

DISTRIBUTION: Native Hymenoptera are found on all the MHI and Laysan and Nihoa in the Northwestern Hawaiian Islands.

ABUNDANCE: Unknown. A lack of systematic surveys prevents any population estimate. However, the loss of native habitats likely means that species within the order are declining.

LOCATION AND CONDITION OF KEY HABITAT: Hymenopterans occur in most terrestrial habitats.

THREATS:

- Competition with non-native wasps and ants.
- Predation by non-native insects.
- Loss or degradation of habitat, especially the loss of native host plants for bees.

CONSERVATION ACTIONS: The goals of conservation actions are not only to protect current populations and key breeding habitats, but also to establish additional populations, thereby reducing the risk of extinction. In addition to common statewide and island conservation actions, specific management directed toward hymenopterans should include the following:

- Conduct surveys to determine the distribution and abundance of known hymenopterans and to document and identify new species.
- Preserve, maintain, and restore habitat for existing populations.

MONITORING: Continue monitoring the status of known populations.

RESEARCH PRIORITIES:

- Conduct systematic and taxonomic assessments of poorly known taxa.
- Conduct studies to document the biology, habitat requirements, and life history of endemic species.
- Refine methods for conducting quantitative survey.

References:

Daly HV, and Magnacca KN. 2003. Insects of Hawai'i Vol 17: Hawaiian *Hylaeus* (Nesoprosopis) bees. University of Hawai'i Press, Honolulu.

Fullaway DT. New species of *Sierola* with explanatory notes. Bishop Museum Occasional Papers 7(7):57-159.

Howarth FG, and Mull WP. 1992. Hawaiian insects and their Kin. University of Hawai'i Press, Honolulu.

Nishida GM, editor. 2002. Hawaiian Terrestrial Arthropod Checklist, 4th edition. Biological Survey, Bishop Museum, Honolulu, Hawai'i.

Takumi RL. 1999. A Systematic Review of the Ectemnius (Hymenoptera: Sphecidae) of Hawai'i, University of California Press, Berkeley.

Zimmerman EC. 2001. Insects of Hawai'i: Volume 1 Introduction. University of Hawai'i Press, Honolulu.