



Photo: DOFAW

## Forest Birds

# Kaua'i 'elepaio

*Chasiempis sclateri*


### SPECIES STATUS:

State Recognized as Endemic  
NatureServe Heritage Rank G2 – Imperiled  
IUCN Red List Ranking – Vulnerable

**SPECIES INFORMATION:** The Kaua'i 'elepaio is a small, adaptable monarch flycatcher (Family: Monarchiade) endemic to the island of Kaua'i. 'Elepaio also occur on the islands of Hawai'i (*C. sandwichensis*) and O'ahu (*C. ibidis*); the latter is federally listed as endangered. Adults have a dark grayish brown crown and back and white underparts with a rusty wash on the upper breast. The bird's name is derived from its primary song which is a shrill whistle given only by males. Little detailed life history information is available from Kaua'i. On Hawai'i and O'ahu, they use virtually all available substrates for foraging, including the ground, logs, rock crevices, snags, and all parts of tress. Equally diverse in the use of foraging maneuvers, 'elepaio capture a wide range of arthropod prey by flycatching, gleaning while perched or hovering, and direct pursuit; foraging maneuvers vary depending on plant species from which prey is being captured and habitat type. For Kaua'i, there is no information on plant species used, although 'ohi'a (*Metrodiseros polymorpha*) is likely. On the island of Hawai'i, pairs remain together throughout the year and long-term pair bonds are common. Unlike Hawaiian honeycreepers, both males and females participate almost equally in all aspects of rearing. Finely woven, cup nests are built in a variety of native and non-native trees. Clutch size is usually two and second nests are attempted, often while fledglings from first are still being fed. Young are fed by parents for at least a month, but remain on their natal territory for up to ten months, which may allow young birds to hone their foraging skills.

**DISTRIBUTION:** Widely distributed above 600 meters (2,000 feet) elevation, but most common above 1,100 meters (3,600 feet) elevation on the Alaka'i Plateau. Original range likely included all forested regions of Kaua'i.

**ABUNDANCE:** In 1984, the island-wide population was estimated at 40,000 birds. The 2000 Kaua'i Forest Bird Survey estimated the population in the Alaka'i and Kōke'e region at nearly 25,000 birds, with no change in the population since 1973. Densities peak in 'ohi'a forest between 1,300 and 1,500 meters (4,500 - 5,000 feet).



**LOCATION AND CONDITION OF KEY HABITAT:** Most common in dense wet 'ōhi'a forests above 1,100 meters (3,600 feet) elevation; uncommon in the drier forests of Waimea Canyon, the Nā Pali coast, and the wet bogs of Wai'ale'ale. 'Elepaio are found in some low elevation valleys with a mix of native and non-native tree species. Common non-native trees in this habitat include Java plum (*Syzygium cumini*), coffee (*Coffea arabica*), kukui (*Aleurites moluccana*), and *Albizia lebeck*. Habitat quality varies considerably. The core of their range, the Alaka'i Plateau and Kōke'e State Park, is managed by the State of Hawai'i as Forest Reserve, Wilderness Preserve, and State Park lands.

**THREATS:** Kaua'i 'elepaio are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For Kaua'i 'elepaio, the following are of particular concern:

- Disease. Avian pox is known to reduce both nesting success and adult survival. On O'ahu, annual survival and reproductive success of birds with active pox lesions are lower compared to healthy birds; no information on the effect of avian malaria.
- Predation. On O'ahu, predation by black rats have been implicated in the loss of nests and death of adult females. Rat control in these populations resulted in large increases in nest success and in the survival of adult females. Other predators, including cats, pueo, and barn owls, occur throughout the forests of Kaua'i.

**CONSERVATION ACTIONS:** Kaua'i 'elepaio likely benefited from management actions to conserve other endangered forest birds including establishment of the Alaka'i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, studies of disease and disease vectors, and public education efforts featuring Kauai's endangered forest birds. In addition to these efforts, future management specific to the Kaua'i 'elepaio may include the following:

- Eradicate or control rats, feral cats, and barn owls in the Alaka'i Wilderness Preserve.
- Prevent the introduction of the small Indian mongoose and other possible predators.
- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.

**MONITORING:** Continue forest bird surveys and habitat monitoring.

**RESEARCH PRIORITIES:** Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquitos. Research priorities specific to the Kaua'i 'elepaio include life history studies to quantify population structure, dispersal patterns, survivorship, nesting phenology, and success of this poorly known subspecies.

**References:**

Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka'i swamp, Kaua'i. *Conservation Biology* 18:716-725.

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: [www.iucnredlist.org](http://www.iucnredlist.org). (Accessed May 2015).

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service. 428 pp.

VanderWerf EA. 1998. 'Elepaio (*Chasiempis sandwichensis*). In The Birds of North America, No. 344 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.