



Photo: Jim Denny

Forest Birds

‘Akeke‘e

Loxops caeruleirostris

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic


NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Critically Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

Kauai Islandwide Recovery Plan – USFWS 2021

SPECIES INFORMATION: The ‘akeke‘e, or Kaua‘i ‘ākepa, is a small, slightly sexually dichromatic, insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to Kaua‘i. Adult males and females are greenish above and yellow below with a yellow crown and a black mask; females are slightly duller than males. Unlike the similar Kaua‘i ‘amakihi (*Hemignathus kauaiensis*), the ‘akeke‘e’s bill is conical. Although not visible in the field, the lower mandible of the ‘akeke‘e is slightly bent to one side which results in the mandible tips being offset; a characteristic shared with the ‘ākepa (*L. coccineus*). The ‘akeke‘e uses its bill to pry open ‘ōhi‘a (*Metrosideros polymorpha*) leaves and flower buds in search of arthropods, primarily spiders, psyllids, and caterpillars. The species is an ‘ōhi‘a specialist and rarely even perches on other trees or shrubs. Its methodical probing of leaf buds is distinctive and can be used to identify the species. ‘akeke‘e are most often observed in pairs or family groups. A study of 22 ‘akeke‘e nests monitored from 2012-2017 found a preference for nesting in small terminal branches of ‘ōhi‘a trees at higher elevation and taller canopy heights, as well as a negative relationship with slope (Fricker et al. 2021). In a sample of eight nests monitored in 2012 and 2013, mean nest height was 11.1 ± 2.3 m, and all nests were also located in the small terminal branches of ‘ōhi‘a (2015). On one occasion a male was observed helping the female build the nest, but on all other occasions, only the presumed female has built the nest (few ‘akeke‘e are color banded so it is often difficult to determine which bird is the female unless she is incubation). The species has been observed to reneest and double brood in at least one instance. Multiple adults and subadults attend nests.



DISTRIBUTION: Formerly found in native forests of the Alaka'i swamp, upper Waimea, and Kōke'e regions mostly above 1,000 meters (3,280 feet) elevation but are now restricted only to the Alaka'i Wilderness Preserve (DOFAW, unpublished data) at a mean elevation of 1298.7m (± 50.6 m; Fricker et al. 2021). Although historically widespread, 'akeke'e apparently did not occur at lower elevations. Occupancy rates in 2012 increased from west to east along the plateau ($\psi = 0.03 \pm 0.10$ to $0.53 \pm$

0.33) but were low throughout its range (Behnke et al. 2016). The estimated suitable habitat range based on surveys from 2012-2017 was 58km² (Fricker et al 2021), which represented a small fraction of the former range. Based on data from 2023 and 2024, the range is now much smaller at approximately 1234 ha (3049 ac; KFBRP, unpublished data), and the species occurs primarily at Halepa'akai and Mohihi .

ABUNDANCE: In the early 1970s the island-wide population was estimated at $5,066 \pm 1,680$ (SE) individuals. The Kaua'i Forest Bird Survey in 2012 suggested that the population was rapidly declining, especially in the periphery of its range; the 2012 population estimate was 945 (95% confidence interval: 460 to 1,547) individuals (Paxton et al. 2016). Surveys in 2018 suggested that the population was relatively stable (Paxton et al. 2020), however the 2023 surveys showed ongoing declines to 765 (95% confidence interval: 297 to 1285) individuals (U.S. Geological Survey unpublished data). Preliminary data from occupancy surveys in 2024 suggest further declines in abundance to fewer than 100 individuals, and ongoing range contraction (Kauai Forest bird Recovery Project, unpublished data). Densities are highest in the interior of the Alaka'i Wilderness Preserve.

LOCATION AND CONDITION OF KEY HABITAT: Occurs above 600 meters (1,950 feet), although populations are densest above 1,100 meters (3,600 feet), in lowland mesic and wet forests dominated by 'ōhi'a, koa (*Acacia koa*), 'ōlapa (*Cheirodendron trigynum*), and lapalapa (*C. platyphyllum*). Most of the current range the southern Alaka'i Wilderness Preserve; the species no longer occurs in Kōke'e State Park or the northern or western Alaka'i. Occupancy is positively correlated with canopy height and maximum 'ōhi'a diameter at breast height.

THREATS:

- Habitat degradation. The spread of non-native plants and degradation by ungulates may reduce habitat suitability. The correlation of occupancy with large tree metrics suggest that damage done by two hurricanes may also limit distribution and abundance.

- Disease. ‘Akeke‘e is highly susceptible to mosquito-borne avian malaria. Two of 11 ‘akeke‘e have tested positive for malarial antibodies since 2011; since malaria is established on the Alaka‘i Plateau, the most likely explanation for this result is high mortality after infection with malaria. Disease, habitat degradation, and predation are the most probable causes of population declines in this species in the last decade.
- Competition. Non-native insects, especially yellow-jackets (*Vespula pensylvanica*) and ants (*Linepithema humile*), may compete with or prey on the native arthropods on which ‘akeke‘e feed. The role of non-native insects in native Hawaiian forests is unclear.
- Predation. Rats (*Rattus* spp.), cats (*Felis catus*), Hawaiian short-eared owls (*Asio flammeus sandwichensis*), and barn owls (*Tyto alba*) occur throughout the forests of Kaua‘i and may prey on young and adults. Rat predation on a female ‘akeke‘e and her young was observed at Halepa‘akai.
- Small population dynamics. The observed hatching failures may indicate genetic issues associated with small population sizes.

CONSERVATION ACTIONS: ‘Akeke‘e likely benefited from actions to conserve other endangered forest birds including the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, habitat monitoring, studies of disease and disease vectors, control of feral ungulates through public hunting and fencing, suppression of rats on two A24 trapping grids over approximately 150 ha (470 ac), and public education featuring Kauai’s endangered forest birds. In 2015-2018, an effort to form a captive flock was initiated by collecting ‘akeke‘e eggs from the wild. Twenty-six eggs were harvested from ten nests; most eggs hatched but many chicks did not survive to fledging, resulting in a 10 adult ‘akeke‘e. The flock has since suffered more attrition, but successful breeding was achieved in 2023 and there are currently seven ‘akeke‘e in human care. In addition to these efforts, future actions specific to the ‘akeke‘e may include the following:

- Resume collections of ‘akeke‘e to increase the size of the captive flock
- Aggressively control ungulates to improve habitat quality, facilitate the recovery of degraded habitat, and potentially reduce breeding habitat for mosquitoes. Control of non-native plants should be part of forest restoration efforts.
- Eradicate or manage mosquito breeding habitat on the Alaka‘i Plateau
- Suppress mosquito populations through releases of incompatible or sterile mosquitoes
- Investigate options to eradicate mosquitoes, including use of genetic manipulation.
- Suppress/control rats, feral cats, and barn owls from the Alaka‘i Wilderness Preserve, Na Pali-Kona Forest Reserve, and Kōke‘e State Park
- Prevent the introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other predators.
- Prevent the spread of Rapid ‘ōhi‘a Death on the Alaka‘i Plateau
- Conduct public outreach and education
- Continue protection and management of wildlife sanctuaries and refuges
- Collect genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning
- Explore translocating birds from Kaua‘i to other locations

MONITORING: Continue Forest bird surveys and habitat monitoring. Conduct species-specific monitoring for ‘akeke‘e (e.g., occupancy surveys with play back, territory monitoring, use of ARUs) that take into account the inability of the HFBS to accurately track populations of rare species

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the ‘akeke‘e include the following:

- Investigate techniques for improving survival and reproduction of the captive flock
- Conduct PVAs for both wild and captive flocks
- Conduct life history studies to quantify the population structure, dispersal patterns, survivorship, nesting phenology, and success of this poorly known species.
- Continue to assess the species’ susceptibility to avian malaria and avian pox.
- Determine sources of mosquitoes and investigate appropriate methods of mosquito control.
- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps)
 - Determine the effects of recently established non-native insects on native arthropods, especially those that are part of the species’ diet.

References:

- Atkinson, CT, RB Utzurrum, DA LaPointe, RJ Camp, LH Crampton, JT Foster, and TW Giambelluca. 2014. Changing Climate and the Altitudinal Range of Avian Malaria in the Hawaiian Islands - an Ongoing Conservation Crisis on the Island of Kaua‘i. *Global Change Biology* 20, 2426–2436, doi: 10.1111/gcb.12535.
- Behnke, L. 2014. Habitat use and conservation implications for Akikiki (*Oreomystis bairdi*) and Akekee (*Loxops caeruleirostris*), two endangered Hawaiian honeycreepers. M.S. Thesis. Colorado State University, Fort Collins, CO.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- Fricker GA, Crampton LH, Gallerani EM, Hite JM, Inman R, Gillespie TW. 2021. Application of lidar for critical endangered bird species conservation on the island of Kauai, Hawaii. *Ecosphere* 12(6).
- Hammond, RL, Crampton, LH, and Foster, JT. 2015. Breeding biology of two endangered forest birds on the island of Kaua‘i. *Condor* 117: 31-40.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Lepson JK, Pratt HD. 1997. ‘Akeke‘e (*Loxops caeruleirostris*). In *The Birds of North America*, No. 295 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.

- Paxton ER, Camp MG, Crampton LH, Leonard DL, VanderWerf EA. 2016. Collapsing avian community on a Hawaiian island. *Science Advances* 2:e1600029.
- Paxton ER, Brinck KW, Crampton LH, Hite J, Constantini M. 2020. 2018 Kaua‘i forest bird population estimates and trends. Technical Report Hawai‘i Cooperative Studies Unit. University of Hawai‘i at Hilo.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.
- U.S. Fish and Wildlife Service. 2021. Kauai Islandwide Recovery Plan. Portland, (OR): U.S. Fish and Wildlife Service. <https://www.fws.gov/node/264716>



Photo: UH EECB

Forest Birds

‘Akiapōlā’au

Hemignathus munroi

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered


State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The ‘akiapōlā’au is a stocky Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Hawai‘i and most famous for their specialized bills, which have a long, decurved upper mandible and a short woodpecker-like lower mandible. Adult males have a bright yellow head and underparts, yellow-green back and wings, and a small, black mask. Adult females are olive above with grayish-yellow to yellow underparts. Males are larger than females and have longer bills. They often join mixed species foraging flocks; the constituent species vary depending on habitat. ‘Akiapōlā’au are mainly insectivorous, with Lepidoptera larva, spiders, and beetle larva being the most important prey items; rarely takes nectar but takes sap from holes it excavates in ‘ōhi‘a (*Metrosideros polymorpha*) trees. Most frequently, creeps along lichen covered and dead branches of koa (*Acacia koa*), kōlea (*Myrsine lessertiana*), māmane (*Sophora chrysophylla*), and naio (*Myoporum sandwicense*) trees tapping branches with their lower mandible to locate prey. Once a food item is located, lower mandible is used similar to that of a woodpecker bill to chisel open a hole. The upper mandible is then used to fish out the prey item. Upper mandible also used to probe natural cracks and crevices. Breeding has been documented year-round, although most activity occurs from May to August with a second peak in November. The species’ open cup nest is most often placed in ‘ōhi‘a trees. Clutch size is usually one, rarely two, and females perform all incubation and brooding. Males provide females and nestlings with the majority of food. Only one fledgling is produced per year, and a long period of parental dependency, usually four to six months, is typical,



but can be as long as 18 months. Family groups consisting of hatch-year and second-year young have been observed. This species is characterized by low annual productivity.

DISTRIBUTION: Occurs in three disjunct populations between 1,500 and 2,000 meters (4,875–6,500 feet) elevation on the Island of Hawai‘i. Original range likely included all forested areas of the island, however now extirpated from their range in Kona and sub-alpine Maunakea.

ABUNDANCE: The Hawaiian Forest Bird Survey (1976-79, 1983), estimated the population at $1,500 \pm 400$ (95% confidence interval). Surveys conducted between 1990 and 1995 estimated the population at 1,109-1,217 birds and most recent analysis puts the population closer to 1,900. Populations at Hakalau NWR were estimated at 1,163 in 2022. Significant declines occurred in two of the four populations known in the 1980s. The Ka‘ū /Kapāpala population decreased from approximately 530 individuals to 44, and a Mauna Kea population dropped from approximately 50 birds to less than 10; in 2000 only three birds remained on Maunakea and this and the Kona populations are now extinct. The Ka‘ū /Kapāpala population has since stabilized or increased at upper elevations.. ‘Akiapōlā‘au however have responded extremely well to koa reforestation, and populations at Hakalau forest have increased by more than 900% in the koa planting areas.

LOCATION AND CONDITION OF KEY HABITAT: Occurs in mesic and wet montane forests dominated by koa and ‘ōhi‘a. The now extinct population of Maunakea ea occurred in subalpine dry forest dominated by māmane and naio. A studies have documented ‘akiapōlā‘au occurring entirely in areas reforested with koa (i.e., second-growth, young forests). Habitat quality varies across the species’ occupied range, but reforestation and fencing have drastically improved some key forestes in the last 50 years. Most remaining populations occur on lands managed by the State of Hawai‘i and the U.S. Fish and Wildlife Service.

THREATS: ‘Akiapōlā‘au are susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For ‘akiapōlā‘au populations, the following are of particular concern:

- Low reproductive potential. ‘Akiapōlā‘au have low annual fledgling production. This life history characteristic may be related to their very specialized foraging strategy. Regardless, the species is very susceptible to factors that reduce population size.
- Disease. Unlike several other honeycreepers found on the island of Hawai‘i (e.g., Hawai‘i ‘amakihi [*H. virens*]), the ‘akiapōlā‘au is absent from most areas below 1,350 meters (4,500 feet). This suggests that the species is particularly susceptible to mosquito-borne avian disease.
- Predation. Predation by rats (*Rattus* spp.), cats (*Felis silvestris*), small Indian mongoose (*Herpestes auropunctatus*), and raptors (Pueo *Asio flammeus sandwichensis*, Barn owl *Tyto alba*, and ‘Io *Buteo solitarius*) may limit ‘akiapōlā‘au populations. Surveys have determined that rat density in the Hakalau Forest National Wildlife Refuge, which supports a significant portion of the ‘akiapōlā‘au population, is high. In addition, the loud, persistent begging of juveniles may make them especially vulnerable to predators.
- Habitat degradation. Habitat loss and degradation from development, logging, and grazing, and feral ungulates has greatly fragmented the species’ habitat.

- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios and stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems.

CONSERVATION ACTIONS: To date, conservation actions specific to ‘akiapōlā‘au have been restricted to annual population surveys of the Hakalau, ‘Ōla‘a/Kīlauea, Kona, and Mauna Kea populations. However, ‘akiapōlā‘au likely have benefited from actions to conserve other endangered forest birds in the Kapāpala Forest Reserve, Hakalau Forest National Wildlife Refuge, Pu‘u Lā‘au, Hawai‘i Volcanoes National Park, and the ‘Ōla‘a/Kīlauea Watershed Partnership. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the ‘akiapōlā‘au may include the following:

- Add Hāmākua, the upper Waiākea kīpuka, Ka‘ū /Kapāpala and south Kona to annual surveys
- Expand fencing efforts of key habitats above 4500ft in the above mentioned reserves
- Continue koa forest restoration and fencing in the Hakalau Forest National Wildlife Refuge.
- Continue restoration of māmane forests on Maunakea.
- Conduct public outreach and education and expand community conservation efforts
- Continue protection and management of wildlife sanctuaries and refuges
- Expand and diversify predator control throughout the species range
- Translocate birds back to Maunakea and other suitable habitats in their former range
- Collect genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning

MONITORING:

- Continue forest bird surveys and habitat monitoring
- Expand the use of Automatic Recording Units (ARU) to remotely monitor populations
- Test survey methods for ‘akiapōlā‘au, and continue regular population surveys with improved methods.
- Monitor small mammal populations to assess effectiveness of control efforts, especially in dry forest sites.
- Conduct range wide population estimates and determine population trends.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to ‘akiapōlā‘au include the following:

- Conduct life history studies to quantify population structure, dispersal patterns, survivorship, nesting phenology, and success
- Document habitat selection, preference, and foraging ecology, particularly in young forests

- Document the response of ‘akiapōlā‘au to control of mammalian predators
- Develop captive propagation techniques.
- Determine the feasibility of ‘akiapōlā‘au re-introductions to suitable locations (e.g., Pu‘u Wa‘awa‘a, Hawai‘i Volcanoes National Park)
- Record vocalizations of distinct populations for archival analysis

References:

- Gorreson, PM, Camp RJ, Reynolds MH, Woodworth BL, Pratt TK. 2009. Status and Trends of Native Hawaiian Songbirds. *In* Conservation Biology of Hawaiian Forest Birds (Pratt TK, Atkinson CT, Banko PC, Jacobi JD, Woodworth BL, editors). Yale University Press.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Kendall SJ, Rounds RA, Camp RJ, and Genz AS. 2022. Forest Bird Populations at the Big Island National Wildlife Refuge Complex, Hawaii. Hawai‘i Cooperative Studies Unit Technical Report HCSU-102. University of Hawai‘i at Hilo, Hawaii, USA. 141 pages.
- Pejchar, L. 2005. Home range size varies with habitat type in a Hawaiian honeycreeper: implications for native *Acacia koa* forestry. *Ecological Applications* 15:1053-1061.
- Pratt TK, Fancy SG, Ralph CJ. 2001 ‘Akiapola‘au (*Hemignathus munroi*) and nukupu‘u (*Hemignathus lucidus*). *In* The Birds of North America, No. 600 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Strommer L. University of Hawai‘i graduate student. Unpublished data.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

‘Akikiki

Oreomystis bairdi

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Critically Endangered


1 Recovery Plan for Hawaiian Forest Birds—USFWS 2006

Kauai Islandwide Recovery Plan – USFWS 2021

SPECIES INFORMATION: The ‘akikiki, or Kaua‘i creeper, is a small, drab Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Kaua‘i. Both males and females are predominantly dark gray to olive above, whitish below. ‘Akikiki have pinkish legs and feet, and their short, slightly decurved bill also is pink. Usually found in pairs, family groups, or small flocks (8 – 12 individuals); during the non-breeding season ‘akikiki join mixed species foraging flocks. ‘Akikiki gleans and probes the bark and lichens and moss on trunks, branches, and twigs of live and dead ‘ōhi‘a (*Metrosideros polymorpha*) and koa (*Acacia koa*) trees for insects and spiders. They usually nest high (9 meters or higher) in the terminal branches of ‘ōhi‘a, but one female has been observed nesting in ‘ōlapa on two occasions. Nest construction begins in early March, and continues into June; males occasionally help with nest building. Most nests contain two eggs, some contain only one egg. Both males and females feed chicks. Despite a long period of parental dependency, many cases of double brooding have been observed, with the male provisioning both the incubating female and the older chicks. Causes of nest failure include predation by rats, poor female attendance, infertility, weather, and hatch failure.

DISTRIBUTION: As of 2024, apparently extirpated throughout most of its recent range in the Alaka‘i Wilderness Preserve, including Halepa‘akai, with a few individuals still occurring in and near the headwaters of Mohihi Stream and side drainages feeding Koai‘e Stream where it turns west. Occupancy rates in 2012 increased from west to east along the plateau, from 0.02 ± 0.07 near Koke‘e State Park to 0.55 ± 0.21 in the southeast part of the range, and were positively correlated with canopy height. The suitable habitat range was recently estimated at $\sim 38\text{km}^2$ and suitable nesting habitat $\sim 18\text{ km}^2$ (Fricker et al. 2021), but newer estimates in 2004 suggest that it is only 94 ha (233 ac; KFBRP, unpubl. data). Historically occupied high- and low-elevation forests, although by the 1960s was most common above 1,140 meters (3,750 feet), and is now restricted to those higher elevations. Subfossil remains suggest a prehistoric island-wide distribution.

ABUNDANCE: The Kaua‘i Forest Bird Survey (KFBS 2000), estimated the population at 2,448 $\pm$ 1,200 (SE) birds. Density estimates were 26 birds per square kilometer; 15 percent lower than in 1981. In 2012, the KFBS estimated density at 8.8 birds per square kilometer, and the population at only 468 (95% confidence interval: 231-916) birds. In 2024, despite extensive surveys through its former range, only four-five individuals were detected, although it is possible that a few more individuals occur on private land or escape detection due to lack of intensive survey effort.



LOCATION AND CONDITION OF KEY HABITAT: Currently restricted to elevations > 1200 m (mean 1304.5 $\pm$ 38.5 m; Fricker et al. 2021) and are found only in the wet tropical rainforest of the eastern Plateau. Occupancy is positively correlated with elevation and large trees (i.e., canopy height) and canopy density. They also exhibit a tendency to nest along lower slopes (e.g., along streams in areas of higher canopy height, and more complex understory (Fricker et al. 2021). The montane forests of Kaua‘i are dominated by ‘ōhi‘a with a subcanopy comprising ‘ōlapa or lapalapa (*Cheiodendron* spp.) and ‘ōhi‘a hā (*Syzygium sandwicensis*). Common understory species include ‘ōhelo (*Vaccinium calycinum*), kanawao (*Broussaisia arguta*), ‘ōhā wai (*Clermontia* spp.), kāwa‘u (*Ilex anomala*), kōlea (*Myrsine lessertiana*), na‘ena‘e (*Dubautia* spp.), and pūkiawe (*Styphelia tamieameiae*). Occupancy is very low in areas invaded by non-native plants.

Hawai‘i's State Wildlife Action Plan
October 1, 2025

THREATS:

- Disease. Mosquitoes likely are ubiquitous on Kaua‘i, and avian malaria and avian pox are likely the most important factors limiting the species’ distribution. Five of 42 ‘akikiki have tested positive for malarial antibodies between 2011 and 2023; some were later resighted, indicating that a few individuals have resistance or tolerance. However, three of the individuals that tested positive were caught in 2022, and one was resighted up to nine months later, but then disappeared. The recent dramatic population declines indicate that avian malaria is a primary threat to this species. Pox lesions have been observed on several birds in the wild, and pox was implicated in the mortality of several individuals transferred to human care in 2023 (KFBRP and SDZWA, unpubl. data).
- Habitat degradation. Pigs (*Sus scrofa*) and goats (*Capra hircus*) have contributed to the spread of non-native plants, but effects to ‘akikiki are unknown. Severe hurricanes in 1982 and 1992 heavily damaged native forests, possibly resulting in short-term reductions in arthropod food resources and long-term damage to forest structure preferred by ‘akikiki, given their affinity for large trees, which are now rare on the landscape
- Natural disasters. Hurricanes in 1982 and 1992 likely caused the death of an unknown number of individuals.
- Competition. Although little evidence exists, it has been suggested that competition with introduced Warbling white-eyes (*Zosterops japonicus*) and Japanese bush warblers (*Horornis diphone*) may negatively affect ‘akikiki. Non-native insects, especially yellow jackets and ants, may compete with or prey on the native arthropods on which ‘akikiki feed. The role of non-native insects in native Hawaiian forests is unclear.
- Predation. Predation by rats (*Rattus* spp.), on eggs has been documented in several instances; predation on chicks and females is suspected in a few cases. Cats (*Felis catus*), Hawaiian short-eared owls (*Asio flammeus sandwichensis*), and barn owls (*Tyto alba*) occur throughout the forests of Kaua‘i, and may prey on young and adults.
- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios and stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems. Some of the observed infertility and failure to hatch may be due to these small population issues.

CONSERVATION ACTIONS: ‘Akikiki likely have benefited from actions to conserve other endangered forest birds including establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, monitoring habitat conditions, studies of disease and disease vectors, control of feral ungulates through public hunting and fencing, control of feral ungulates through public hunting and fencing, suppression of rats on two A24 trapping grids over approximately 150 ha (470 ac), and public education efforts. Between 2015 and 2018, eggs were collected to initiate captive flock of about 45 individuals. Six adults and subadults were added to the flock between 2021 and 2023, but they did not survive. Overall, mortality and reproduction are approximately even in the captive population, thus the current flock numbers still about 43 birds. In addition to these efforts, future management specific to the recovery of ‘akikiki may include:

- Continue to collect wild ‘akikiki adults, subadults, chicks, and eggs; Aggressively control ungulates to improve habitat quality and facilitate recovery of degraded, but potential, habitat. It could also reduce breeding habitat for mosquitoes. Control of non-native plants should be part of forest restoration efforts.
- Reduce or manage mosquito breeding habitat on the Alaka‘i Plateau
- Suppress mosquito populations through releases of incompatible or sterile mosquitoes
- Investigate options to eradicate mosquitoes, including use of genetic manipulations.
- Suppress/control rats, feral cats, and barn owls from the Alaka‘i Wilderness Preserve, Na Pali-Kona Forest Reserve, and Koke‘e State Park.
- Tranlocations to areas with low disease prevalence on other islands
- Prevent the introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other potential predators.
- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.
- Prevent the spread of Rapid Ohia Death on the Alaka‘i Plateau
- Collect genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning

MONITORING: Conduct species-specific monitoring for ‘akikiki (e.g., occupancy surveys with play back, territory monitoring) that take into account the inability of the HFBS to track populations of rare species.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the ‘akikiki include the following:

- Investigate techniques for improving survival and reproduction of the captive flock
- Conduct PVAs for captive flock, including impact of reintroduction back into the wild once malaria is controlled
- Explore the use of surrogate species such as ‘alawī to restore wild behaviors and or to establish birds in the wild on Hawai‘i Island
- Investigate and evaluate different strategies for reintroducing the population to the wild
- Continue to assess the susceptibility of this species to avian malaria and avian pox.
- Determine sources of mosquitoes and investigate methods of mosquito control.
- Determine the effects of recently established non-native insects on native arthropods, especially those that are part of the species’ diet.
- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps).
- Assess the impact of introduced rodents on food sources important to ‘akikiki.
- Investigate the feasibility of food supplementation.

References:

- Atkinson, CT, RB Utzurrum, DA LaPointe, RJ Camp, LH Crampton, JT Foster, and TW Giambelluca. 2014. Changing Climate and the Altitudinal Range of Avian Malaria in the Hawaiian Islands - an Ongoing Conservation Crisis on the Island of Kaua‘i. *Global Change Biology* 20, 2426–2436, doi: 10.1111/gcb.12535.

- Behnke, L. 2014. Habitat use and conservation implications for Akikiki (*Oreomystis bairdi*) and Akekee (*Loxops caeruleirostris*), two endangered Hawaiian honeycreepers. M.S. Thesis. Colorado State University, Fort Collins, CO.
- Foster JT, Scott JM, Sykes PW. 2000. 'Akikiki (*Oreomystis bairdi*). In The Birds of North America, No. 552 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka'i swamp, Kaua'i. *Conservation Biology* 18:716-725.
- Fricker GA, Crampton LH, Gallerani EM, Hite JM, Inman R, Gillespie TW. 2021. Application of lidar for critical endangered bird species conservation on the island of Kauai, Hawaii. *Ecosphere* 12(6).
- Hammond, RL, Crampton, LH, and Foster, JT. 2015. Breeding biology of two endangered forest birds on the island of Kaua'i. *Condor* 117: 31-40.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.
- U.S. Fish and Wildlife Service. 2021. Kauai Islandwide Recovery Plan. Portland, (OR): U.S. Fish and Wildlife Service. <https://www.fws.gov/node/264716>



Photo: Robby Kohley

Forest Birds 'Ākohekohe

Palmeria dolei

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled


IUCN Red List Ranking—Critically Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The 'ākohekohe, or crested honeycreeper (Family: Fringillidae), is the largest extant honeycreeper on Maui Nui (Lāna'i, Moloka'i, Maui, and Kaho'olawe). Although primarily black, the plumage of the 'ākohekohe is striking. Depending on their location, feathers are tipped with orange-yellow, gray, silver, or white. Orange feathers surround the eyes and extend over the nape, orange or yellow-white feathers cover the thighs, the epaulettes are white with orange tips, and there is a distinctive plume of white feathers that curl forward over the bill. They do not sing, but produce a random series of buzzes, croaks, and whistles. They are primarily nectarivorous, feeding mainly on 'ōhi'a (*Metrosideros polymorpha*), but also from the flowers of other trees and shrubs, such as 'ākala (*Rubus hawaiensis*) and kōlea (*Myrsine* species). Like 'apapane (*Himatione sanguinea*) and 'i'iwi (*Vestiaria coccinea*), 'ākohekohe are strong fliers and will move from low to high elevations in search of blooming 'ōhi'a. Arthropods, mainly gleaned from 'ōhi'a, are also part of the species' diet. They spend up to 70 percent of the day foraging. They aggressively defend feeding and nesting territories year-round. Females build open-cup nests primarily in 'ōhi'a, incubate the clutch of one or two eggs, and brood nestlings; male feeds female on nest. Fledglings can forage independently 10 to 14 days after leaving the nest. Pairs successfully fledge two to three broods per season.

DISTRIBUTION: Restricted to a 24 square kilometer (9 square mile) area on the northeastern slope of Haleakalā at 1,600 to 2,300 meters (5,200 – 7,000 feet). Subfossil evidence indicates they once occurred in Maui's lowland dry forests, and historically they also once occurred on eastern Moloka'i. They currently occupy 2 percent of their historical range.

ABUNDANCE: The Hawai'i Forest Bird Survey from 2022 estimated the population at 1768 ± 315 (95% CI 1193–2411) individuals, a 78% decline from 2001.



LOCATION AND CONDITION OF KEY HABITAT: Wet and mesic montane forests dominated by ‘ōhi‘a and ‘ōlapa (*Cheirodendron trigynum*); koa (*Acacia koa*) and kāwa‘u (*Ilex anomala*) occur at lower densities. Nearly all birds occur in forest between 1,700 and 2,100 meters (5,700 – 6,900 feet) elevation in rugged, steep terrain with a dense understory. The entire known range of the species occurs within State (e.g., Forest Reserve and Natural Area Reserve), Federally (e.g., National Park), or Privately (e.g. The Nature Conservancy) managed lands.

THREATS: ‘Ākohekohe are susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For ‘ākohekohe, the following are of particular concern:

- Disease. Similar to ‘apapane and ‘i‘iwi, movements between low- and high-elevation foraging sites may increase these birds’ exposure to mosquito-borne diseases.
- Habitat degradation. Feral pig (*Sus scrofa*), deer (*Axis axis*), and goats (*Capra hircus*) damage understory vegetation may reduce the availability of nectar-producing plants important to ‘ākohekohe, especially those flowering when ‘ōhi‘a nectar is less available.
- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios and

stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems.

CONSERVATION ACTIONS: Captive propagation of ‘ākohekohe was attempted decades ago but was not successful. ‘Ākohekohe likely benefited from actions to conserve other endangered forest bird species on the northeastern slope of Haleakalā, including fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future actions specific to ‘ākohekohe may include the following:

- Establish a second population to reduce the chances that a catastrophe could result in the species’ extinction. Potential re-introduction sites (e.g., West Maui and Moloka‘i) are limited because of the presence of mosquitoes; therefore, it has been prioritized to research potential locations for a conservation translocation on Hawai‘i Island (e.g. Ka‘ū FR, Hilo FR, Hakalau NWR, Pu‘u Maka‘ala NAR, or Pu‘u Wa‘awa‘a FBS) which has more high elevation native protected forest than Maui.
- Capture and hold a small number of individuals to develop and adaptively manage captive holding conditions to reduce stress and promote breeding/cohabitation. These efforts would inform if and/or how captive holding can be used as part of the conservation strategy for this species.
- Develop and deploy mosquito control techniques as soon as possible to reduce the mosquito vector of avian disease.
- Continue fence management and feral animal control to improve understory conditions in occupied habitat and potentially facilitate expansion of ‘ākohekohe populations if and when mosquito control occurs.
- Conduct public outreach and education. Incorporate Hawaiian knowledge and practices in conservation strategies
- Continue protection and management of wildlife sanctuaries and refuges. Collect genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning
- Develop and deploy new tools and evaluate above approaches.

MONITORING: Continue forest bird surveys, habitat monitoring, and use of automated recording units (ARU).

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to ‘ākohekohe include the following:

- Determine if disease-resistant individuals exist, and if so, if resistance is passed to offspring. Disease-resistant individuals could be used to establish new populations.
- Determine the role of ‘ākohekohe in transmitting disease between high- and low-elevation habitats.
- Determine if a conservation translocation is feasible to Hawai‘i Island and develop translocation and field holding protocols.

References:

- Berlin KE, VanGelder EM. 1999. 'Akohekohe (*Palmeria dolei*). In The Birds of North America, No. 400 (Poole A, Gill F, editors). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Judge, SW, Camp RJ, Warren CW, Berthold LK, Mounce HL, Hart PJ, and Monello RJ. 2021. Population Estimates and Trends of Three Maui Island-endemic Hawaiian Honeycreepers. Journal of Field Ornithology. <https://onlinelibrary.wiley.com/doi/full/10.1111/jfo.12364>
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.
- Wang AX, Paxton EH, Mounce HL, Hart PJ. 2020. Divergent movement patterns of adult and juvenile 'ākohekohe, an endangered Hawaiian Honeycreeper. Journal of Field Ornithology.



Photo: DOFAW

Forest Birds

‘Alalā

Corvus hawaiiensis

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic


NatureServe Heritage Rank: GXC-Presumed Extinct/

Captive Population

IUCN Red List Ranking—Extinct in the Wild

Revised Recovery Plan for the ‘Alalā —USFWS 2009

SPECIES INFORMATION: Historically at least five crow species (Family: Corvidae) occurred in Hawai‘i, only the ‘alalā, or Hawaiian crow survives. Like other crows, ‘alalā are raucous, gregarious and vocal; young, captive-raised birds often engage in tug-of-war with sticks. Like many corvids, ‘alalā are long-lived with a life span of 20 or more years. The diet primarily consists of native and introduced fruits, invertebrates, and eggs and nestlings of other forest birds, as well as nectar, flowers and carrion. Seasonal movements in response to weather and availability of food plants (e.g., ‘ie‘ie [*Freycinetia arborea*]) have been noted. Although individuals form long-term pair bonds, extra-pair copulations have been observed. Nests are predominantly constructed in ‘ōhi‘a (*Metrosideros polymorpha*) trees. Both sexes participate in nest construction, although only females incubate eggs and brood young. Clutch size ranges from two to five, although usually only one or two nestlings fledge. Fledglings typically cannot fly and often remain near the ground for long periods, likely increasing their susceptibility to disease (i.e., toxoplasmosis) and predation. Juveniles depend on their parents for at least eight months and remain with their family group until the following breeding season. Large flock’s characteristic of American crows (*C. brachyrhynchos*) have not been reported, but there are historical reports of small local flocks after the breeding season.



DISTRIBUTION: No individuals are known to exist in the wild. Historically occurred in high- and low-elevation forests of the western and southeastern regions of the island of Hawai‘i.

ABUNDANCE: World population of 108 individuals in 2024, housed entirely in the Keauhou and Maui Bird Conservation Centers.

LOCATION AND CONDITION OF KEY HABITAT: Historically, ‘alalā occupied dry and seasonally wet ‘ōhi‘a and ‘ōhi‘a/koa (*Acacia koa*) forests between 300 and 2,500 meters (1,000 – 8,200 feet) elevation. Because the last wild individuals were confined to a small subset of the species’ former range, specific knowledge of key habitat requirements are unknown. Currently, all potential habitat is degraded. The presence of non-native mammalian predators and birds, which can act as disease reservoirs, further reduces habitat quality. Core areas of the species’ former range are now managed by the State of Hawai‘i and the U.S. Fish and Wildlife Service.

THREATS: ‘Alalā are susceptible to the same factors that threaten other native Hawaiian forest birds, including: loss and degradation of habitat, predation by introduced mammals, and disease. For ‘alalā populations, the following are of particular concern:

- Predation. The small Indian mongoose, rats, and feral cats prey on ‘alalā. The ‘io (*Buteo solitarius*) and presumably pueo (*Asio flammeus sandwichensis*) also prey on juvenile and adults. ‘Io have been documented killing captive-raised birds released into the wild. Fledglings are unable to fly and this likely contributes to high rates of predation.
- Shooting. Many ‘alalā were killed around farms between 1890 and 1930. Despite legal protection in 1931, shooting of individuals occurred into the 1980s.
- Disease. Population declines were noted between 1890 and 1910, a period when other native bird populations declined, presumably because of the introduction of rats and increased rates of deforestation. In general ‘alalā have proven resistant to avian malaria and pox virus, but they are susceptible to toxoplasmosis spread by the parasite’s obligate host, domestic cats (*Felis catus*)
- Habitat degradation. Habitat conversion by human activity as well as by grazing ungulates and seed predation by rats has severely degraded former ‘alalā habitat. These changes may have limited food or nesting resources and may have increased the vulnerability of ‘alalā to predation by ‘io. Currently, little suitable habitat exists for the species.
- Population size. Small populations are plagued by a variety of potentially irreparable problems which fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios.
- Captive-breeding. There is some evidence that captive-reared birds lack important foraging and predator-avoidance behaviors.

CONSERVATION ACTIONS: The ‘alalā has been legally protected by the State of Hawai‘i since 1931 and was listed as federally endangered in 1967. A captive propagation program was established in 1973; crows are now housed at the Keauhou Bird Conservation Center and the Maui Bird Conservation Center. The ‘Alalā Recovery Team was formed to facilitate the species recovery, and a related second group, the ‘Alalā Partnership, was formed to facilitate program

implementation on private lands. Between 1993 and 1998, 27 captive-raised juvenile ‘āla‘ā were released at McCandless Ranch. Of these, 21 died in the wild and six were recaptured and returned to the captive flock. Predator control was ongoing during the release program. Intensive field studies of the wild population and released juveniles were conducted between 1992 and 2002. In 1999, the Kona Forest Unit of Hakalau Forest National Wildlife Refuge was acquired, with the goal of restoring habitat in the core of the species’ historic range. Recent release efforts were made from 2016 to 2019 in the Pu‘u Maka‘ala Natural Area Reserve on Hawai‘i Island. This is a fenced area, free of ungulates, creating a dense understory containing many plants suitable for ‘āla‘ā to eat. This set of releases showed improvements and were considered successful due to birds surviving for multiple years and one pair observed nesting. ‘Āla‘ā mortalities increased in 2020 due to predation from ‘io or Hawaiian hawk (*Buteo solitarius*) and other factors, ultimately causing the remaining five individuals to be brought back into captivity. Five ‘āla‘ā were released on Maui within the Kipahulu Forest Reserve in 2024. The birds have fared well without the presence of ‘io and any new findings from this release can apply to future releases. Additionally, efforts have been made to research ‘io behavior to help inform site selection for Hawai‘i island. Future releases will also occur elsewhere on Maui Nui or at a site on Hawai‘i Island.

Restoration of future re-introduction sites is ongoing and re-introductions are expected to occur by 2027 within the Ka‘ū Forest Reserve. In addition to the above efforts, ‘āla‘ā likely will benefit from management activities to conserve other endangered forest birds on the island of Hawai‘i including fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the ‘āla‘ā should include the following:

- Continue restoration of future reintroduction areas throughout historic range.
- Maintain and increase the captive flock without further loss of genetic diversity.
- Continue protection and management of wildlife sanctuaries and refuges.
- Evaluating habitat on additional islands for releases.
- Expand and improve collecting genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning.

MONITORING: The captive flock is monitored. Wild populations are intensively monitored using a suite of remote and in the field techniques including game cameras, transmitters, and direct observation.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the ‘āla‘ā include the following:

- Review all data from studies on captive and wild populations.
- Determine methods to increase the reproductive output of captive individuals.
- Conduct field studies to determine if understory restoration will reduce the ability of ‘io to prey on ‘āla‘ā.
- Establish a set of habitat criteria to inform releases of birds at any particular site.

- Develop methods to habituate captive-raised individuals to respond appropriately to mammalian and avian predators, and sources of toxoplasmosis.
- Determine potential reintroduction sites on other islands.
- Develop methods to assess the impact of ‘alalā on other rare taxa such as snails and forest birds and mitigation strategies.

References:

- Banko PC, Ball DL, Banko WE. 2002. Hawaiian crow (*Corvus hawaiiensis*). In The Birds of North America, No. 648 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Berger AJ. 1981. Hawaiian birdlife. Honolulu: University of Hawai‘i Press. 260 pp.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- U.S. Fish and Wildlife Service. 2009. Revised recovery plan for the ‘Alala (*Corvus hawaiiensis*). Portland, (OR): U.S. Fish and Wildlife Service. Xiv+105 pp.



Photo: Jim Denny

Forest Birds

‘Anianiau

Magumma parva


SPECIES STATUS:

State Recognized as Endemic

NatureServe Heritage Rank G2—Imperiled

IUCN Red List Ranking—Endangered

SPECIES INFORMATION: The ‘anianiau is the smallest Hawaiian honeycreeper (Family: Fringillidae). Endemic to Kaua‘i, the ‘anianiau also is one of the most common native birds of the island’s high-elevation forests. Adult males are brilliant yellow; females also are yellow, but may be duller, although some bright females have been observed. Constantly on the move, ‘anianiau feed on nectar from ‘ōhi‘a (*Metrosideros polymorpha*), ‘ōhelo (*Vaccinium* spp.), ‘alani (*Melicope* spp.), and other native and introduced plants. They also glean arthropods from the outer canopy and smaller twigs and branches of ‘ōhi‘a and koa (*Acacia koa*) trees as well as from the foliage of shrubs, vines, and the fronds of tree ferns (*Cibotium* spp.). Nectar, spiders, and Lepidoptera larvae compose the bulk of the species’ diet. ‘Anianiau are occasionally seen in small flocks, especially at favored nectar sources. Males sing a sweet, high-pitched trill, and predominantly defend breeding territories that may be as small as 9 meters (29.3 feet) in diameter. Both sexes build the open-cup nest, females incubate eggs and brood young, and males provision females, generally away from the nest. No information on post-fledgling behavior or dependency. ‘Anianiau nest success rate in 2012 and 2013 was 0.56 (± 0.09 , Hammond et al. 2015).



DISTRIBUTION: May occur as low 774 m elevation in native forests of the Kōke‘e and Alaka‘i regions, but are becoming scarce in the former regions. However, the lowest elevation detection in 2024 was at 1131 m (KFBRP, unpubl. data). Original range likely included all forested regions of Kaua‘i. Notable range contractions of more than 60% occurred between 2018 and 2023 with the ‘anianiau now occupying a range of approximately 3648 ha (9014 ac; Kaua‘i

Forest Bird Recovery Project, unpubl. data).

ABUNDANCE: In the early 1970s the island-wide ‘anianiau population was estimated at 24,000 $\pm$ 3,000 (SE) individuals. The Kaua‘i Forest Bird Survey (2000) estimated the population within the Alaka‘i Wilderness Preserve and Kōke‘e region at close to 35,000 individuals, with a significant population increase between 1981 and 2000. Although the 2008 surveys suggested that the population appeared stable in its current range, the population was estimated around only 10,000 individuals in 2018 (Paxton et al. 2020). The 2023 surveys suggested further declines to 3,827 individuals (DOFAW, unpublished data). .

LOCATION AND CONDITION OF KEY HABITAT: Mesic and wet forests above 1,000 meters (2,000 feet), with the highest densities above 1,200 meters (3,600 feet). These forests are dominated by ‘ōhi‘a, koa, ‘ōlapa (*Cheirodendron trigynum*), and lapalapa (*C. platyphyllum*). At lower elevations, where the species historically occurred, native habitats are severely degraded. Although public hunting reduces the number of feral ungulates in the most accessible parts of the species’ range, hunting is not an effective method to prevent habitat degradation across its entire range. Occupied habitats are largely managed by the State of Hawai‘i, although a few individuals may occur on private lands around the Plateau.

THREATS:

- Disease. Thirteen of 129 ‘anianiau sampled between 2011 and 2023 tested positive for malarial parasites (*Plasmodium relictum*) in areas where parasites were common in other species. Since malaria is established on the Alaka‘i Plateau, the most likely explanation for this low prevalence is high mortality after infection with malaria. Avian malaria is likely the primary threat to the species. Research comparing mortality between Kaua‘i honeycreepers (‘anianiau and ‘akikiki) and non-honeycreepers (puaiohi and Kaua‘i ‘elepaio), found that the honeycreepers were much more susceptible to avian malaria (Whitaker 2022).
- Habitat degradation. The species is tolerant to habitat alteration, but it is most common in undisturbed native forest. Introduction of non-native plants is the most important threat as ‘anianiau density is negatively related to the presence of non-native shrubs.
- Competition. Competition with introduced Japanese white-eyes (*Zosterops japonicus*) may negatively affect ‘anianiau. Non-native insects, especially yellow-jackets (*Vespula pensylvanica*) and ants (*Linepithema humile*), may compete with or prey on the native arthropods on which ‘anianiau feed. The role of non-native insects in native forest ecosystems is unclear.
- Mammalian predators. Rats (*Rattus* spp.), cats (*Felis catus*), Hawaiian short-eared owls (*Asio flammeus sandwichensis*), and barn owls (*Tyto alba*) occur throughout the forests of Kaua‘i. Hammond et al. 2015 found evidence of ‘anianiau nest predation in the form of trail camera pictures, egg-shell fragments in the nest and even chick remains.

CONSERVATION ACTIONS: ‘Anianiau likely benefit from actions to conserve other endangered forest birds including establishment of the Alaka‘i Wilderness Preserve, regular

surveys of forest bird populations, habitat monitoring, studies of disease and disease vectors, control of feral ungulates through public hunting and fencing, suppression of rats on two A24 trapping grids over approximately 150 ha (470 ac), and public education featuring Kauai's endangered forest birds. Conservation efforts should include continued protection and management of wildlife sanctuaries and refuges. Since the population is declining rapidly, collection of 'anianiau adults and eggs has been implemented to establish an ex-situ insurance population, with 13 birds currently in human care. In addition to these efforts, future actions specific to the 'anianiau may include the following:

- Aggressively control ungulates to improve habitat quality, facilitate the recovery of degraded habitat, and potentially reduce breeding habitat for mosquitoes. Control of non-native plants should be part of forest restoration efforts.
- Eradicate or manage mosquito breeding habitat on the Alaka'i Plateau
- Suppress mosquito populations through releases of incompatible or sterile mosquitoes
- Investigate options to eradicate mosquitoes, including use of genetic manipulation
- Translocate individuals to habitat with low levels of disease prevalence on Hawai'i Island
- Suppress/control rats, feral cats, and barn owls from the Alaka'i Wilderness Preserve, Na Pali-Kona Forest Reserve, and Kōke'e State Park
- Prevent the introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other predators.
- Prevent the spread of Rapid Ohia Death on the Alaka'i Plateau
 - Conduct public outreach and education.
 - Continue protection and management of wildlife sanctuaries and refuges
 - Expand and improve collecting genetic samples from as many individuals as possible for bio-banking, genetic rescue, and cloning

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the 'anianiau include:

- Conduct life history studies to quantify the population structure, dispersal patterns, survivorship, nesting phenology and success of this poorly known species.
- Investigate techniques for improving survival and reproduction of the captive flock
- Conduct PVAs for both wild and captive flocks
- Investigate the feasibility of 'anianiau translocation to Hawai'i Island
- Determine species specific avian malaria mortality rate and if potential resistance is passed to offspring. Disease-resistant individuals could be used as founders for new populations.
- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps)
- Assess the impact of introduced rodents on food resources (e.g., floral, invertebrates) important to 'anianiau

- Determine if competition with Warbling white-eyes or Japanese Bush-warblers occurs, and if so, its effect on ‘anianiau populations.
- Determine the effects of recently established non-native insects on native arthropods, especially on those arthropods that are part of the species’ diet.
- Determine the status of populations outside of the greater Alaka‘i Swamp region.

References:

- Camp, RJ and PM Gorresen. 2011. Design of forest bird monitoring for strategic habitat conservation on Kaua‘i Island, Hawai‘i. Hawai‘i Cooperative Studies Unit Technical Report HCSU-022. University of Hawai‘i at Hilo.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- Hammond RL, Crampton LH, Foster JT. 2015. Nesting success of native and introduced forest birds on the island of Kauai. *Journal of Avian Biology* 47: 252-262.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Lepson JK. 1997. ‘Anianiau (*Hemignathus parvus*). In *The Birds of North America*, No. 312 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Paxton ER, Camp MG, Crampton LH, Leonard DL, VanderWerf EA. 2016. Collapsing avian community on a Hawaiian island. *Science Advances* 2:e1600029.
- Paxton ER, Brinck KW, Crampton LH, Hite J, Constantini M. 2018 Kaua‘i forest bird population estimates and trends. Technical Report Hawai‘i Cooperative Studies Unit. University of Hawai‘i at Hilo.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. *Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation*. Lawrence, (KS): Cooper Ornithological Society.
- Whitaker K. 2022. Avian malaria on Kaua‘i: prevalence and survival as a function of sex, age, and species. Master’s thesis, Northern Arizona University.



Photo: Eric Nishibayashi

Forest Birds

‘Apapane

Himatione sanguinea

SPECIES STATUS:


State Recognized as Endemic
NatureServe Heritage Rank G3—Vulnerable
IUCN Red List Ranking—Least Concern

SPECIES INFORMATION: The ‘apapane is a small, crimson, primarily nectarivorous Hawaiian honeycreeper (Family: Fringillidae) and is an important ‘ōhi‘a (*Metrosideros polymorpha*) pollinator. It is the most abundant and widely distributed Hawaiian honeycreeper, and is often seen flying above the canopy in search of patches of flowering ‘ōhi‘a. Wide-ranging movements may facilitate disease transmission among native forest birds. ‘Apapane often forage in conspecific flocks, likely to overwhelm ‘i‘iwi (*Vestiaria coccinea*) and ‘ākohekohe (*Palmeria dolei*), which often defend flower-rich trees. Outside the breeding season, ‘apapane also join mixed-species flocks. They feed on insects, which they glean from outer foliage and twigs in the upper- and mid-canopy. Sexual chasing and courtship feeding often precede nest building, a task shared by both male and female. Pairs defend small territories around nests. Females incubate three eggs and brood young; males feed females away from the nest. Both parents feed nestlings, and fledglings may remain with their parents for up to four months.

DISTRIBUTION: Occurs in native forests on the islands of Hawai‘i, Maui, and Kaua‘i. On O‘ahu, occurs in the Ko‘olau Range from 300 meters (975 feet) to summit at 946 meters (3,075 feet), and are less common in the Wai‘anae Range above 600 meters (1,950 feet). Rare on Moloka‘i and Lāna‘i. Historically were common at low elevations on all islands with appropriate habitat.

ABUNDANCE: Based on Hawaiian Forest Bird Surveys (1976-1981):

1,080,000 $\pm$ 25,000 (95% confidence interval) birds on island of Hawai‘i, 110,000 $\pm$ 9,000 on Maui (86% on Haleakalā), 39,000 $\pm$ 5,000 on Moloka‘i, 540 $\pm$ 213 on Lāna‘i, and 30,000 $\pm$ 1,500 on Kaua‘i (O‘ahu was not included in surveys). On Kaua‘i, populations declined after the 1992 hurricane but have significantly increased since, estimated at 64,972 $\pm$ 2,014 (SE) birds in



2000 then at 95,511 in 2023 (95% CI: 85,433-107,052; DOFAW, unpublished data). Rare on Moloka‘i and Lāna‘i.

Maui nui updated numbers-

East Maui- 228,480 ± 19,855 individuals

West Maui- we don’t have an updated estimate, but they are the most commonly detected species on counts. 4.53 detections per station.

Lanai- they continue to be detected on Lanai in small numbers, based on surveys from 2023.

Molokai- 81,896 ± 10, 836

LOCATION AND CONDITION OF KEY HABITAT: Mesic and wet forests dominated by ‘ōhi‘a and koa (*Acacia koa*), primarily at elevations greater than 1,250 meters (4,100 feet) but can be found as low as 150m (500ft). The primary reason for this limitation is the high density of cold-intolerant *Culex* mosquitoes, an important disease vector, below this elevation. Occupied habitats also contain kōlea (*Myrsine lessertiana*), naio (*Myoporum sandwicense*), and hapu‘u tree ferns (*Cibotium* spp.). Māmane (*Sophora chrysophylla*) is common in high-elevation foraging habitat. Although much of the species’ current range is under State or Federal jurisdiction, habitat protection and restoration efforts vary considerably.

THREATS: Although populations appear stable on the islands of Hawai‘i, Maui, and Kaua‘i, they are susceptible to the same factors that threaten other native Hawaiian forest birds including habitat loss and degradation, predation by introduced mammals, and disease. For ‘apapane the following is of particular concern:

- **Disease.** Of Hawai‘i’s native forest birds, ‘apapane have the highest prevalence of avian malaria. Individuals infected with avian pox also are more likely to be infected with malaria. Foraging movements may increase their exposure to disease. ‘Apapane breed in mid-elevation forests, which suggests some disease resistance.
- **Predation.** Predation on adults and nests by rats (*Rattus* spp.), cats (*Felis catus*), the small Indian mongoose (*Herpestes auropunctatus*), and owls (Pueo; *Asio flammeus sandwichensis*, Barn owl; *Tyto alba*) may limit the species. Likely because ‘apapane are so abundant, observations have been made of direct depredation events by these predators.

CONSERVATION ACTIONS: ‘Apapane likely benefit from actions to conserve other endangered forest birds on northeastern Haleakalā, Hakalau Forest National Wildlife Refuge, Hawai‘i Volcanoes National Park, the ‘Ōla‘a/Kīlauea Watershed Partnership, and Alaka‘i Wilderness Preserve and surrounding areas. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. Future actions specific to the protection of ‘apapane may include the following:

- Control mosquitos and non-native predators across native forest habitat.
- Conduct public education and outreach.
- Continue protection and management of wildlife sanctuaries and refuges.
- Encourage more native tree planting in backyards, community parks, and landscaping.

MONITORING: Continue traditional forest bird surveys, expand use of automatic recording units (ARU), and habitat monitoring on all islands.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to ‘apapane include the following:

- Determine if disease-resistant individuals exist and if so, if resistance is passed to offspring. Disease-resistant birds could be used to found new populations.
- Determine the role of ‘apapane in disease transmission between high- and low-elevation habitats.
- Conduct life history studies to quantify the population structure, dispersal patterns, survivorship, nesting phenology, and success of this poorly known species, especially with small populated locations like Lāna‘i.
- Explore reintroducing ‘apapane to the Papahānaumokuākea Marine National Monument.
- Conduct range wide population estimates and determine population trends.
- Evaluate metrics and requirements to allow for reestablishment of cultural use of the species.

References:

- Fancy SG, Ralph CJ. 1997. ‘Apapane (*Himatione sanguinea*). In The Birds of North America, No. 296 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org.
- Judge SW, Camp RJ, Warren CC, Berthold LK, Mounce HL, Hart PJ, Monello RJ. 2019. Pacific island landbird monitoring annual report, Haleakalā National Park and East Maui Island, 2017. Natural Resource Report. NPS/PACN/NRR—2019/1949. National Park Service. Fort Collins, Colorado.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Smith L, Judge S, Camp R, Genz A, Berry L, Berthold L, Mounce H. 2023. The Status and Trends of Forest Birds on Moloka‘i. Poster Presentation. Hawai‘i Conservation Conference. Honolulu, HI.

Forest Birds

Moloka‘i ‘Ō‘ō

Moho bishopi

SPECIES STATUS:

Federally Listed as Extinct

State Recognized as Extinct

NatureServe Heritage Rank GH—Possibly Extinct

IUCN Red List Ranking—Extinct

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: Known only from Moloka‘i, Bishop’s ‘ō‘ō is a large, noisy honeyeater (Family: Meliphagidae). This striking species is black with yellow ear patches, under tail coverts, and maxillary tufts; sexes are similar. The bird’s vocalizations have been described as varied and “unlike any other native bird.” Bishop’s ‘ō‘ō appears to be primarily nectarivorous, preferring lobelia (Campanulaceae) flowers. Little is known about this species’ life history and nothing is known about its nesting biology.


DISTRIBUTION: Unknown. Probably extinct. Historic range of Bishop’s ‘ō‘ō likely included all native forests of eastern Moloka‘i. Subfossils suggest it may have occurred on Maui. Sightings in the 1980s of a possible ‘ō‘ō species on Maui were never confirmed.

ABUNDANCE: Bishop’s ‘ō‘ō was last observed in 1904 and is probably extinct. No information on historical abundance.

LOCATION AND CONDITION OF KEY HABITAT:

Unknown. Bishop’s ‘ō‘ō occupied the montane forests of eastern Moloka‘i. The areas where the species was last observed are managed by the State of Hawai‘i as a Natural Area Reserve or by private conservation entities (e.g., The Nature Conservancy) as a Natural Area Partnership Preserve.

THREATS: Unknown. However, Bishop’s ‘ō‘ō likely were susceptible to the same factors that threaten other native Hawaiian forest birds including: loss and degradation of habitat, predation by introduced mammals, and disease, and possible human exploitation. For Bishop’s ‘ō‘ō populations, the following likely were of particular concern:



- Predation. The introduction of black rats in Hawai‘i seems to have been the primary early driver of extinction. Rats were first introduced to O‘ahu and Maui, and then around the late 1890s were introduced to Moloka‘i and Hawai‘i resulting in rapid waves of extinction on both islands. ‘Ō‘ō may have also nested in cavities making them even more susceptible to the highly arboreal pest.
- Disease. The fact that no habitat above 1,250 meters (4,100 feet) occurs on Moloka‘i suggests disease may have played an important role in the species’ decline.

CONSERVATION ACTIONS: If the species persists, it likely benefits from management activities to conserve other forest birds on eastern Moloka‘i, including the establishment and management of protected areas, regular surveys of forest bird populations, monitoring of habitat conditions, and studies of disease and disease vectors. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continuing to protect and manage wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to Bishop’s ‘ō‘ō.

References:

- I.A.E. Atkinson. 1977. A Reassessment of Factors, Particularly *Rattus rattus* L., That Influenced the Decline of Endemic Forest Birds in the Hawaiian Islands. University of Hawaii Press
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Sykes PW, Kepler AK, Kepler CB, Scott JM. 2000. Kaua‘i o‘o (*Moho braccatus*), O‘ahu o‘o (*Moho apicalis*), Bishop’s o‘o (*Moho bishopi*), Hawai‘i o‘o (*Moho nobilis*), and kioea (*Chaetoptila angustipluma*). In *The Birds of North America*, No. 535 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

Hawai'i 'ākepa

Loxops coccineus

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic


NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The Hawai'i 'ākepa is a small, insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Hawai'i. Separate species of 'Ākepa also are known from Maui (*L. ochraceus*) and O'ahu (*L. rufus*); both of which are likely extinct.. After three-four years, males obtain their bright orange adult plumage; subadult plumage is dull brownish orange, although individual variation is high. Females are grayish-green with a yellow-orange breast band. The lower mandible of the 'ākepa is slightly bent to one side which results in the mandible tips being offset; a characteristic shared with the 'akeke'e (*L. caeruleirostris*). The bend can be to the left or right, and depending on the direction of the bend, individuals also possess an accompanying leg asymmetry; the leg opposite the curve in the mandible is slightly longer than the other leg. Together, these adaptations likely improve the species foraging efficiency in the tightly bunched liko or leaf buds of 'ōhi'a. They often join mixed-species foraging flocks, particularly those with 'alawī (*Loxops mana*). They feed mainly on 'ōhi'a (*Metrosideros polymorpha*) leaf clusters, but also on koa (*Acacia koa*) leaves and seed pods, where it uses its bill to pry open leaf and flower buds in search of small arthropods. 'Ākepa are obligate cavity nesters, with most nests placed in natural cavities found in old-growth 'ōhi'a and koa trees. Females build nests, incubate eggs, and brood nestlings, and males deliver food to the female on and off the nest. Both parents feed the young, which remain with their parents for two to three months after fledging.

DISTRIBUTION: Occurs in four disjunct populations above 1,300 meters (4,300 feet) elevation. The majority of the population is found on the windward side of Maunakea within the Hakalau Forest National Wildlife Refuge and the Hilo Forest Reserve, but substantial



populations persist in windward Mauna Loa, and within the Ka‘ū Forest Reserve. A small remnant population of ~50 individuals also persists within the Pu‘u Wa‘awa‘a Forest Bird sanctuary. Original range likely included all forested regions of the island.

ABUNDANCE: The Hawaiian Forest Bird Survey (1976-79, 1983), estimated the population at $14,000 \pm 2500$ (95% confidence interval) birds. Judge et al. in 2018 using combined totals from Camp et al. 2009, Gorresen et al. 2005, and Camp et al. 2016 to estimate that the total range wide population for the species in 2016 was 16,428 (10,074 – 25,198). As of 2024 Populations in Ka‘ū and within the ‘Ōla‘a-Kīlauea region appear to have declined, however populations in windward Maunakea appear stable with populations at Hakalau NWR alone estimated to be 7,221 (4,164 – 11,173). The south Kona and Hualālai populations were estimated at 660 ± 250 birds, but as of 2024 are only known from ~50 birds on Hualālai..

LOCATION AND CONDITION OF KEY HABITAT: Occurs in ‘ōhi‘a and ‘ōhi‘a/koa forests above 1,300 meters (4,300 feet). Density appears to be related to the number of available cavities, and because cavities primarily occur in older, large trees, old-growth forests may be preferred. The highest density of ‘ākepa occurs in the Pua ‘Ākala tract of Hakalau Forest National Wildlife Refuge, which has numerous large trees but a degraded understory. Many areas occupied by the species have been degraded by feral ungulates. Most of the current range of the Hawai‘i ‘ākepa is managed by State and Federal agencies or private conservation partnerships.

THREATS:

- Habitat degradation and loss. Logging and ranching has fragmented and reduced the amount of suitable habitat. Breeding density may be limited by nest-site availability and current levels of food availability may limit populations. In forest fragments, the large trees required for nesting may be more susceptible to windfall and desiccation. The slow growth rate of ‘ōhi‘a complicates management for ‘ākepa. In addition, habitat fragmentation may prevent or restrict natural re-colonization of former range.
- Disease. The Hawai‘i ‘ākepa is not found below 1,300 meters (4,300 feet), which suggests that it is particularly susceptible to mosquito-borne diseases.
- Predation. Cavity nests may be vulnerable to rat predation, although nest success is high at Pua ‘Ākala in the Hakalau Forest NWR, where rat densities are high.

CONSERVATION ACTIONS: Completed or ongoing actions specific to the Hawai‘i ‘ākepa include: demographic and reproductive studies have determined the importance of old-growth trees for nesting and that the species will use artificial cavities for nesting, and captive propagation techniques have been developed. In addition, Hawai‘i ‘ākepa likely benefit from management activities to conserve other endangered forest birds in Hakalau Forest National Wildlife Refuge, the Kona unit of the Hakalau Forest National Wildlife Refuge, ‘Ōla‘a/Kīlauea Watershed Partnership, Kapāpala Forest Reserve, and Pu‘u Wa‘awa‘a Wilderness Sanctuary. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the Hawai‘i ‘ākepa may include the following:

- Aerially broadcast rodenticides to increase nestling and adult female survival.
- Conduct public education and outreach.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the Hawai‘i ‘ākepa include:

- Continue studies designed to refine the suitability of artificial cavities and evaluate their potential to facilitate the establishment of new populations.
- Determine the factors affecting the growth form of regenerating ‘ōhi‘a and potential methods for protecting old-growth trees from wind and desiccation.
- Identify disease-resistant individuals. Determining if genetic markers or genotypes are associated with resistance would allow targeted translocations of individuals possessing this genotype into populations lacking disease resistance.
- Conduct range wide population estimates and determine population trends.

References:

- Judge, Seth & Camp, Richard & Hart, Patrick & Kichman, Scott. 2018. Population estimates of the Endangered Hawai‘i ‘Ākepa (*Loxops coccineus*) in different habitats on windward Mauna Loa. *Journal of Field Ornithology*. 89.
- Kendall SJ, Rounds RA, Camp RJ, and Genz AS. 2022. Forest Bird Populations at the Big Island National Wildlife Refuge Complex, Hawaii. Hawai‘i Cooperative Studies Unit Technical Report HCSU-102. University of Hawai‘i at Hilo, Hawaii, USA. 141 pages.
- Lepson JK, Freed LA. 1997. ‘Akepa (*Loxops coccineus*). In *The Birds of North America*, No. 294 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

Hawai'i 'amakihi

Chlorodrepanis virens

SPECIES STATUS:

State Listed Endangered on Lāna'i

State Recognized as Endemic

NatureServe Heritage Rank G3—Vulnerable

SPECIES INFORMATION: The Hawai'i 'amakihi is a small, generalist Hawaiian honeycreeper (Family: Fringillidae). Until 1995, the Hawai'i 'amakihi, and the O'ahu (*H. flavus*) and Kaua'i 'amakihi (*H. kauaiensis*) were considered a single species: the common 'amakihi (*H. virens*). Plumage of all species is similar; males are yellow-green to olive with black lores. Females are generally similar, but duller. All have decurved bills. Plumage of males is bright yellow-green above, and there is some inter-island variation, especially among females. The Hawai'i 'amakihi is brighter and smaller than the Kaua'i 'amakihi. Hawai'i 'amakihi are generalized foragers that glean arthropods from the leaves, blossoms, twigs, branches, and less frequently from tree trunks, ferns, and shrubs. Feeds on nectar predominately from the flowers of 'ōhi'a (*Metrosideros polymorpha*), māmane (*Sophora chrysophylla*), and native lobelias (Campanulaceae), but also forages on flowers of a number of other native and non-native plants. They also eat fruit from native and non-native plants, but predominately from pilo (*Coprosma* spp.). Forages alone, in pairs, in family groups, or in mixed flocks. Courtship behavior is somewhat complex and includes courtship chases, advertising displays, and courtship feeding. Pairs remain together for successive breeding seasons. Pair selects nest site; female builds an open-cup nest and lays two or three eggs. Only females incubate eggs and brood nestlings. Males deliver food to females who then feed nestlings. Fledglings are dependent on parents for up to three months. The Hawai'i 'amakihi usually raise two broods in a season.

DISTRIBUTION: Occurs as low as 300m and as high as 2,900 meters (1,000 – 9,500 feet) on Hawai'i, Maui and Moloka'i; not common below 500 meters (1,625 feet). Widely distributed on Hawai'i and Maui, rare on Moloka'i. Original range likely included all forested regions of the above islands as well as those on Lāna'i, where it was last seen in 1976.

ABUNDANCE: The Hawaiian Forest Bird Survey (1976-1983) estimated the population at $870,000 \pm 5,612$ (95% confidence interval) birds on the island of Hawai'i, $44,000 \pm 1,786$ birds on east Maui, $3,000 \pm 408$ on west Maui, and $1,800 \pm 357$ birds on Moloka'i. Populations on Hawai'i and Maui are probably stable; the Moloka'i population is probably declining.

Updated Maui nui numbers-

East Maui- $77,776 \pm 3,694$ individuals. Densities have declined in some locations on East Maui.

West Maui- did not have a density estimate in 2020 surveys but continue to be common, but not as widely distributed as ‘apapane.

Moloka‘i- $1,085 \pm 510$ birds.

LOCATION AND CONDITION OF KEY HABITAT: A range of habitats including native shrubland and dry, mesic, and wet forests in montane and subalpine communities. Densities are highest on the island of Hawai‘i in subalpine ‘ōhi‘a scrub in Ka‘ū, and in māmane/naio (*Sophora chrysophylla* and *Myoporum sandiwickense*) forests on Mauna Kea. ‘Amakihi also are common in koa (*Acacia koa*) reforestation areas at higher elevations. On Maui, they are common in subalpine dry communities dominated by ‘ōhi‘a, māmane, pūkiawe (*Styphelia tamieameiae*) and ‘a‘ali‘i (*Dodonea viscosa*). They also occupy some non-native tree plantations on Maui, near areas where native vegetation persists. Habitat on Moloka‘i is restricted to the ‘ōhi‘a forests of the eastern half of the island and in lower elevation hala (*Pandanus tectorinus*) dominated areas near Kalaupapa. The condition of this habitat varies considerably. Much of the species’ current range is under State or federal jurisdiction.

THREATS: Although populations appear stable they are susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease.

CONSERVATION ACTIONS: Hawai‘i ‘amakihi likely have benefited from management actions to conserve other endangered forest birds in the native habitat on East Maui, Hakalau Forest National Wildlife Refuge, Hawai‘i Volcanoes National Park, and the ‘Ōla‘a/Kīlauea Watershed Partnership. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. Future management specific to the Hawai‘i ‘amakihi may include the following:

- Control mosquitos and non-native predators across native forest habitat.
- Translocate captive-bred individuals to Lāna‘i and Kaho‘olawe.
- Conduct public education and outreach.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring. Conduct range wide population estimates and determine population trends.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Currently, the U.S. Geological Survey’s Biological Resources Division is conducting genetic analyses to determine the species’ phylogenetic status and examining the relationship between genetic diversity and disease resistance. Additional research priorities include the following:

- Quantify population structure, dispersal patterns, survivorship, nesting phenology and success, especially for Maui and Moloka‘i populations.

- Determine if competition with Japanese white-eyes (*Zosterops japonicus*) occurs, and if so, its effect on Hawai'i 'amakihi populations.
- Conduct translocation experiments using Hawai'i 'amakihi to help reestablish this and other Hawaiian honeycreeper populations.
- Evaluate metrics and requirements to allow for reestablishment of cultural use of the species.

References:

- Judge SW, Camp RJ, Warren CC, Berthold LK, Mounce HL, Hart PJ, Monello RJ. 2019. Pacific island landbird monitoring annual report, Haleakalā National Park and East Maui Island, 2017. Natural Resource Report. NPS/PACN/NRR—2019/1949. National Park Service. Fort Collins, Colorado.
- Fukunaga, K, Berthold L, Warren C, Mounce H. 2019. Tracking Hawai'i 'Amakihi (*Chlorodrepanis virens wilsoni*) in Nakula Natural Area Reserve on East Maui: home range estimation and evaluation of techniques for upcoming Kiwiku (Pseudonestor xanthophrys) conservation translocation. Poster Presentation. Hawai'i Conservation Conference. Honolulu, HI. Photo of poster. Poster wording.
- Lindsey GD, VanderWerf EA, Baker H, Baker PE. 1998. Hawai'i (*Hemignathus virens*), Kaua'i (*Hemignathus kauaiensis*), O'ahu (*Hemignathus chloris*) and greater 'amakihi (*Hemignathus sagittirostris*). In *The Birds of North America*, No. 360 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Smith L, Judge S, Camp R, Genz A, Berry L, Berthold L, Mounce H. 2023. The Status and Trends of Forest Birds on Moloka'i. Poster Presentation. Hawai'i Conservation Conference. Honolulu, HI



Forest Birds

‘Alawī

Loxops mana

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Ranking G2—Imperiled

IUCN Red List Ranking—Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The ‘Alawī is a small, inconspicuous Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Hawai‘i. Adults are predominately olive-green above, dull buff below, and have a dark gray mask extending around the eyes; males are brighter. Their similarity to Hawai‘i ‘amakihi (*Chlorodrepanis virens*), Hawai‘i ‘ākepa (*Loxops coccineus coccineus*), and introduced Japanese white-eyes (*Zosterops japonicus*) complicates field identification. Unlike many Hawaiian forest birds, their life history is well known. Outside the breeding season, they frequently join mixed-species foraging flocks and forages over home ranges that average 5-11 hectares (up to 17.3 acres). They glean insects, spiders, and other invertebrates from the branches, trunks, and foliage of live ‘ōhi‘a (*Metrosideros polymorpha*) and koa (*Acacia koa*) trees. During the breeding season, the species’ home range averages 4 to 7 hectares (10 – 17 acres) and a 10 – 20 meter (33 – 66 feet) territory around the nest is defended. Most nests are open cup structures, but about 15 percent are placed in cavities or in bark crevices. Females build nests, incubate eggs, and brood nestlings. Males deliver food to the female on and off the nest. Both parents feed the young for approximately one month. ‘Alawī re-nest after nest failures and pairs may raise two broods in a season. Nest success is very low, but adults have high annual survival.

DISTRIBUTION: Occurs in four disjunct populations above 1,500 meters (5,000 feet) on the island of Hawai‘i. Historically occurred across the island above 1,070 meters (3,500 feet) elevation. Last detected in the Pu‘u Wa‘awa‘a Forest Bird Sanctuary in 2016, and nearly extirpated from the Kona District. Extinct in Kohala.

ABUNDANCE: The Hawaiian Forest Bird Survey (1976-79, 1983), estimated the population at $12,500 \pm 2,000$ (95% confidence interval) birds. The largest population consisted of 9,888 (6,115 – 14,974) at Hakalau Forest in 2019. .

LOCATION AND CONDITION OF KEY HABITAT: Most commonly in mesic and wet forests dominated by ‘ōhi‘a and koa, with a subcanopy of ‘ōlapa (*Cheirodendron trigynum*),

pūkiawe (*Styphelia tameiameia*), ‘ōhelo (*Vaccinium* spp.), ‘akala (*Rubus hawaiiensis*), kōlea (*Myrsine* spp.), kāwa‘u (*Ilex anomala*), and hapu‘u tree ferns (*Cibotium* spp.). Habitat conditions vary across the species’ range, with much of it degraded by grazing ungulates, especially cattle and feral pigs. Most of the current range of the ‘alawī is within the boundaries of State and Federally owned lands.

THREATS:

- Predation. Nest success is very low (11 to 50 percent) and rat (*Rattus* spp.) predation may be partially responsible. ‘Alawī place their nests near the main trunks of trees which may facilitate predation by rats.
- Disease. The alawī’s absence below 1,350 meters (4,500 feet) elevation suggests that it may be particularly susceptible to mosquito-borne avian disease.
- Habitat loss and degradation. Logging and grazing ungulates have reduced, degraded, and fragmented suitable forest habitats. Habitat fragmentation may be a dispersal barrier preventing or restricting recolonization of the species’ former range.
- Competition. Competition with Japanese white-eyes (*Zosterops japonicus*) may negatively affect ‘alawī.

CONSERVATION ACTIONS: Past or ongoing actions specific to the ‘alawī include studies on productivity, recruitment, and survival, and development of captive propagation techniques. They likely have benefited from actions to conserve other endangered forest birds in the Hakalau Forest National Wildlife Refuge, the Kona unit of the Hakalau Forest National Wildlife Refuge, ‘Ōla‘a/Kīlauea Watershed Partnership, Kapāpala Forest Reserve, and Pu‘u Wa‘awa‘a Wildlife Sanctuary. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. Future management specific to ‘alawī may include the following:

- Reintroduce the ‘alawī to managed areas in their former range (e.g., Mauna Loa strip in Hawai‘i Volcanoes National Park).
- Control rodents to enhance nestling and female survival. Aerial broadcast of rodenticides would be the most effective method to treat broad areas.
- Increase public education to engender support for conservation of forest birds.
- Continue and expand protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the ‘alawī include:

- Determining the efficacy and health implications of broadcast rodenticide.
- Explore landscape use of predator control via lethal traps.
- Conduct range wide population estimates and determine population trends.
- Determine if ‘alawī can be a surrogate species for ‘akikiki.

References:

- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Kendall SJ, Rounds RA, Camp RJ, and Genz AS. 2022. Forest Bird Populations at the Big Island National Wildlife Refuge Complex, Hawaii. Hawai'i Cooperative Studies Unit Technical Report HCSU-102. University of Hawai'i at Hilo, Hawaii, USA. 141 pages.
- Lepson JK, Woodworth BL. 2001. Hawai'i creeper (*Oreomystis mana*). In The Birds of North America, No. 680 (A. Poole and Fr. Gill, eds.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Photo: Bret Nainoa Mossman

Forest Birds

Hawai'i 'elepaio

Chasiempis sandwichensis

SPECIES STATUS:

State Recognized as Endemic

NatureServe Heritage Rank G3—Vulnerable


IUCN Red List Ranking—Vulnerable

Revised Recovery Plan for Hawaiian Forest Birds—USFWS
2006

SPECIES INFORMATION: The Hawai'i 'elepaio is a small, adaptable monarch flycatcher (Family: Monarchiade) endemic to the island of Hawai'i. 'Elepaio also occur on Kaua'i (*C. sclateri*) and O'ahu (*C. ibidis*); the latter is federally listed as endangered. Adults are dark brown above and white below with variable amounts of brown streaking; there is considerable variation in pattern based on elevation and precipitation levels; males have black throats and females have white throats. The bird's name is derived from its primary song which is a shrill whistle given only by males. 'Elepaio forage in the air, on the ground, logs, rock crevices, snags, and all parts of trees. They use a diversity of foraging maneuvers likely dependent on habitat type: they capture arthropods by flycatching, glean while perched or hovering, and in direct pursuit, and may prefer 'ōhi'a (*Metrosideros polymorpha*) and kāwa'u (*Ilex anomala*) for foraging. Pairs remain together year-round, and long-term pair bonds are common; one pair was together for 11 years. Unlike Hawaiian honeycreepers, both males and females participate almost equally in all aspects of rearing. Finely woven cup nests are built in 'ōhi'a and in other trees in proportion to their availability. Clutch size is usually two, and second nests are attempted, often while fledglings from first are still being fed. Young are fed by parents for at least a month but remain on their natal territory for up to ten months.

DISTRIBUTION: Occurs in most forested areas above 600 meters (2,000 feet). Isolated populations occur in Kohala and on the western slope of Mauna Kea. Historical distribution likely included all forested areas of the island.

ABUNDANCE: The Hawaiian Forest Bird Surveys (1976-79, 1983) estimated the islandwide population at more than 270,000 birds. The island of Hawai'i contains four populations (~150,000 birds) in windward Hawai'i Island, ~12,080 birds in Kohala, one population in Kona and an isolated population on the south slope of Mauna Kea. The Kona and Maunakea



populations have both declined since the 1980s in some areas by as much as 90%. As of 2019 the population at Hakalau Forest National Wildlife Refuge was 21,833 birds with another 5,315 in the Kona unit.

LOCATION AND CONDITION OF KEY HABITAT: A variety of forest types and elevations, but most common in wet or mesic forests at higher elevations. Highest densities occur in ‘ōhi‘a or mixed ‘ōhi‘a-koa (*Acacia koa*) forests above 1,100 meters (3,600 feet). Much of the current range is managed by State and federal agencies or private conservation partnerships.

THREATS:

- Predation. On O‘ahu, predation by black rats (*Rattus rattus*) has been implicated in the loss of nests and death of adult females, and rat control in these populations resulted in large increases in nest success and in the survival of adult females.
- Disease. Avian pox reduces nesting success and adult survival. On O‘ahu, annual survival and reproductive success of birds with active pox lesions are lower compared to healthy birds; no information is available on the effect of avian malaria.
- Habitat loss and degradation. Historical habitat loss and degradation, especially at low elevations, is a major cause of declines. In Hakalau Forest National Wildlife Refuge, population densities are lower in degraded, open forests than in intact, dense forests.

CONSERVATION ACTIONS: Hawai‘i ‘elepaio likely have benefited from management actions to conserve other endangered forest birds species at Hakalau Forest National Wildlife Refuge, Hawai‘i Volcanoes National Park, Pu‘u Lā‘au, and the ‘Ōla‘a/Kīlauea Watershed Partnership. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the Hawai‘i ‘elepaio may include the following:

- Protect and restore high elevation native forests and eliminate feral ungulates and non-native invasive plants.
- Conduct public education and outreach about the benefits of rodent control.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the Hawai‘i ‘elepaio include the following:

- Continue to screen birds for disease resistance. If resistant birds are identified, translocation and/or captive propagation of these birds may help recover populations.
- Continue to develop techniques for captive propagation.
- Conduct range wide population estimates and determine population trends.

References:

Hawai‘i’s State Wildlife Action Plan
October 1, 2025

- Burnett K, Camp RJ, Hart PJ. 2021. Current Distribution and Abundance of Kohala Forest Birds in Hawai'i. *Journal of Field Ornithology*.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Kendall, S. J., R. J. Camp, A. Wang, L. Berry, and L. Nietmann. 2023. Forest bird populations at the Pu'u Wa'awa'a Forest Bird Sanctuary and Pu'u Wa'awa'a Forest Reserve, Hawai'i. Hawai'i Cooperative Studies Unit Technical Report 108. University of Hawai'i at Hilo. 57 pages.
- Kendall SJ, Rounds RA, Camp RJ, and Genz AS. 2022. Forest Bird Populations at the Big Island National Wildlife Refuge Complex, Hawaii. Hawai'i Cooperative Studies Unit Technical Report HCSU-102. University of Hawai'i at Hilo, Hawaii, USA. 141 pages.
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.
- VanderWerf EA. 1998. 'Elepaio (*Chasiempis sandwichensis*). In *The Birds of North America*, No. 344 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Forest Birds

‘I‘iwi

Drepanis coccinea

SPECIES STATUS:

Federally Listed as Threatened

State Listed as Endangered on O‘ahu, Moloka‘i, Lāna‘i

State Recognized as Endemic

NatureServe Heritage Rank G4/T1/TH—Apparently Secure/

Possibly Extinct on O‘ahu, Moloka‘i and Lāna‘i/Critically imperiled on Kaua‘i

IUCN Red List Ranking—Vulnerable

SPECIES INFORMATION: The ‘i‘iwi is one of the most beautiful of the extant Hawaiian honeycreepers (Family: Fringillidae). Both males and females are vermillion red, with a black tail and wings, and a long, decurved pink bill. Native Hawaiians created feather adornments such as ‘Ahu ‘ula using hundreds of thousands of ‘i‘iwi feathers; such pieces signified power, spirituality, and prestige. Like ‘apapane (*Himatione sanguinea*), ‘i‘iwi often fly long distances in search of flowering ‘ōhi‘a (*Metrosideros polymorpha*). ‘I‘iwi are important pollinators for a wide swath of native plants especially the rare and endangered mints and members of the bell-flower family (Mā‘ohi‘ohi, ‘ōha wai, hāhā, ‘ōpelu, etc). Their diet consists primarily of nectar from a variety of native and non-native flowers and the presence of non-native flowers may have contributed to increases in some populations. In addition to nectar, ‘i‘iwi also eat small arthropods. Both sexes defend small nesting territories and may defend important nectar resources. Courtship chases and feeding may precede breeding. Nest sites are in terminal branches of ‘ōhi‘a trees and both sexes build the open-cup nest. Only females incubate eggs (typically two) and brood young. Young are mostly provisioned by female; males feed females off the nest. Despite their widespread distribution, little is known about their life history.

DISTRIBUTION: Occurs above 1,250 meters (4,500 feet) elevation on the islands of Hawai‘i, Maui, and Kaua‘i; and may occur at reduced densities below. Limited to East Maui and only a few small areas within the ‘Alaka‘i Wilderness on Kaua‘i. Seemingly extinct on O‘ahu and Moloka‘i. Historically, ‘i‘iwi were common down to low elevations on all the Main Hawaiian Islands.

ABUNDANCE: The ‘I‘iwi (*Drepanis coccinea*) is vulnerable to disease and habitat loss due to Rapid Ohia Death and appears to be declining in many areas; as such it was listed as threatened by U.S. Fish & Wildlife Service in October 2017. The following island population estimates are based on Paxton et al. (2013): 543,009 ± 26,697 (95% confidence interval) birds on island of Hawai‘i, 50,252 ± 3,437 individuals on East Maui, 176 ± 74 on West Maui,^[000] and 2,551 ± 617 on Kaua‘i. However, preliminary data from 2023 suggests the Kaua‘i population declined to 161 (95% CI: 47-304), with a range of only 492 ha (1215 ac);DOFAW, unpublished data). They have

not been detected on Molokaʻi since 2010 and on Oʻahu since 2018. The population is declining in certain locations, but shows evidence of increases in some areas of Hawaiʻi Island and East Maui. The species' wide-ranging foraging complicates population estimates and the determination of long-term trends.

LOCATION AND CONDITION OF KEY HABITAT: Mesic and wet forest dominated by ʻōhiʻa and koa (*Acacia koa*). Loss and degradation of habitat and high densities of cold-intolerant *Culex* mosquitoes, an important disease vector, in lowland areas restrict most birds to elevations above 1,250 meters (4,100 feet). Habitats with the highest ʻiʻiwi densities also support kōlea (*Myrsine lessertiana*), naio (*Myoporum sandwicense*), and hapuʻu tree ferns (*Cibotium* spp.). Māmane (*Sophora chrysophylla*) is common in high-elevation foraging habitat. Although much of the species' current range is under State or Federal jurisdiction, habitat quality and habitat protection and restoration varies considerably.

THREATS: Although populations appear stable on the islands of Hawaiʻi and Maui, they are drastically declining on Kauaʻi and likely susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For ʻiʻiwi, the following is of particular concern:

- Disease. ʻIʻiwi are very susceptible to avian malaria and avian pox. Nine of ten individuals died within 37 days after receiving a single bite from mosquitoes infected with *Plasmodium*. Individuals infected with pox also are more likely to be infected with malaria. Because the highest points on Molokaʻi and Oʻahu are below 1,250 meters (4,100 feet), this susceptibility likely explains the severe population declines noted on these islands. Foraging movements may increase their exposure to disease.

CONSERVATION ACTIONS: ʻIʻiwi likely have benefited from actions to conserve other endangered forest birds on northeastern Haleakalā, Hakalau Forest National Wildlife Refuge, Alakaʻi Wilderness Preserve and surrounding areas, Hawaiʻi Volcanoes National Park, and the ʻŌlaʻa/Kīlauea Watershed Partnership. Future actions specific to the protection of ʻiʻiwi may include the following:

- Control mosquitoes across native forest habitat.
- Aggressively control ungulates to improve habitat quality, facilitate the recovery of degraded habitat, and potentially reduce breeding habitat for mosquitoes. Control of non-native plants should be part of forest restoration efforts.
- Eradicate or manage mosquito breeding habitat in areas where ʻiʻiwi occur. Investigate options to eradicate mosquitoes, including use of genetic manipulation.
- Suppress/control rats, feral cats, and barn owls in areas where ʻiʻiwi occur.
- Prevent the introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other predators on Kauaʻi.
- Prevent the spread of Rapid ʻōhiʻa Death in prime ʻiʻiwi habitat.
- Conduct public education and outreach.
- Continue protection and management of wildlife sanctuaries and refuges.
- Collection for conservation breeding program, especially of the Kauaʻi population.
- Supplementation of the Kauaʻi population via translocation from other islands.

- Reintroduce birds to Moloka‘i, O‘ahu, and other areas where they have been extirpated if habitat can be cleared of mosquitoes.

MONITORING: Continue forest bird surveys and habitat monitoring on all islands.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats (*Rattus* spp.) and feral cats (*Felis catus*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to ‘i‘iwi include the following:

- Determine if disease-resistant birds exist, and if so, determine if resistance is passed to offspring. Disease-resistant birds could be used to establish new populations.
- Determine the role of ‘i‘iwi in transmitting disease between low and high elevations.
- Conduct life history studies to quantify the population structure, dispersal patterns, survivorship, nesting phenology and success of this poorly known species.
- Investigate genetic structure and disease prevalence of populations in mid-to- lower elevations, such as the population within West Maui, which had individuals translocated from East Maui in 2001.
- Study the genetics of the Kaua‘i population to determine how it may differ since Kaua‘i population has the greatest distance separating it from other islands’ populations.
- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps)
- Assess the impact of introduced rodents on food resources (e.g., floral, invertebrates) important to ‘i‘iwi.
- Evaluate recovery priorities, metrics, and requirements to best allow for reestablishment of cultural use of the species.

References:

- Burnett K, Camp RJ, Hart PJ. 2021. Current Distribution and Abundance of Kohala Forest Birds in Hawai‘i. *Journal of Field Ornithology*.
- Becker CD, Massey G, Groombridge JJ, Hammond R. 2010. Moving ‘I‘iwi (*Vestiaria coccinea*) as a Surrogate for Future Translocations of Endangered ‘Akohekohe (*Palmeria dolei*). Technical Report #172.
- Fancy S G, Ralph CJ. 1998. ‘I‘iwi (*Vestiaria coccinea*). In *The Birds of North America*, No. 327 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists’ Union.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Kendall, S. J., R. J. Camp, A. Wang, L. Berry, and L. Nietmann. 2023. Forest bird populations at the Pu‘u Wa‘awa‘a Forest Bird Sanctuary and Pu‘u Wa‘awa‘a Forest Reserve, Hawai‘i. Hawai‘i Cooperative Studies Unit Technical Report 108. University of Hawai‘i at Hilo. 57 pages.

Judge SW, Camp RJ, Warren CC, Berthold LK, Mounce HL, Hart PJ, Monello RJ. 2019. Pacific island landbird monitoring annual report, Haleakalā National Park and East Maui Island, 2017. Natural Resource Report. NPS/PACN/NRR—2019/1949. National Park Service. Fort Collins, Colorado.

Paxton, EH, Gorresen, PM, Camp RJ. 2013. Abundance, distribution, and population trends of the iconic Hawaiian Honeycreeper, the ‘I‘iwi (*Vestiaria coccinea*) throughout the Hawaiian Islands: U.S. Geological Survey Open-File Report 2013-1150.

Smith L, Judge S, Camp R, Genz A, Berry L, Berthold L, Mounce H. 2023. The Status and Trends of Forest Birds on Moloka‘i. Poster Presentation. Hawai‘i Conservation Conference. Honolulu, HI.

USFWS 2023? New recovery plan for iiwi

VanderWerf EA, Rohrer JL. 1996. Discovery of an ‘I‘iwi population in the Ko‘olau Mountains of O‘ahu. ‘Elepaio 56:25–28

Forest Birds



Jeremy Snell Courtesy of Bernice Pauahi Bishop Museum

Kākāwahie

Paroreomyza flammea

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank GH—Extinct


IUCN Red List Ranking—Extinct

Revised Recovery Plan for Hawaiian Forest Birds—

USFWS 2006

SPECIES INFORMATION: The kākāwahie, or Moloka‘i creeper, is a small, sexually dichromatic, insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to the forests of eastern Moloka‘i. Males are scarlet red; females are a warm light brown on top and pale gray below with an orange red breast. Juvenile males are a patchy dull rusty color. The species’ Hawaiian name translates as “woodchopping” and apparently describes the species chipping call. Kākāwahie forage in groups, gleaning invertebrates from leaves, bark, and epiphytes in wet ‘ōhi‘a (*Metrosideros polymorpha*) forests. Little is known about the species’ breeding biology, but it is assumed to be similar to that of the Maui creeper (*P. montana*). First described in 1889, the last bird was observed in May 1963

DISTRIBUTION: Unknown. Probably extinct. Last observed on the west rim of Pelekunu Valley. Kākāwahie were common in native forests of eastern Moloka‘i at the end of the 19th century. Original range likely included all forested regions of Moloka‘i



ABUNDANCE: Unknown. The last kākāwahie was observed in 1963 and the species is probably extinct.

LOCATION AND CONDITION OF

KEY HABITAT: Unknown. Was known to occur in wet ‘ōhi‘a (*Metrosideros polymorpha*) forests from low to high elevations, and other heavily wooded native areas of eastern Moloka‘i. The areas where the species was last observed are managed by the State as

Natural Area Reserves or by The Nature Conservancy.

THREATS: Unknown. However, kākāwahie likely were susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For kākāwahie, the following likely was of particular concern:

- Disease. This species rapid decline and the fact that no habitat above 1,250 meters (4,100 feet) occurs on Moloka‘i suggests disease may have played an important role in the species’ decline.

CONSERVATION ACTIONS: If the species persists, it likely benefits from actions to conserve other endangered forest birds of eastern Moloka‘i including the establishment of the protected areas, regular surveys of forest bird populations, habitat monitoring, and studies of disease and disease vectors. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue Forest bird surveys and habitat monitoring on all islands.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to kākāwahie.

References:

- Baker PE, Baker H. 2000. Kākāwahie (*Paroreomyza flammea*) and O‘ahu alauahio (*Paroreomyza maculata*). In *The Birds of North America*, No. 503 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.

Forest Birds

Kāma'o

Myadestes myadestinus

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank GH – Possibly Extinct

IUCN Red List Ranking — Extinct

Y Plan for the Hawaiian Forest Birds—USFWS 2006




Jeremy Snell Courtesy of Bernice Pauahi Bishop Museum

SPECIES INFORMATION: The kāma'o, or large Kaua'i thrush, is one of two Hawaiian solitaires (family: Turdidae) endemic to Kaua'i. The species was noted for flying upward, singing a few loud notes, and then suddenly dropping into the understory. Like all adult Hawaiian solitaires, the kāma'o has olive-brown and gray plumage with rusty brown patches in the wings, but it lacks the white-eye ring and pinkish legs of the smaller puaiohi or small Kaua'i thrush (*M. palmeri*). The species' complex song is composed of a melodic series of liquid warbles, trills, and whistles, and was often heard before dawn and after dusk. The diet is reported to consist of fruits and berries.. Life history characteristics are mostly unknown, but are presumed similar to the 'ōma'o (*M. obscurus*). Breeding is thought to occur in spring, although no nest has been described.

DISTRIBUTION: Has not been observed below 1,100 meters (3,500 feet) since the mid-1960s. Extinct, but formerly, it was concentrated in the uppermost regions of the Alaka'i Wilderness Preserve. Historically was found in moist forests near sea level on northern Kaua'i as well as upland, interior mountain forests.

ABUNDANCE: Probably extinct. The Hawaiian Forest Bird Survey (1981), estimated the population at 24 ± 30 (SE) individuals. The last kāma'o was observed in 1989, and was not observed during the 2000, 2008, 2018, or 2023 Kaua'i Forest Bird Surveys. Historically, the kāma'o was extremely common.



LOCATION AND CONDITION OF KEY HABITAT: Most recent sightings were in open canopy forests of ‘ōhi‘a (*Metrosideros polymorpha*) and ‘ōlapa (*Cheirodendron* spp.). Based on the diet and life history of the ‘ōma‘o, a diverse understory including epiphytes, tree ferns, and fruiting plants such as ‘ie‘ie, ‘ōhā wai (*Clermontia* spp.), and ‘ōhelo (*Vaccinium* spp.) would likely be high-quality habitat. Because ‘ie‘ie, an important food plant, does not do well in high-elevation forests, if the species persists it may be restricted to marginal habitat. The area where the species was last observed is managed by the State of Hawai‘i as a Wilderness Preserve.

THREATS: Kāma‘o are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For kāma‘o, the following are of particular concern:

- Disease. Mosquito-borne disease is probably the most important factor in the decline of the kāma‘o. Pox lesions were noted on this species in mid-19th century.
- Habitat degradation. The presence of native forest with abundant fruit-bearing plants below the species current range demonstrates that habitat degradation cannot entirely explain the species extirpation from lowland areas. However, several invasive plants and feral pigs (*Sus scrofa*) have degraded the understory of many native forests.
- Competition. Non-native birds, especially ecologically similar species (e.g., white-rumped shama [*Copsychus malabaricus*]), may have contributed to the species’ decline.
- Predation. If kāma‘o, like many solitaires, are cavity or low platform nesters, their nests would be very susceptible to rats (*Rattus* spp.).
- Non-native arthropods. Recently introduced non-native insects, especially yellow jackets (*Vespula pensylvanica*) and Argentine ants (*Linepithema humile*), may compete with the kāma‘o’s native arthropod prey or disrupt the pollination of the species’ food plants. Introduced herbivorous insects also could reduce the abundance of food plants.

CONSERVATION ACTIONS: If the species persists, it likely benefits from efforts to conserve other endangered forest birds on Kaua‘i, including the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, habitat monitoring, studies of disease and disease vectors, and public education efforts featuring Kauai’s endangered forest birds. In addition to these efforts, future management specific to the recovery of the kāma‘o may include:

- Aggressively control ungulates to improve the quality of kāma‘o habitat and facilitate the recovery of degraded, but potential, habitat. Control of non-native plants should be part of forest restoration efforts.
- Conduct eradication of rats and feral cats from the Alaka‘i Wilderness Preserve.
- Prevent introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other possible predators.
 - Conduct public outreach and education.
 - Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to

control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to kāmā‘o.

References:

- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.
- Wakelee KM, Fancy SG. 1999. ‘Oma‘o (*Myadestes obscurus*), kama‘o (*Myadestes myadestinus*), oloma‘o (*Myadestes lanaiensis*), and ‘amaui (*Myadestes woahensis*). In *The Birds of North America*, No. 460 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.

Forest Birds



Jeremy Snell Courtesy of Bernice Pauahi Bishop Museum

Kaua'i 'akialoa

Akialoa procerus

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank GX—Presumed Extinct

IUCN Red List Ranking—Extinct


Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The Kaua'i 'akialoa is perhaps the most morphologically specialized of the Hawaiian honeycreepers (Family: Fringillidae) having a decurved bill that is up to half the length of their body. Both sexes are mostly olive-green; males being somewhat brighter, slightly larger, and have a longer bill. The life history of the Kaua'i 'akialoa is poorly known and mostly based on observations from the turn of the last century. The species principally foraged for arthropods on the trunks and branches of 'ōhi'a (*Metrosideros polymorpha*) and koa (*Acacia koa*) trees, and hāpu'u tree ferns (*Cibotium* spp.) by using its bill to probe bark crevices, decaying wood, epiphytes, and organic matter; observed to insert its entire bill into crevices. Foraging behavior has been described as being similar to that of a woodpecker or creeper. They also took nectar from 'ōhi'a and lobelia (Campanulaceae) flowers. Nothing is known about its breeding biology. 'Akialoa also occurred on the islands of Hawai'i, O'ahu, and Lāna'inone have been observed for at least 60 years.

DISTRIBUTION: Unknown. Probably extinct. The Kaua'i 'akialoa was last seen in the Alaka'i wilderness preserve. Original range likely included all forested regions of Kaua'i.

ABUNDANCE: Unknown. Last observed in 1969, and is probably extinct. Extensive surveys in 1989, 1994, 1996, 2000, 2005, 2006, 2008, 2018, and 2023 did not detect the species. Historically, the species was reported to be fairly common.

LOCATION AND CONDITION OF KEY HABITAT: Unknown. In the late



1800s, the Kaua‘i ‘akialoa occurred in most forests on Kaua‘i from between 200 and 1,500 meters (650 – 4,875 feet) elevation. The species was last observed in the Alaka‘i Wilderness Preserve.

THREATS: Causes of the decline of this species are unknown. However, ‘akialoa likely were susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For Kaua‘i ‘akialoa populations, the following likely was of particular concern:

- Disease. Avian pox lesions are noted in historic accounts and occur on museum specimens. Perkins (1903) noted Kaua‘i ‘akialoa was “grievously affected by...swelling on the legs and feet, as well as on the head at the base of the bill, and on the skin around the eyes.”
- Predation. Following the introduction of non-native mammals, particularly black rats, to Kaua‘i in the late 1800s or early 1900s this species and most other Kaua‘i endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Hawai‘i, Maui, Moloka‘i, and Lāna‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: If the species persists, it likely benefits from actions to conserve other endangered forest birds on Kaua‘i, including the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, habitat monitoring, studies of disease and disease vectors, control of feral ungulates through public hunting, and public education efforts featuring Kauai’s endangered forest birds. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to Kaua‘i ‘akialoa.

References:

- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. *Conservation Biology* 18:716-725.
- 0Atkinson I. A. E. 1977. A Reassessment of Factors, Particularly *Rattus rattus*., That Influenced the Decline of Endemic Forest Birds in the Hawaiian Islands. *Pacific Science*.

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Lepson JK, Johnston SM. 2000. Greater 'akialoa (*Hemignathus ellisianus*) and lesser 'akialoa (*Hemignathus obscurus*). In *The Birds of North America*, No. 512 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.

Perkins RC. 1903. Vertebrata. Pp. 365-466 In *Fauna Hawaiiensis*. Volume 1, Part IV (Sharp D, editor). Cambridge, UK: University Press.

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

Kaua'i 'amakihi

Chlorodrepanis kawaiensis


SPECIES STATUS:

State Recognized as Endemic
NatureServe Heritage Rank G3—Vulnerable
IUCN Red List Ranking—Endangered

SPECIES INFORMATION: The Kaua'i 'amakihi is a small, generalist Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Kaua'i. Until 1995, the Kaua'i 'amakihi, Hawai'i (*H. virens*), and O'ahu 'amakihi (*H. falvus*) were considered a single species: common amakihi (*H. virens*). The plumage of all species is similar; males are yellow-green to olive with black lores. Females are generally similar, but duller. All have decurved bills. The Kaua'i 'amakihi is the duller of the three species, is larger and has a longer, heavier and more decurved bill than the other species. They glean arthropods from branches and trunks as well as leaves, often hanging upside down to examine the underside of branches. Also probes bark crevices for food items, pierces the bases of 'ōhi'a (*Metrosideros polymorpha*) flowers for nectar, and feeds on the fruit of native and non-native plants. Males display (i.e., rapidly move around the female and sing), chase, and feed females prior to breeding. Both sexes build the nest, typically in a non-blooming 'ōhi'a tree, although the female does most of the construction while the male sings in adjacent trees. Kaua'i 'elepaio nests are lower than most other forest bird species on Kaua'i at 5.1 ± 2.0 m, ($n = 40$, range 1.3–8.4 m; Hammond et al. 2015). Females incubate a clutch of three eggs and brood nestlings at night and during inclement weather. Both parents feed nestlings. The mean incubation period of Kaua'i 'elepaio is 19.7 ± 1.5 d ($n = 23$, range = 17–23 and the mean nestling period is 15.0 ± 1.2 d ($n = 24$, range = 13–17 d, for a total nesting period of 35 d (Hammond et al. 2015). Fledglings are accomplished fliers. No information on post-fledgling behavior or parental dependency. Although weather has been implicated in nest failure, nest survival probability is high (0.63 ± 0.08), but of nests that fail, predation is the cause in 60% of cases (Hammond et al. 2015).

DISTRIBUTION: Occurs above 600 meters (2,000 feet) in the forests of Waimea Canyon, Nā Pali Plateau, the Alaka'i Swamp, and Makaleha Mountains. Original range likely included all forested areas of Kaua'i

ABUNDANCE: In the early 1970s the population was estimated at $10,743 \pm 970$ (SE) birds. A survey in the late 1980s estimated 15,000 to 20,000 birds. The Kaua'i Forest Bird Survey (2000) estimated the population in the Alaka'i Swamp and Kōke'e State Park area at greater than 40,000 birds and reported a significant population increase between 1981 and 2000. However, data from the Alakai Plateau suggest a general decline and possible range constriction since 2008, and in 2023 the population was estimated to be 9,307 (95% CI: 6,852-11,588; DOFAW, unpublished data).



LOCATION AND CONDITION OF KEY HABITAT: Wet and mesic montane forests above 600 meters (2,000 feet) dominated by 'ōhi'a, koa (*Acacia koa*), 'ōlapa (*Cheirodendron trigynum*), and lapalapa (*C. platyphyllum*). At lower elevations where the species historically occurred, native habitats are severely degraded. Although public hunting reduces the number of feral ungulates in accessible areas of the species' range, it is not effective in preventing habitat degradation. Occupied habitats above Waimea Canyon, in and west of the Alaka'i Swamp, are managed by the State of Hawai'i.

THREATS: Kaua'i 'amakihi are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. Although many Kaua'i 'elepaio tolerate malaria well enough to be captured post-infection, positive malaria status is negatively associated with body condition and the species sometimes shows pox lesions (KFBRP, unpubl. data).

CONSERVATION ACTIONS: Kaua'i 'amakihi likely benefit from actions to conserve other endangered forest birds including establishment of the Alaka'i Wilderness Preserve, regular surveys of forest bird populations, habitat monitoring, studies of disease and disease vectors, control of feral ungulates through public hunting control of feral ungulates through public hunting and fencing, suppression of rats on two A24 trapping grids over approximately 150 ha (470 ac), and education about Kauai's endangered forest birds. Future management specific to the Kaua'i 'amakihi may include the following:

- Aggressively control ungulates to improve habitat quality.
- Control non-native plants that degrade native habitat.
- Reduce breeding habitat for mosquitoes on the Alaka'i Plateau
- Suppress mosquito populations through releases of incompatible or sterile mosquitoes.
- Investigate options to eradicate mosquitoes, including use of genetic manipulations.

- Prevent the spread of Rapid Ohia Death on the Alaka'i Plateau
- Suppress/control rats (*Rattus* spp.), feral cats (*Felis catus*), and barn owls (*Tyto alba*) from the Alaka'i Wilderness Preserve and Koke'e Region.
- Prevent the introduction of the small Indian mongoose (*Herpestes auropunctatus*) and other non-native predators.
- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquitoes. Research priorities specific to the Kaua'i 'amakihi include the following:

- Conduct life history studies to quantify the population structure, dispersal patterns, survivorship, nesting phenology, and success of this poorly known species.
- Conduct studies to investigate the apparent population fluctuations in this species
- Determine the species' susceptibility to avian malaria and avian pox.
- Investigate the impact of rat predation on food resources (e.g., flowers, invertebrates) for 'amakihi.
- Investigate sub-lethal impacts disease (malaria, pox) on 'amakihi fecundity
- Determine sources of mosquitoes and investigate methods of mosquito control.
- Determine the effects of recently established non-native insects on native arthropods, especially those that are part of the species' diet. Determine the status of populations outside of the greater Alaka'i Swamp region.
- Conduct phylogenetic analyses to determine the relationship to other 'amakihi species.

References:

- Camp, RJ and PM Gorresen. 2011. Design of forest bird monitoring for strategic habitat conservation on Kaua'i Island, Hawai'i. Hawai'i Cooperative Studies Unit Technical Report HCSU-022. University of Hawai'i at Hilo.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka'i swamp, Kaua'i. *Conservation Biology* 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Lindsey GD, VanderWerf EA, Baker H, Baker PE. 1998. Hawai'i (*Hemignathus virens*), Kaua'i (*Hemignathus kauaiensis*), O'ahu (*Hemignathus chloris*) and greater 'amakihi (*Hemignathus sagittirostris*). In *The Birds of North America*, No. 360 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.



Forest Birds

Kaua'i 'elepaio


Chasiempis sclateri

SPECIES STATUS:

State Recognized as Endemic
NatureServe Heritage Rank G2—Imperiled
IUCN Red List Ranking—Vulnerable

SPECIES INFORMATION: The Kaua'i 'elepaio is a small, adaptable monarch flycatcher (Family: Monarchiade) endemic to the island of Kaua'i. 'Elepaio also occur on the islands of Hawai'i (*C. sandwichensis*) and O'ahu (*C. ibidis*); the latter is federally listed as endangered. Adults have a dark grayish brown crown and back and white underparts with a rusty wash on the upper breast. The bird's name is derived from its primary song which is a shrill whistle given only by males. Like other 'elepaio on Hawai'i and O'ahu, Kaua'i 'elepaio use virtually all available substrates for foraging, including the ground, logs, rock crevices, snags, and all parts of trees. Equally diverse in the use of foraging maneuvers, 'elepaio capture a wide range of arthropod prey by flycatching, gleaning while perched or hovering, and direct pursuit; foraging maneuvers vary depending on plant species from which prey is being captured and habitat type. For Kaua'i, there is little information on plant species used, although 'ōhi'a (*Metrodiseros polymorpha*) is likely and nests in 'alani (*Melicope* spp.) are common. On the island of Hawai'i, pairs remain together throughout the year and long-term pair bonds are common. Unlike Hawaiian honeycreepers, both males and females participate almost equally in all aspects of rearing. Finely woven, cup nests are built in a variety of native and non-native trees. Clutch size is usually two and second nests are attempted, often while fledglings from first are still being fed. Young are fed by parents for at least a month, but remain on their natal territory for up to ten months, which may allow young birds to hone their foraging skills.

DISTRIBUTION: Widely distributed above 600 meters (2,000 feet) elevation, but most common above 1,100 meters (3,600 feet) elevation on the Alaka'i Plateau. Original range likely included all forested regions of Kaua'i.



ABUNDANCE: In 1984, the island-wide population was estimated at 40,000 birds. The 2000 Kaua‘i Forest Bird Survey estimated the population in the Alaka‘i and Kōke‘e region at nearly 25,000 birds, with no change in the population since 1973. In 2018, the estimated population size was 51,903 birds (95% CI: 29,203-76,403, Paxton et al. 2020). In 2023, the population was estimated around 36,255 (95% CI: 29,894-43,463; U.S. Geological Survey, unpublished data). Unlike other Kaua‘i forest bird species, which have experienced steep overall population declines over the last 40 years, reduced Kaua‘i ‘elepaio densities in Kōke‘e have been compensated by increased densities in the Alaka‘i Wilderness Area during the same time period (DOFAW, unpubl., Paxton et al. 2020). Densities peak in ‘ōhi‘a forest between 1,100 and 1,300 meters (3,600 - 4,500 feet).

LOCATION AND CONDITION OF KEY HABITAT: Most common in dense wet ‘ōhi‘a forests above 1,100 meters (3,600 feet) elevation; uncommon in the drier forests of Waimea Canyon, the Nā Pali coast, and the wet bogs of Wai‘ale‘ale. ‘Elepaio are found in some low elevation valleys with a mix of native and non-native tree species. Common non-native trees in this habitat include Java plum (*Syzygium cumini*), coffee (*Coffea arabica*), kukui (*Aleurites moluccana*), and *Albizia lebbbeck*. Habitat quality varies considerably. The core of their range, the Alaka‘i Plateau and Kōke‘e State Park, is managed by the State of Hawai‘i as Forest Reserve, Wilderness Preserve, and State Park lands.

THREATS: Kaua‘i ‘elepaio are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For Kaua‘i ‘elepaio, the following are of particular concern:

- Disease. Avian pox is known to reduce both nesting success and adult survival. On O‘ahu, annual survival and reproductive success of birds with active pox lesions are lower compared to healthy birds; no information on the effect of avian malaria.
- Predation. On O‘ahu, predation by black rats (*Rattus rattus*) have been implicated in the loss of nests and death of adult females and on Hawai‘i and Kaua‘i, the loss of nests (Banko et al. 2019, Hammond et al. 2015). Rat control in these populations resulted in large increases in nest success and in the survival of adult females. Other predators, including cats (*Felis catus*), Hawaiian short-eared owls (*Asio flammeus sandwichensis*), and barn owls (*Tyto alba*) occur throughout the forests of Kaua‘i.

CONSERVATION ACTIONS: Given the increased population densities in the interior of Kaua‘i, where most management occurs, Kaua‘i ‘elepaio likely benefited from management actions to conserve other endangered forest birds including establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, eradication of ungulates within ungulate-proof fences, suppression of rats on A24 trapping grids, studies of disease and disease vectors, and public education efforts featuring Kauai’s endangered forest birds. In addition to these efforts, future management specific to the Kaua‘i ‘elepaio may include the following:

- Eradicate or control rats, feral cats, and barn owls in the Alaka‘i Wilderness Preserve.
- Prevent the introduction of the small Indian mongoose and other possible predators.
- Prevent the spread of Rapid ‘ōhi‘a Death on the Alaka‘i Plateau

- Suppress mosquito populations through releases of incompatible or sterile mosquitoes
- Investigate options to eradicate mosquitoes, including use of genetic manipulation.
- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquitos. Research priorities specific to the Kaua'i 'elepaio include life history studies to quantify population structure, dispersal patterns, survivorship, nesting phenology, and success of this poorly known subspecies.

- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps).
- Assess the impact of introduced rodents on food sources important to 'elepaio
- Investigate sub-lethal impacts disease (malaria, pox) on 'elepaio fecundity

References:

- Banko, Paul & Jaenecke, Kelly & Peck, Robert & Brinck, Kevin. (2019). Increased nesting success of Hawaii Elepaio in response to the removal of invasive black rats. *The Condor*. 121. duz003. 10.1093/condor/duz003.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka'i swamp, Kaua'i. *Conservation Biology* 18:716-725.
- Hammond RL, Crampton LH, Foster JT. 2015. Nesting success of native and introduced forest birds on the island of Kauai. *Journal of Avian Biology* 47: 252-262.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service. 428 pp.
- Paxton ER, Brinck KW, Crampton LH, Hite J, Constantini M. 2020. 2018 Kaua'i forest bird population estimates and trends. Technical Report Hawai'i Cooperative Studies Unit. University of Hawai'i at Hilo.
- VanderWerf EA. 1998. 'Elepaio (*Chasiempis sandwichensis*). In *The Birds of North America*, No. 344 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Forest Birds

Kaua'i nukupu'u *Hemignathus hanapepe*

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank G1T1—Critically Imperiled

IUCN Red List Ranking—Critically Endangered

(Potentially Extinct)


Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The Kaua'i nuku pu'u is a medium, short-tailed Hawaiian honeycreeper (Family: Fringillidae) with a long, thin decurved bill; the lower mandible is half the length of the upper mandible. Nukupu'u also are known from O'ahu (*H. lucidus*) and Maui (*H. affinis*); the O'ahu and Maui species are certainly extinct. Adult males are olive green with a yellow head, throat, and breast and have a small black mask; females are grayish green above and whitish below. Little is known about the species' life history. Often joins mixed species foraging flocks, especially those with 'akikiki (*Oreomystis bairdi*). Apparently would creep along tree trunks, especially those of ōhi'a (*Metrosideros polymorpha*) and koa (*Acacia koa*) trees, searching the bark and dead wood for arthropod prey; also may have taken nectar. Accounts vary regarding bill use. Either hammered surfaces, similar to its congener the 'akiapōlā'au (*H. munroi*), or used its upper mandible to fish out prey from crevices, catching them with its tongue and lower mandible. No information on the species' breeding biology, but likely is similar to the 'akiapōlā'au.

DISTRIBUTION: Unknown. Probably extinct. All recent Kaua'i nukupu'u sightings are from a small area of southwestern Kaua'i between 610 and 1,220 meters (2,000–4,000 feet) elevation. The species' historic range also appeared very restricted.

ABUNDANCE: Unknown. Probably extinct. There are fewer than a dozen historical records of the Kaua‘i nukupu‘u. Extensive surveys in 1989, 1994, 1996, 2000, 2005, 2008, 2018, and 2023 did not detect the species.

LOCATION AND CONDITION OF KEY HABITAT: Mesic and wet montane forests. Habitat conditions of the species’ historic range vary and all presumably support *Culex* mosquitoes. Areas where Kaua‘i nukupu‘u were most recently been observed are managed by the State of Hawai‘i.



THREATS: Unknown. However, the Kaua‘i nukupu‘u likely was susceptible to the same factors that threaten other native Hawaiian forest birds including habitat loss and degradation, predation by introduced mammals, and disease. For Kaua‘i nukupu‘u, the following likely was of particular concern:

- Disease. The precipitous decline of all nukupu‘u taxa suggests that mosquito-borne diseases played an important role in the species’ demise.
- Predation. Following the introduction of non-native mammals, particularly black rats, to Kaua‘i in the late 1800s or early 1900s this species and most other Kaua‘i endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Hawai‘i, Maui, Moloka‘i, and Lāna‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: If the species persists, it likely benefits from management activities to conserve other endangered forest birds on Kaua‘i, including the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, studies of disease and disease vectors, control of feral ungulates through public hunting, and public education efforts featuring Kauai’s endangered forest birds. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in

native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to Kaua‘i nukupu‘u.

- If mosquitoes can be sufficiently controlled, explore introducing ‘Akiapōlā‘au to Kaua‘i to fill the vacant niche.

References:

- Atkinson I. A. E. 1977. A Reassessment of Factors, Particularly *Rattus rattus*., That Influenced the Decline of Endemic Forest Birds in the Hawaiian Islands. Pacific Science.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. Conservation Biology 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Pratt TK, Fancy SG, Ralph CJ. 2001 ‘Akiapola‘au (*Hemignathus munroi*) and nukupu‘u (*Hemignathus lucidus*). In The Birds of North America, No. 600 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Photo: Mark Collins

Photo: Robert Shallenberger

Forest Birds

‘Ō‘ō‘ā‘ā

Moho braccatus

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank GH—Possibly Extinct

IUCN Red List Ranking—Extinct


Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: Endemic to its namesake, the Kaua‘i ‘ō‘ō is the smallest of the five known species of Hawaiian honeyeaters (Family: Meliphagidae), and has the least gaudy plumage of the four Hawaiian species of *Moho*. ‘Ō‘ō eat a variety of arthropods, snails, ‘ōlapa (*Cheirodendron* spp.) fruits, and nectar from the flowers of ‘ōhi‘a (*Metrosideros polymorpha*), lobelia, as well as other species. Early naturalists reported the species extensively feeding on the flower bracts of ‘ie‘ie (*Freycinetia arborea*), a species abundant in lowland forests, but not in upper elevation forests. Little is known about the species’ breeding biology. The only known nests were in cavities in large ‘ōhi‘a snags.

DISTRIBUTION: Unknown.

Possibly extinct. Was last observed in stream valleys of the central Alaka‘i Wilderness Preserve. Historically, the Kaua‘i ‘ō‘ō occurred in forest habitat throughout the island.

ABUNDANCE: Was last observed in 1987, and may be extinct. Extensive surveys in 1989, 1994, 1996, 2000, 2008, 2018, and 2023 did not detect the species. Was very common up to the end of 19th century.



LOCATION AND CONDITION OF KEY HABITAT: Unknown. The last sightings were in dense ‘ōhi‘a forests of the Alaka‘i swamp. Ironically, this habitat may have been low-quality or marginal habitat. ‘Ie‘ie, an important food plant, common in the lower elevation forests

previously occupied by ‘ō‘ō, is not found in the upper elevations forests where the species was last observed. Extensive damage to forests by hurricanes in 1982 and 1992 may have further reduced the suitability of high-elevation forests, especially given the species’ apparent dependence on large snags for nest sites. The area where the species was last observed is managed by the State of Hawai‘i as a Wilderness Preserve.

THREATS: Unknown. However, the species was likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For Kaua‘i ‘ō‘ō, the following likely were of particular concern:

- Disease. The precipitous decline of all Hawaiian *Moho* species suggests that disease played a role in this species’ decline.
- Predation. Following the introduction of non-native mammals, particularly black rats, to Kaua‘i in the late 1800s or early 1900s this species and most other Kaua‘i endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Hawai‘i, Maui, Moloka‘i, and Lāna‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: If the species persists, it likely benefits from management activities to conserve other endangered forest birds on Kaua‘i, including the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, studies of disease and disease vectors, control of feral ungulates through public hunting, and public education efforts featuring Kauai’s endangered forest birds. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the USFWS *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring on all islands.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct there are no research priorities specific to Kaua‘i ‘ō‘ō.

References:

- Atkinson I. A. E. 1977. A Reassessment of Factors, Particularly *Rattus rattus*., That Influenced the Decline of Endemic Forest Birds in the Hawaiian Islands. Pacific Science.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka‘i swamp, Kaua‘i. Conservation Biology 18:716-725.

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

Sykes PW, Kepler AK, Kepler CB, Scott JM. 2000. Kaua'i o'o (*Moho braccatus*), O'ahu 'o'o (*Moho apicalis*), Bishop's 'o'o (*Moho bishopi*), Hawai'i 'o'o (*Moho nobilis*), and kioea (*Chaetoptila angustipluma*). In *The Birds of North America*, No. 535 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.

U.S. Fish and Wildlife Service. 2006. Revised Recovery Plan for Hawaiian Forest Birds. Portland, (OR): U.S. Fish and Wildlife Service.

Forest Birds

Maui ‘ākepa

Loxops ochraceus

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank G1/TH— Presumed Extinct

IUCN Red List Ranking—Critically Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006



Jeremy Snell Courtesy of Bernice
Pauahi Bishop Museum

SPECIES INFORMATION: The Maui ‘ākepa is a small, insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of Maui. ‘Ākepa also are known from the island of Hawai‘i (*L. coccineus*) and O‘ahu (*L. rufus*); the latter subspecies is certainly extinct. Adults are less colorful compared to the Hawai‘i ‘ākepa, and adult males have two color morphs bright florescent orange and yellow. There is no information on the plumage sequence of the Maui species, however the adult males on Hawai‘i take 3-4 years to reach their definitive plumage. The lower mandible of the ‘ākepa is slightly bent to one side which results in the mandible tips being offset; a characteristic shared with the ‘akeke‘e (*L. caeruleirostris*) and the other ‘ākepa. Although almost nothing is known about its life history, the Maui ‘ākepa likely forages in a similar manner to the Hawai‘i ‘ākepa, and was most often observed in ‘ōhi‘a (*Metrosideros polymorpha*) forests. Although based on a single observation, in stark contrast to the Hawai‘i ‘ākepa, the Maui subspecies apparently builds an open-cup nest in the terminal foliage of ‘ōhi‘a trees. Declared extinct by U.S. Fish & Wildlife Service in 2023.


DISTRIBUTION: Unknown. Probably extinct. Last sightings occurred on the northeastern slopes of Haleakalā between 1,700 and 2,000 meters (5,500 – 7,000 feet) elevation. Historical distribution is poorly known, but like many Hawaiian forest birds the original range likely included all forested regions of the island.

ABUNDANCE: The Hawaiian Forest Bird Survey (1980), estimated the population at 230 ± 290 (95% confidence interval) individuals, but reliance on auditory detections may have biased this estimate. No ‘ākepa were found during the Hawai‘i Rare Bird Search (1995 - 1999). The last reliable sightings occurred in 1970, and the species has not been detected in subsequent surveys.

LOCATION AND CONDITION

OF KEY HABITAT: Virtually all sightings of the Maui ‘ākepa in this century were in wet montane ‘ōhi‘a

forests on the northeastern slopes of Haleakalā. These forests vary from intact to degraded. Areas where the Maui ‘ākepa have most recently been observed are managed by the State of Hawai‘i or by the National Park Service.



THREATS: Unknown. However, the Maui ‘ākepa was likely susceptible to the same factors that threaten other native Hawaiian forest birds including loss and degradation of habitat, predation by introduced mammals, and disease. For Maui ‘ākepa, the following was likely of particular concern:

- Disease. The precipitous decline of the Maui and O‘ahu subspecies and the restriction of the Hawai‘i ‘ākepa to high-elevation forests suggests that disease played an important role in the demise of Maui ‘ākepa.
- Predation. Following the introduction of non-native mammals, particularly black rats, to Maui in the early to late 1800s this species and most other Maui endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Hawai‘i, Maui, Moloka‘i, Lāna‘i, and Kaua‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions. If this species nested in cavities, it may have also been particularly susceptible to predators.

CONSERVATION ACTIONS: If the species persists, it likely benefits from actions to conserve other endangered forest birds on northeastern Haleakalā. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring and studies of disease and disease vectors. Should this species be rediscovered, the Rare Bird Recovery Protocol contained in the U.S. Fish and Wildlife Service (USFWS) *Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to the Maui ‘ākepa.

- Explore introducing Hawai‘i ‘ākepa to Maui to fill the now vacant niche left by this species.

References:

- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Lepson JK, Freed LA. 1997. ‘Akepa (*Loxops coccineus*). In The Birds of North America, No. 294 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Photo: Brooks Rownd

Forest Birds

Maui ‘alauahio

Paroreomyza montana

SPECIES STATUS:

State Recognized as Endemic


NatureServe Heritage Rank G4—Apparently Secure

IUCN Red List Ranking—Endangered

SPECIES INFORMATION: The Maui ‘alauahio, or Maui creeper, is a small insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to Maui. A subspecies also occurred on Lāna‘i but was last seen in 1937 and is presumed extinct. Adult males are predominantly olive-green above and have a bright yellow face, throat, and belly; the amount and intensity of yellow varies among individuals. Adult females are similar, but generally not as bright; both have short, fine straight bills. Adult plumage is not attained for several years. The Maui ‘alauahio gleans invertebrates from woody and leafy parts of a variety of plants. Adults defend an approximately 1-hectare (2.5 acre) home range against conspecifics year-round and will chase other birds and predators from the vicinity of their nests. They are socially monogamous and pair for life, although extra-pair copulations have been confirmed through genetic analysis. Females choose the nest site and build open-cup nests. Clutch size is two, and birds will renest after a failure, although double brooding has not been documented. Only females incubate eggs and brood nestlings. They do not breed until their third year, and young birds (i.e., helpers) associate with breeding pairs. Helpers are usually offspring from the previous year and feed the female, nestlings, and fledglings. Family groups can be as many as six individuals. Fledglings are fed for two to three months, and young remain with their parents in family groups for 18 - 20 months. They will make alarm calls when predators are in the area.

DISTRIBUTION: Broadly distributed in native and non-native forest on the slopes of Haleakalā at elevations ranging from 1200 to 2300 meters (3,900- 7,500 feet). . Historically common in West Maui and on Lāna‘i; these populations are now extirpated. Fossil evidence suggests they were common across the south side of East Maui and in lowland forests, but they no longer exist in these locations today. Range is estimated to be 107 square kilometers (41 square miles).

ABUNDANCE: The Hawai‘i Forest Bird Survey from 2022 estimated the population at 99,060 ± 9510 individuals (95% confidence interval); however surveys did not include all inhabited areas, so the population is likely higher. This estimate represents a 41% decrease since 1992.



LOCATION AND CONDITION OF KEY HABITAT: Primarily wet and mesic montane forests dominated by ‘ōhi‘a (*Metrosideros polymorpha*), although they also occur in subalpine māmane shrubland (*Sophora chrysophylla*), and in dry and mesic forests dominated by pine (*Pinus* spp.) and eucalyptus (*Eucalyptus* spp.) (e.g., Polipoli State Park and Hosmer Grove); all populations occur above 1200 meters (3,900 feet) elevation. Habitat conditions vary greatly across the species’ range. The species’ range is managed by the State of Hawai‘i, (i.e., Forest Reserve and Natural Area Reserve), the National Park Service, and private landowners including The Nature Conservancy. All entities are current members of the East Maui Watershed Partnership. Management efforts vary.

THREATS:

- Predation. Rats (*Rattus* spp.) and mongooses (*Herpestes auropunctatus*) have been observed depredating nests and females. Female behavior of begging near nests may make them particularly susceptible to rats. Additionally, their nests occur lower in the canopy than other honeycreeper species.
- Disease. Susceptibility to avian malaria has been documented, and likely prevents the establishment of populations in lowland areas.
- Habitat degradation. Current fencing around protected areas is not effective in excluding axis deer (*Axis axis*). Currently, deer populations on Maui are growing and threaten to further degrade forests occupied by the ‘alaupaho.


CONSERVATION ACTIONS: Maui ‘alauahio likely have benefited from actions to conserve endangered forest birds on northeastern Haleakalā including fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies on disease and disease vectors. In addition, ongoing fencing, ungulate control, and habitat restoration on Department of Hawaiian Homelands at Kahikinui could possibly allow for the re-establishment of a population there in the future. In general, actions should include continued protection and management of wildlife sanctuaries and refuges. Mosquito control implementation will benefit populations and could likely add to an increase in range and population and allow for reintroductions to areas where they used to be found. A possible reintroduction to West Maui, or Lāna‘i and translocations to O‘ahu and Moloka‘i could allow for an expansion of range as well as being a safeguard for the species.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for Hawaiian forest birds include improving methods for controlling rats, mongooses, and feral cats (*Felis silvestris*) in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquitoes. Research priorities specific to Maui ‘alauahio include development of a translocation protocol to facilitate reintroduction into restored forests. Diet studies to determine overlap with non-native species as well as determining targeted invertebrate recovery.

References:

- Baker H, Baker PE. 2000. Maui ‘alauahio (*Paroreomyza montana*). In The Birds of North America, No. 504 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Judge SW, Camp RJ, Warren CC, Berthold LK, Mounce HL, Hart PJ, Monello RJ. 2019. Pacific island landbird monitoring annual report, Haleakalā National Park and East Maui Island, 2017. Natural Resource Report. NPS/PACN/NRR—2019/1949. National Park Service. Fort Collins, Colorado.
- Judge, SW, Camp RJ, Warren CW, Berthold LK, Mounce HL, Hart PJ, and Monello RJ. 2021. Population Estimates and Trends of Three Maui Island-endemic Hawaiian Honeycreepers. Journal of Field Ornithology.
- Peck RW, Banko PC, Cappadonna J, Steele C, Leonard DL, Mounce HL, Becker CD, Swinnerton KJ. 2015. An assessment of arthropod prey resources at Nakula Natural Area Reserve, a potential site of reintroduction for Kiwīkiu (*Pseudonestor xanthophrys*) and Maui ‘Alauahio (*Paroreomyza montana*). Hawai‘i Cooperative Studies Unit, University of Hawai‘i at Hilo. Technical Report HCSU-059.
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Warren CC, Motyka PJ, Mounce HL. 2015. Home range sizes of two Hawaiian honeycreepers: implications for proposed translocation efforts. Journal of Field Ornithology 86(4):305–316.



Jeremy Snell Courtesy of Bernice

Forest Birds

Maui nukupu‘u

Hemignathus affinis

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank G1T1—Presumed Extinct

IUCN Red List Ranking—Critically Endangered

(Possibly Extinct)

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006


SPECIES INFORMATION: The Maui nukupu‘u is a large, short-tailed Hawaiian honeycreeper (Family: Fringillidae) with a long, thin decurved bill; the lower mandible is half the length of the upper mandible. Nukupu‘u also are known from O‘ahu (*H. lucidus*) and Kaua‘i (*H. hanapepe*); all three species are certainly extinct. Adult males are olive green with a yellow head, throat, and breast and have a small black mask; females are olive green above and variable yellow-gray below. Little is known about the species’ life history. Often joins mixed species foraging flocks. Apparently would creep along large ‘ōhi‘a (*Metrosideros polymorpha*) limbs searching epiphytes, moss, bark, and dead wood for arthropod prey; may also have taken nectar. Hammered bark with lower mandible, similar to its congener the ‘akiapōlā‘au (*H. munroi*), and used its upper mandible to fish out prey from excavations. No information on the species’ reproduction, but is likely similar to ‘akiapōlā‘au. Declared extinct by U.S. Fish & Wildlife Service in 2023.

DISTRIBUTION: Poorly known. Probably extinct. Most recent sightings between 1,100 and 2,100 meters (3,600–6,900 feet) elevation in the Kīpahulu Valley and the northeastern slope of Haleakalā. Historic range apparently very restricted, although subfossil evidence suggests the species may have occurred in dry forests.

ABUNDANCE: Unknown. Probably extinct. Based on a single sighting, the Hawaiian Forest Bird Survey (1980), estimated the population at 28 ± 56 (95% confidence interval) individuals. More recent surveys have failed to detect the Maui nukupu‘u. Historically considered uncommon.

LOCATION AND CONDITION OF KEY HABITAT:

Mixed ‘ōhi‘a/koa (*Koa acacia*) forests and mixed shrub montane wet forests between 1,100 and 2,100 meters (3,600–6,900 feet). Historic and fossil evidence indicates that its range was much broader and remnant populations may have been surviving in marginal habitat. Habitat conditions of the species’ former range vary. Areas where nukupu‘u were most recently sighted are managed as a Forest Reserve by the State of Hawai‘i or by the National Park Service.



THREATS: Unknown. However, the Maui nukupu‘u was likely susceptible to the same factors that threaten other native Hawaiian forest birds including loss and degradation of habitat, predation by introduced mammals, and disease. For Maui nukupu‘u, the following was likely of particular concern:

- Disease. The precipitous decline of all nukupu‘u taxa suggests that disease played an important role in the species’ decline.
- Predation. Following the introduction of non-native mammals, particularly black rats, to Maui in the early to late 1800s this species and most other Maui endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Hawai‘i, Maui, Moloka‘i, Lāna‘i, and Kaua‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: If the species persists, it likely benefits from actions to conserve other endangered forest birds on northeastern Haleakalā. These efforts include the establishment of the 3,000 hectares (7,500 acres) Hanawā Natural Area Reserve in 1986, the formation of the East Maui Watershed Partnership, fencing, ungulate and small mammal control, forest restoration, habitat monitoring and studies of disease and disease vectors. Should this species be rediscovered, the Rare Bird Recovery Protocol in the U.S. Fish and Wildlife Service (USFWS) *Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to Maui nukupu‘u.

- Explore the translocation of ‘Akiapōlā‘au to Maui to fill the vacant niche left by this species.

References:

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Pratt TK, Fancy SG, Ralph CJ. 2001 ‘Akiapola‘au (*Hemignathus munroi*) and nukupu‘u (*Hemignathus lucidus*). In The Birds of North America, No. 600 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service. 428 pp.



Photo: Ryan Wagner

Forest Birds

Kiwikiu

Pseudonestor xanthophrys

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered


State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking— Critically Endangered

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The kiwikiu or Maui parrotbill is a Hawaiian honeycreeper endemic to Maui, with a short tail and a relatively large, parrot-like bill. Adults are mostly olive-green above with a yellow breast, belly and cheeks, and a bright yellow line above their eyes (i.e., supercilium). Males are typically brighter than females, although individuals vary. Males are larger than females in size and bill. They feed on a variety of shrubs and small trees, especially ‘ākala (*Rubus hawaiiensis*), kanawao (*Broussaisia arguta*), ‘ōhi‘a (*Metrosideros polymorpha*), _kōlea (*Myrsine* species), pilo (*Coprosma* species) and hoi kuahiwi (*Smilax melastomifolia*). They use their large parrot-like bills to glean prey from moss-covered branches or chisel, crack, crush, dig, and tear bark and softer wood in search of beetle and Lepidoptera larvae and pupae. They also open fruit in search of insects. Pairs defend relatively large 11.8-to-14.5-hectare year-round home ranges. Females build nests, incubate eggs, and brood young. Clutch size is usually one, and females feed nestlings with food delivered by males. Males feed the fledglings. They will renest after a nest failure but are not known to attempt another nest during the breeding season if the first is successful. Development of bill and acquisition of foraging techniques is prolonged and young remain with parents for 5 to 18 months. Because of this long period of dependency, kiwikiu are often seen in small groups and males can be seen provisioning juveniles from current and previous years. Pairs are typically monogamous and invidious can live longer than 16 years.



DISTRIBUTION: Restricted to a ~30 square kilometer (11.5 square mile) on the northeastern slopes of Haleakalā between 1,600 and 2,200 meters (5,200 – 7,200 feet). Subfossils indicate they once occurred island-wide including at low elevations and leeward (southeastern) forests and on the island of Moloka‘i.

ABUNDANCE: The Hawai‘i Forest Bird Survey from 2022 estimated the population at 157 ± 67 (95% confidence interval) birds, a 71% decline in population since 2001

LOCATION AND CONDITION OF KEY HABITAT: Mid-to-upper-elevation montane wet forests dominated by ‘ōhi‘a, and in a few mesic areas dominated by ‘ōhi‘a and koa (*Acacia koa*), with a dense, diverse native understory and subcanopy of ferns, sedges, epiphytes, shrubs, and small to medium trees. Most of the range is managed by the National Park Service, State of Hawai‘i, The Nature Conservancy (TNC). Their range exists within the East Maui Watershed Partnership.

THREATS:

- Low reproduction. Unlike many Hawaiian honeycreepers, kiwikiu have low annual fledgling production. This results from a low reproductive potential (one fledgling per

year) coupled with low reproductive success due to habitat limitations and weather. This life history characteristic may be related to their very specialized foraging strategy. Regardless, the species is susceptible to factors that reduce population size.

- Disease. Despite the availability of seemingly suitable habitat below 1,600 meters (5,200 feet), kiwikiu are not found in these areas, suggesting that disease may be restricting populations to higher elevations. Furthermore, nearly all of the kiwikiu died of avian malaria during the 2019 translocation to the southeast slope of Haleakalā, even though the kiwikiu had come from and were taken to mid-to high elevation forest, which was typically known for lower disease rates, suggesting that disease in these forests is higher than previously thought.
- Predation. Predation on adults and nests by rats (*Rattus* spp.), cats (*Felis silvestris*), the small Indian mongoose (*Herpestes auropunctatus*), and owls (*Asio flammeus sandwichensis*, *Tyto alba*) may limit the species. High rat densities have been reported in the Hanawī area, which also supports a large proportion of the kiwikiu population.
- Habitat loss. Historical accounts suggest that kiwikiu favored koa for foraging. Logging and ranching have resulted in the loss of large areas of mesic koa forest, and their current range is restricted to wet forests where koa density is relatively low and wet weather can lead to nest failure. Thus like many endangered Hawaiian forest birds, kiwikiu may be restricted to suboptimal habitat.
- Habitat degradation. Damage to understory vegetation by feral pigs (*Sus scrofa*) likely reduces habitat suitability and may contribute to reduced food availability and low reproductive success. Habitat degradation also may increase exposure of nests to inclement weather.
- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former is usually most problematic. Demographic factors include skewed sex ratios and stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems.

CONSERVATION ACTIONS: The kiwikiu benefits from management efforts to conserve other endangered forest birds on northeastern Haleakalā, such as the establishment of the 3,000 hectare (7,500 acre) Hanawī Natural Area Reserve in 1986, the formation of East Maui Watershed Partnership and Maui Forest Bird Recovery Project, fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. Future actions specific to the recovery of the kiwikiu may include the following:

- Develop and deploy mosquito control techniques as soon as possible to reduce the mosquito vector of avian disease.
- Capture enough birds with a suitable age structure and sex ratio to establish a viable captive population. Continue to evaluate and adaptively manage captive care to maintain wild behaviors and encourage parental breeding behaviors. If/when mosquito control is deployed on Maui and meets efficacy success criteria, release worthy candidates back on Maui. In 1997, a captive breeding program was initiated. As of 2024, 8 kiwikiu are in captive care at the Maui Bird Conservation Center and efforts are ongoing to continue bringing suitable individuals into holding to prevent outright extinction in the wild from avian disease.

- Explore translocating birds to Hawai‘i Island to establish a 2nd population and reduce the species risk of extinction.
- Protect and restore habitat in high-elevation disease-free areas.
- Implement fencing and ungulate control in low-elevation habitat from the Hanawā Natural Area Reserve to TNC’s Waikamoi Preserve, to facilitate the recovery of the understory and subcanopy vegetation and eventually result in high-quality kiwikiu habitat. Once mosquito control has been implemented, this could be a possible location for kiwikiu to reestablish a population.
- Establish a continuous corridor of suitable habitat around Haleakalā by connecting conservation lands on the southern and western parts of the mountain. Restoration of koa forests to this area would be a key element to this effort.
- Restore, fence, and eradicate ungulates from the remnant mesic koa forests on the State Forest Reserve and Department of Hawaiian Home Lands in the Kahikinui region of southern Haleakalā. Restoration of this area would be a cost-effective starting point to providing the kiwikiu with high-quality habitat for future reintroductions.
- Conduct public outreach and education about the importance of invasive species control and forest restoration. Incorporate Hawaiian knowledge and practices in conservation strategies
- Continue protection and management of wildlife sanctuaries and refuges.
- Develop and deploy new tools and evaluate above approaches.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats, mongooses, and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquitoes. Research priorities specific to the kiwikiu include the following:

- Evaluate the effect of predator control on reproduction and survival of kiwikiu.
- Further refine captive breeding techniques and evaluate experimental reintroduction sites. Evaluation should include mosquito surveys and determination of disease prevalence in lower elevation sites and possible sites on other islands.
- Investigate invertebrate communities where kiwikiu do not currently inhabit to design and implement large-scale restoration. Diet studies to determine overlap with non-native species as well as determining targeted invertebrate recovery.

References:

- Brinck KW, Camp RJ, Gorresen PM, Leonard DL, Mounce HL, Iknayan KJ, Paxton EH. 2012. 2011 Kiwikiu (Maui Parrotbill) and Maui Alauahio abundance estimates and the effect of sampling effort on power to detect a trend. Hawaii Cooperative Studies Unit, University of Hawaii at Hilo. Technical Report HCSU-035.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Judge, S.W., R.J. Camp, C.W. Warren, L.K. Berthold, H.L. Mounce, P.J. Hart, and R.J. Monello. 2021. Population Estimates and Trends of Three Maui Island-endemic Hawaiian Honeycreepers. *Journal of Field Ornithology*.

- Peck RW, Banko PC, Cappadonna J, Steele C, Leonard DL, Mounce HL, Becker CD, Swinnerton KJ. 2015. An assessment of arthropod prey resources at Nakula Natural Area Reserve, a potential site of reintroduction for Kiwikiu (*Pseudonestor xanthophrys*) and Maui 'Alauahio (*Paroreomyza montana*). Hawai'i Cooperative Studies Unit, University of Hawai'i at Hilo. Technical Report HCSU-059.
- Mounce HL, Leonard DL, Swinnerton KJ, Becker CD, Berthold LK, Iknayan KJ, Groombridge JJ. 2013. Determining productivity of Maui Parrotbills, an endangered Hawaiian honeycreeper. *Journal of Field Ornithology* 84(1):32-39.
- Mounce, HL, Iknayan, KJ, Leonard, DL, Swinnerton, KJ, and Groombridge, JJ. 2014. Management implications derived from long term re-sight data: annual survival of the Maui Parrotbill *Pseudonestor xanthophrys*. *Bird Conservation International* 24:316-326
- Mounce HL, Raisin C, Leonard DL, Wickenden H, Swinnerton KJ, Groombridge JJ. 2015. Spatial genetic architecture of the critically-endangered Maui Parrotbill (*Psuedonestor xanthophrys*): management considerations for reintroduction strategies. *Conservation Genetics* Volume 16, Issue 1: Page 71-84.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Simon JC, Baker PE, Baker H. 1997. Maui parrotbill (*Pseudonestor xanthophrys*). In *The Birds of North America*, No. 311 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Warren CC, Motyka PJ, Mounce HL. 2015. Home range sizes of two Hawaiian honeycreepers: implications for proposed translocation efforts. *Journal of Field Ornithology* 86(4):305–316.
- Warren CC, Berthold LK, Mounce HL, Luscomb P, Masuda B, Berry L. 2021. 2019 Kiwikiu Conservation Translocation Report. Pacific Cooperative Studies Unit Technical Report #203. University of Hawai'i at Mānoa. Honolulu, HI. 103 pages.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

O'ahu 'alauahio

Paroreomyza maculata

SPECIES STATUS:

Federally Listed as Endangered

States Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank GH—Possibly Extinct

IUCN Red List Ranking—Critically Endangered


(Possibly Extinct)

Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The O'ahu 'alauahio is a small, sexually dichromatic, insectivorous Hawaiian honeycreeper (Family: Fringillidae) endemic to O'ahu. Males are olive-green above with bright yellow underparts and forehead, females are grayish-green above and yellowish-white below. The plumage of this species is very similar to that of the O'ahu 'amakihi (*Hemignathus flavus*). The song of the O'ahu 'alauahio has never been described. The O'ahu 'alauahio is insectivorous and forages by methodically searching and probing the bark of large branches and tree trunks. Like other Hawaiian creepers, 'alauahio joins foraging flocks during non-breeding seasons. Little is known about the life history or breeding biology of the species, but it is likely similar to that of the Maui species (*P. montana*).

DISTRIBUTION: Unknown. Probably extinct. Historical range is poorly known as it was likely declining when first discovered, but it historically occurred in both the Ko'olau and Wai'anae ranges.

ABUNDANCE: Unknown. Probably extinct. The last well-documented sighting was of two birds in 1985. O'ahu was not included in the Hawai'i Rare Bird Search in the late 1990s, so it is possible that the O'ahu 'alauahio still exists in remote valleys.



LOCATION AND CONDITION OF

KEY HABITAT: Unknown. Once occupied mid-elevation forests of 'ōhi'a (*Metrosideros polymorpha*) and koa (*Acacia koa*). Historically, it was reported to prefer large koa trees, but they also were reported in areas without koa. All recent observations occurred in mixed

Hawai'i's State Wildlife Action Plan
October 1, 2025

koa/‘ōhi‘a forest between 300 and 600 meters (1,000 – 2,000 feet). Forests where the O‘ahu ‘alauahio historically occurred are largely managed by the State of Hawai‘i, the U.S. Fish and Wildlife Service (USFWS), or the U.S. Military.

THREATS: Unknown. However, the O‘ahu ‘alauahio likely were susceptible to the same factors that threaten other native Hawaiian forest birds including habitat loss and degradation, predation by introduced mammals, and disease. For O‘ahu ‘alauahio, the following was likely of particular concern:

- Disease. The fact that no habitat above 1,250 meters (4,100 feet) occurs on O‘ahu, and that historical accounts report the species only above 350 meters (1,500 feet) suggests that disease played an important role in the species’ decline.
- Predation. Following the introduction of non-native mammals, particularly black rats, to O‘ahu in the early 1800s this species and most other endemics precipitously declined. Similar patterns of extinction and population decline were observed on Maui, Hawai‘i, Moloka‘i, Lāna‘i, and Kaua‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: Unfortunately, few forest birds remain on O‘ahu, and little if any, specific management is directed at forest birds other than the O‘ahu ‘elepaio (*Chasiempis sandwichensis ibidis*). The O‘ahu Forest National Wildlife Refuge in the Ko‘olau Range is near some of the most recent sightings, but whether the species still exists in the area is unknown. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the USFWS *Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Initiate regular forest bird surveys and habitat monitoring on O‘ahu.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats (*Rattus* spp.) and feral cats (*Felis silvestris*) in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is likely extinct, there are no research priorities specific to O‘ahu ‘alauahio.

References:

- Baker PE, Baker H. 2000. Kākāwahie (*Paroreomyza flammea*) and O‘ahu alauahio (*Paroreomyza maculata*). In The Birds of North America, No. 503 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.

Forest Birds



Photo: Bryan Shirota

O'ahu 'amakihi

Hemignathus flavus


SPECIES STATUS:

State Recognized as Endemic
NatureServe Heritage Rank G3—Vulnerable
IUCN Red List Ranking—Vulnerable

SPECIES INFORMATION: The O'ahu 'amakihi is a small, generalist Hawaiian honeycreeper (Family: Fringillidae) endemic to the island of O'ahu. Until 1995, the O'ahu 'amakihi, and the Hawai'i (*H. virens*) and Kaua'i amakihi (*H. kauaiensis*), were considered a single species: the common 'amakihi (*H. virens*). Plumage of all species is similar; males are yellow-green to olive with black lores. Females are similar, but duller. All have decurved bills. The plumage of some male O'ahu amakihi is variable in having yellow above eyes and more yellow breasts, and compared to the other species, female O'ahu 'amakihi have two dull wing bars. The O'ahu 'amakihi is brighter and smaller than the Kaua'i 'amakihi but nearly identical to the Hawai'i species. O'ahu 'amakihi are generalized foragers that take arthropods from a variety of trees and substrates. The species often gleans arthropods from leaves and twigs, less frequently from larger branches and trunks. Feeds on nectar and fruit from a variety of native and non-native plants and has been observed eating sap from koa (*Acacia koa*) trees. Only three nests have been found; thus, the species' reproductive biology is poorly known, but is likely similar to Hawai'i 'amakihi.

DISTRIBUTION: Occurs in two disjunct populations between 50 and 300 meters (180 – 1,000 feet), although is most numerous above 200 meters (650 feet). In recent years, range has increased to include some residential areas. Original range likely included all forested regions of O'ahu. Especially in areas where homeowners are planting native trees.

ABUNDANCE: A 1991 survey conducted by the State of Hawai‘i Division of Forestry and Wildlife estimated the O‘ahu ‘amakihi population at between 20,000 and 60,000 individuals. Audubon Christmas bird counts from 1958 to 1985 documented a decrease in detections. Despite this, populations may be increasing in some areas. The population was estimated at 49,500 birds in the Ko‘olau range, and 2,300 in Wai‘anae in 2009.



LOCATION AND CONDITION OF KEY HABITAT:

Occurs in a variety of habitats from very wet forests in the Ko‘olau Mountains to dry forests in the Wai‘anae Mountains. They are more common in sheltered forests in valleys at middle elevations. Unlike other Hawaiian passerines, the range of the O‘ahu ‘amakihi extends to low-elevation forest dominated by non-native plant species. Among introduced forests, ‘amakihi are most abundant in areas dominated by guava (*Psidium guajava*) or kukui (*Aleurites moluccana*). Most of the species’ range is managed by the U.S. Fish and Wildlife Service, U.S. Army, and the State of Hawai‘i.

THREATS: Although O‘ahu ‘amakihi populations appear stable, they are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For O‘ahu ‘amakihi, the following is of particular concern:

- **Fire.** Non-native plants and military training activities often result in wildfires that threaten O‘ahu ‘amakihi habitat on military lands.

CONSERVATION ACTIONS: O‘ahu ‘amakihi likely have benefited from management activities to conserve other endangered forest birds including the establishment of the O‘ahu Forest National Wildlife Refuge in the Ko‘olau mountains, rat control directed at protecting nesting O‘ahu ‘elepaio (*Chasiempis sandwichensis ibidis*), fencing and ungulate control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the O‘ahu ‘amakihi may include the following:

- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Initiate regular forest bird surveys on O‘ahu and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to

control mosquito populations. Because the O‘ahu ‘amakihi appears to be surviving and possibly thriving in disturbed habitat and at elevations below the point where mosquitoes commonly occur, they provide potentially important research opportunities. Research priorities specific to the O‘ahu ‘amakihi include the following:

- Quantify the population structure, dispersal patterns, survivorship, nesting phenology, and success. Studies comparing life history characteristics between native and non-native habitats would be particularly useful.
- Identify and study disease-resistant populations, focusing on the genetic basis of resistance.
- Explore the impact of native plant reforestation and urban forestry on ‘amakihi populations.

References:

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Lindsey GD, VanderWerf EA, Baker H, Baker PE. 1998. Hawai‘i (*Hemignathus virens*), Kaua‘i (*Hemignathus kauaiensis*), O‘ahu (*Hemignathus chloris*) and greater ‘amakihi (*Hemignathus sagittirostris*). In *The Birds of North America*, No. 360 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Photo: Bryan Shiota

Forest Birds

O'ahu 'elepaio

Chasiempis ibidis

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic


NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Endangered

Revised Recovery Plan for Hawaiian Forest Birds

—USFWS 2006

SPECIES INFORMATION: The O'ahu 'elepaio is a small, adaptable monarch flycatcher (Family: Monarchiade) endemic to the island of O'ahu. Other species of 'Elepaio occur on Kaua'i (*C. sclateri*) and the island of Hawai'i (*C. sandwichensis*). Males and females are dark brown above and white below with variable light brown streaks on breast and conspicuous white wing bars, tail feather tips, and throat. Both sexes have variable amounts of blackening markings, but males tend to have more. The bird's name is derived from its primary song which is a shrill whistle given only by males. On the island of Hawai'i, 'elepaio use virtually all available substrates for foraging including the ground, logs, rock crevices, snags, and all parts of tress. Equally diverse in the use of foraging maneuvers, 'elepaio capture a wide range of arthropod prey by flycatching, gleaning while either perched or hovering, and direct pursuit; foraging maneuvers vary depending on plant species from which prey is being captured, and habitat. O'ahu 'elepaio use a variety of native and non-native trees for foraging. Pairs remain together throughout the year, and long-term pair bonds are common. Breeding season on O'ahu is January through July compared to March through August on the island of Hawai'i. Unlike Hawaiian honeycreepers, both males and females participate almost equally in all aspects of rearing. Finely woven cup nests are built in a variety of native and non-native trees. Clutch size is usually two and second and third nests are attempted after failures, but rarely is a second nest attempted if the first is successful. Fecundity is low even in areas where predators are controlled. Young are fed by parents for at least a month, but remain on their natal territory for up to ten months which may allow young birds to hone their foraging skills.



DISTRIBUTION: Occurs in the Ko‘olau Range between 100 to 550 meters (325 – 1,800 feet) elevation, and in the Wai‘anae Range between 500 to 850 meters (1,625 – 2,775 feet) elevation. Dispersal between the ranges is unlikely. Each subpopulation consists of several populations; the amount of dispersal among these is likely low. Original distribution likely included all forested areas of O‘ahu.

ABUNDANCE: In 2013, the population was estimated at 1,261 (95% confidence interval = 1,205-1,317) birds. It had previously been estimated at 1,200 to 1,400 birds. Although Audubon Christmas bird counts from the 1960s through the 1980s provided strong evidence of a dramatic population decline, numbers are now so low that the rate of decline since the 1990s cannot be determined.

LOCATION AND CONDITION OF KEY HABITAT: Occurs in a variety of forest types and across a range of elevations, primarily in valleys and particularly those with tall riparian vegetation, a continuous canopy, and dense understory. Common native plant species where ‘elepaio occur include papala kēpau (*Pisonia umbellifera*), lama (*Diospyros sandwicensis*), māmakī (*Pipturus albidus*), kaulu (*Sapindus oahuensis*) and ‘āla‘a (*Pouteria sandwicensis*). Common introduced plants in ‘elepaio habitat include strawberry guava (*Psidium cattleianum*), common guava (*P. guajava*), kukui (*Aleurites moluccana*), mango (*Mangifera indica*), and Christmas berry (*Schinus terebinthifolius*). O‘ahu ‘elepaio are not found in very wet forests, on windswept summits, or in very dry scrubland. Much of their current range is managed by the U.S. military or by the State of Hawai‘i.

THREATS: O‘ahu ‘elepaio are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For O‘ahu ‘elepaio, the following threats are of particular concern:

- Predation. Predation by black rats (*Rattus rattus*) have been implicated in the loss of nests and death of adult females. Rat control in O‘ahu populations resulted in large increases in nest success and in survival of adult females.
- Low reproductive potential. The species’ low annual productivity, even in quality habitat, makes it very susceptible to factors that reduce population size.
- Disease. Avian pox (*Poxvirus avium*) reduces both annual survival and reproductive success of birds with active pox lesions compared to healthy birds; no information on the effect of avian malaria (*Plasmodium relictum*).
- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios and stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems.
- Fire. Wildfires resulting from military activities threaten two populations.

CONSERVATION ACTIONS: Conservation efforts already undertaken to protect the O‘ahu ‘elepaio include the following: listing as an endangered species by both the U.S. Fish and

Wildlife Service (USFWS) and the State of Hawai‘i, the initiation of long term population and demographic surveys which have identified the most serious threats to its survival, and ongoing rat control at the Honolulu Forest Reserve (DOFAW), at Schofield Barracks West Range and Mākua Military Reservation (U.S. Army Environmental Division), in Honouliuli Preserve (DOFAW) and in Lualualei Valley (U.S. Navy and USDA). In addition, the O‘ahu ‘elepaio also benefits from management activities designed to conserve other endangered forest birds including the establishment of the O‘ahu Forest National Wildlife Refuge in the Ko‘olau Mountains, fencing and ungulate control, forest restoration, habitat monitoring and studies on disease and disease vectors. In addition to these efforts, future management specific to the O‘ahu ‘elepaio should include the following:

- Continue and expand rat control.
- Protect remaining forests on O‘ahu, including through fire prevention.
- Conduct public outreach about the importance and benefits of rodent control.
- Continue demographic studies, especially in the largest populations.
- Continue protection and management of wildlife sanctuaries and refuges.
- Explore translocations to suitable habitat in Maui Nui.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the O‘ahu ‘elepaio include the following:

- Identify disease resistance and transmission patterns. If resistant individuals are identified, translocation and/or captive propagation of these individuals may help recover populations.
- Determine genetic population structure.
- Identify areas most suitable for re-introduction of populations or for creation of habitat dispersal links between existing populations.
- Continue efforts to develop techniques for captive propagation using surrogate species (e.g., Hawai‘i ‘elepaio (*Chasiempis sandwichensis sandwichensis*)).

References:

- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- VanderWerf EA. 1998. ‘Elepaio (*Chasiempis sandwichensis*). In The Birds of North America, No. 344 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- VanderWerf EA, Rohrer JL, Smith DG, Burt MD. 2001. Current distribution and abundance of the O‘ahu ‘elepaio. Wilson Bulletin 113:10-16.
- VanderWerf EA, Lohr MT, Titmus AJ, Taylor PE, Burt MD. 2013. Current distribution and abundance of the O‘ahu Elepaio (*Chasiempis ibidis*). Wilson Journal of Ornithology 125:600-608.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Photo: Top – Moloka'i, bottom – Lāna'i. Bret Mossman courtesy of the Bernice Pauahi Bishop Museum.

Forest Birds

'Oloma'o

Myadestes lanaiensis

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank GH

—Known Only from Historical Occurrences

IUCN Red List Ranking—Critically Endangered


(Possibly Extinct)

Revised Recovery Plans for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The 'oloma'o is one of five species of Hawaiian solitaires (family: Turdidae). All adult Hawaiian solitaires have olive-brown and gray plumage. 'Oloma'o are prolific singers often singing into the night, and the species engages in song flights. Their song, like that of many thrushes, is melodious. Similar to other Hawaiian solitaires, the species often trembles their wings when perched. 'Oloma'o are very philopatric, seldom leaving their small home range. The species feeds on a variety of small fruits and insects. Little is known about their breeding biology, but it is presumed to be similar to the 'ōma'o (*M. obscurus*).

DISTRIBUTION: Unknown. Probably extinct. The historic range included the native forests of eastern Moloka'i and Lāna'i. Historically the species also may have occurred on Maui, where subfossils of Hawaiian solitaires are abundant. Recent sightings (2023-2024) have been reported from Pelekunu and other remote valleys but not substantiated.

ABUNDANCE: Probably extinct. Since 1907, 'oloma'o have been observed on a dozen occasions, most recently in 1988. The Hawaiian Forest Bird Surveys (1979-1980) estimated the population at 19 ± 38 (SE) individuals. However, 'oloma'o were not detected during surveys in the late 1980s and 1990s or more recently. There is little information on historical abundance.



LOCATION AND CONDITION OF KEY HABITAT: Unknown. Historically occupied closed, wet and mesic ‘ōhi‘a (*Metrosideros polymorpha*) forests across a broad elevation range. The areas where the species was last observed are managed by the State of Hawai‘i as a Natural Area Reserve or by private conservation entities (e.g., The Nature Conservancy).

THREATS: Unknown. However, ‘oloma‘o likely were susceptible to the same factors that threaten other native Hawaiian forest birds including loss and degradation of habitat, predation by introduced mammals, and disease. For ‘oloma‘o, the following was likely of particular concern:

- Disease. The fact that no habitat above 1,250 meters (4,100 feet) occurs on Moloka‘i or Lāna‘i suggests disease may have played an important role in the species decline.
- Predation. Following the introduction of non-native mammals, particularly black rats, to Moloka‘i in the late 1800s and Lāna‘i in the early 1900s this species and most other endemics precipitously declined. Similar patterns of extinction and population decline were observed on O‘ahu, Maui, Hawai‘i, and Kaua‘i. Within about a decade of the introduction of black rats to each respective island there were catastrophic population declines and multiple extinctions.

CONSERVATION ACTIONS: If the species persists, it likely benefits from management activities to conserve other endangered forest birds on eastern Moloka‘i, including the establishment and management of protected areas, regular surveys of forest bird populations, monitoring of habitat conditions, and studies of disease and disease vectors. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Revised Recovery Plan for Hawaiian Forest Birds* would be implemented, management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species is probably extinct there are no research priorities specific to oloma‘o.

References:

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.

Wakelee KM, Fancy SG. 1999. 'Oma'o (*Myadestes obscurus*), kama'o (*Myadestes myadestinus*), oloma'o (*Myadestes lanaiensis*), and 'amaui (*Myadestes woahensis*). In *The Birds of North America*, No. 460 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Forest Birds

‘Ōma‘o

Myadestes obscurus


SPECIES STATUS:

State Recognized as Endemic
NatureServe Heritage Ranking G3—
Vulnerable
IUCN Red List Ranking—Vulnerable

SPECIES INFORMATION: One of five species of Hawaiian solitaires (family: Turdidae), the ‘ōma‘o is endemic to the island of Hawai‘i. They often perch silently for long periods and are usually detected by their song; however males perform a flight-song display known as “skylarking.” Like all adult Hawaiian solitaires, ‘ōma‘o have drab olive-brown and gray plumage. Diet consists primarily of fruits of native and introduced understory plant species, although they also take koa (*Acacia koa*) flowers from the canopy and prey on invertebrates, including earthworms, snails, spiders, and insects. The life history is well-studied. Both sexes defend small nesting territories. Nests are built by females in a variety of locations (e.g., cavities, trunk forks); females also perform most incubation and brooding. Clutch size is one or two eggs, and double brooding occurs. The young remain in natal territories for four to six months after fledging. A male-biased sex-ratio exists, but its significance to populations is unknown.

DISTRIBUTION: Primarily occurs in two populations on the eastern and southern slopes of the island of Hawai‘i at elevations greater than 1,000 meters (3,300 feet). A third, smaller population occurs in alpine scrub between 2,000 and 3,500 meters (6,500 – 11,500 feet). Currently occupies about 30 percent of their former range, which historically included habitats between 300 and 3,000 meters (1,000 – 9,750 feet).

ABUNDANCE: The Hawaiian Forest Bird Surveys (1976-79, 1983) estimated the population at 170,000 birds. Based on more recent surveys, populations appear stable and may be increasing below 1,200 meters (3,450 feet).



LOCATION AND CONDITION OF KEY HABITAT: Mesic and wet montane ‘ōhi‘a (*Metrosideros polymorpha*) or mixed ‘ōhi‘a and koa forests in the Hāmākua, Ka‘ū, and Kīlauea districts. These forests support important food plants, including ‘ōlapa (*Cheirodendron trigynum*), kōlea (*Myrsine lessertiana*), kāwa‘u (*Ilex anomala*), naio (*Myoporum sandwicense*), pilo (*Coprosma* spp.), pūkiawe (*Styphelia tameiameia*), ‘ōhelo (*Vaccinium* spp.), and ‘ākala (*Rubus hawaiiensis*). In the small alpine scrub population on Mauna Loa, pūkiawe, ‘ōhelo, kūkaenēnē (*Coprosma ernodeoides*), and ‘a‘ali‘i (*Dodonea viscosa*) are important food plants. Although most of the current range occurs on State and federal lands, habitat conditions vary considerably.

THREATS: ‘Ōma‘o are likely susceptible to the same factors that threaten other native Hawaiian forest birds, including loss and degradation of habitat, predation by introduced mammals, and disease. For ‘ōma‘o, the following are of particular concern:

- Disease. The prevalence of disease in areas tested is low and five ‘ōma‘o exposed to malaria recovered quickly, suggesting a greater disease resistance compared to other native forest birds. However, the species’ disappearance from lower elevations is the pattern of decline noted in other Hawaiian birds susceptible to mosquito-borne diseases.
- Predation. Nests are very accessible and vulnerable to predation by rats (*Rattus* spp.). Predation by native raptors also is likely.
- Habitat degradation. ‘Ōma‘o occur at lower densities in degraded habitat. Pigs (*Sus scrofa*) and other ungulates likely destroy important food plants.

CONSERVATION ACTIONS: ‘Ōma‘o likely have benefited from management efforts designed to conserve other endangered forest birds and native habitat at Hakalau Forest National Wildlife Refuge, Hawai‘i Volcanoes National Park, and the ‘Ōla‘a/Kīlauea Watershed Partnership. These efforts include fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future actions specific to the protection of ‘ōma‘o may include the following:

- Protect and restore native forests above 1,500 meters (4,500 feet), and eliminate feral ungulates and non-native plants.
- Conduct control or eradication of rats and feral cats in areas occupied by ‘ōma‘o.
- Conduct public education and outreach.
- Continue protection and management of wildlife sanctuaries and refuges.
- Restore ‘ōma‘o to former parts of their range.
- Translocate birds to other islands such as O‘ahu and Maui Nui to fill the niche left vacant by those island’s extinct thrush species.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to ‘ōma‘o include the following:

- Identify disease-resistant individuals.

- Develop improved techniques to control alien weed species.

References:

IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

Wakelee KM, Fancy SG. 1999. ‘Oma’o (*Myadestes obscurus*), kama’o (*Myadestes myadestinus*), oloma’o (*Myadestes lanaiensis*), and ‘amaui (*Myadestes woahensis*). In The Birds of North America, No. 460 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Photo: Mike Scott

Forest Birds

‘Ō‘ū

Psittirostra psittacea

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Critically Endangered

(Possibly Extinct)

Revised Recovery Plan for Hawaiian Forest Birds—


USFWS 2006

SPECIES INFORMATION: The ‘ō‘ū is a heavy-bodied Hawaiian honeycreeper (family: Fringillidae) with a distinctive thick, pink, parrot-like bill. Adults are olive-green with whitish undertail coverts; males have a bright yellow head. Early naturalist noted that ‘ō‘ū had a strong musky odor, which is retained in museum specimens. Like several of Hawaii's nectivorous birds, ‘ō‘ū are strong fliers and ranged widely in search of fruit. ‘Ie‘ie (*Freycinetia arborea*) inflorescences apparently are an important part of the species’ diet, although ‘ō‘ū also feed on the fruits of *Clermontia* spp. as well as other native fruits. Geometrid caterpillars are an important food item during the breeding season. Little is known of the species life history and its nesting and breeding habits have not been described.

DISTRIBUTION:

Possibly extinct. Occupied forests between 900 and 1,500 meter (3,000 and 5,000 feet, respectively) elevations on the islands of Kaua‘i and Hawai‘i. Historically widespread, ‘ō‘ū formerly occurred on all the Main Hawaiian Islands in low- to high-elevation forests. They are now presumed extirpated on every island except possibly on Kaua‘i and Hawai‘i.

ABUNDANCE: Almost certainly extinct. The Hawaiian Forest Bird Surveys (1976-1981) estimated the population at 400 ± 300 (95% confidence interval) birds on the island of Hawai‘i and nine or fewer individuals on Kaua‘i. ‘Ō‘ū



have not been detected during more recent surveys, although unconfirmed sightings are occasionally reported.

LOCATION AND CONDITION OF KEY HABITAT: Although known from a range of forest types, all recent observations have occurred in mid-elevation mesic to wet ‘ōhi‘a (*Metrosideros polymorpha*) forests with an understory of ‘ie‘ie (*Freycinetia arborea*), tree ferns (*Cibotium* spp.), ‘ōlapa (*Cheirodendron* spp.), kāwa‘u (*Ilex anomala*), kolea (*Myrsine* spp.), and pilo (*Coprosma* spp.). All recent sighting of ‘ō‘ū have occurred on lands managed by the State of Hawai‘i.

THREATS:

- Habitat degradation. Pigs (*Sus scrofa*) degrade the understory of wet forest destroying food plants.
- Disease. ‘Ō‘ū primarily occurred in low- to mid-elevation forests where the effects of mosquito-borne diseases was most severe. The species’ foraging movements may have increased their exposure to disease.
- Predation. In addition to potentially depredating nests, rats (*Rattus* spp.) may also compete with ‘ō‘ū by reducing the availability of fruits.
- Natural disasters. In 1984, a large portion of the Upper Waiākea Forest Reserve was inundated by a lava flow from Mauna Loa. This flow occurred in an area where the most recent observations of the species were noted and destroyed high quality ‘ō‘ū habitat. In 1982 and 1992, two strong hurricanes struck Kaua‘i, devastating native forest habitat. The ‘ō‘ū, has not been observed on Kaua‘i since 1992.

CONSERVATION ACTIONS: If the species persists, it likely benefits from management efforts to conserve other endangered forest birds on Hawaii and Kaua‘i. On Hawai‘i, these activities have included fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. On Kaua‘i, these activities included the establishment of the Alaka‘i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, studies of disease and disease vectors, and public education efforts featuring Kauai’s endangered forest birds. Should this species be rediscovered, the Rare Bird Recovery Protocol outlined in the U.S. Fish and Wildlife Service (USFWS) *Recovery Plan for Hawaiian Forest Birds* would be implemented, and management in anticipation of that possibility should include continued protection and management of wildlife sanctuaries and refuges.

MONITORING: Continue forest bird surveys and habitat monitoring on Hawai‘i and Kaua‘i.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include developing improved methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Given that this species may be extinct, there are no research priorities specific to ‘ō‘ū.

References:

- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Snetsinger TJ, Reynolds MH, Herrmann CM. 1998 'O'u (*Psittirostra psittacea*) and Lana'i hookbill (*Dysmorodrepanis munroi*). In The Birds of North America, No. 335-336 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Forest Birds

Palila

Loxioides bailleui

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic

NatureServe Heritage Rank G1—Critically Imperiled


IUCN Red List Ranking—Critically Endangered

Revised Recovery Plan for Hawaiian Forest Birds

—USFWS 2006

Critical Habitat Designated 1977

SPECIES INFORMATION: The palila is a finch-billed Hawaiian honeycreeper (Family: Fringillidae) whose life history and survival is linked to māmane (*Sophora chrysophylla*), an endemic dry-forest tree in the legume family. Males and females are similar, with a yellow head and breast, greenish wings and tail, a gray back, and white underparts. Males have a black mask, and females have less yellow on the back of their heads and a gray mask. Approximately 90 percent of the palila's diet consists of immature māmane seeds; the remainder consists of māmane flowers, buds, leaves, and naio (*Myoporum sandwicense*) berries. Caterpillars and other insects comprise the diet of nestlings, but also are eaten by adults. Māmane seeds have been found to contain high levels of toxic alkaloids, and palila use particular trees for foraging, suggesting that levels of alkaloids may vary among trees. Individuals will move limited distances in response to the availability of māmane seeds. Palila form long-term pair bonds, and males perform low advertisement flights, sing, chase females, and engage in courtship feeding prior to breeding. Females build nests, usually in māmane trees, and males defend a small territory around the nest tree. Females mostly incubate eggs, brood nestlings and feed young with food delivered by male. First-year males sometimes help a pair by defending the nest and feeding the female and nestlings. Limited genetic testing found no evidence that helpers father the nestlings they were assisting, although more data are needed. Fledglings are dependent on their parents for three to four months, during this time they learn and practice foraging skills. The availability of green māmane seeds



strongly influences the number of nesting attempts in a given year. In poor years, not all pairs will attempt to nest.

DISTRIBUTION: Mostly restricted to the western slopes of Mauna Kea between 2,000 and 2,850 meters (6,500 – 9,250 feet) elevation. Historically, palila were common in all māmane forests. Currently, the species occupies approximately 10 percent of their historical range on the island of Hawai‘i. Subfossil evidence indicates palila also occurred in māmane forest on O‘ahu and Kaua‘i.

ABUNDANCE: Annual surveys between 1998 and 2005 yielded a mean population estimate of $3,268 \pm 190$ (SE) birds. In 2014, the estimate was $2,070 \pm 209$ birds. Populations have continued to decline and in 2024 was estimated to number 666 individuals. Population estimates are variable among years, which may be an artifact of survey techniques or timing.

LOCATION AND CONDITION OF KEY HABITAT: Restricted to māmane and māmane/naio forests. Densities are highest at 2,300 meters (7,550 feet) with large māmane trees and a high proportion of native shrubs. Up to 96 percent of the population and nearly all successful breeding occurs in a 30 square kilometer (11.5 square mile) area on the southwestern slope of Mauna Kea that has high quality habitat and steep terrain. The latter is important, especially during the breeding season because māmane at different elevations flower and fruit at different times, ensuring that māmane seeds are always available. Most habitat in the species’ range is severely degraded by grazing ungulates, particularly mouflon sheep, and the spread of non-native plant species, especially fire-prone grasses. Most of the palila’s current range occurs in the Mauna Kea Forest Reserve and is managed by the State of Hawai‘i.

THREATS:

- Feral ungulates. Historically, large numbers of sheep (*Ovis* spp.) grazed on Mauna Kea, reducing the density and productivity of māmane trees, limiting the regeneration of other native plants, and causing soil erosion. More recently the introduction of mouflon sheep has further degraded habitat.
- Invasive plants. Soil disturbance caused by sheep facilitated the spread of invasive plants. Fire-adapted grasses, such as fountain grass (*Pennisetum clandestinum*), are especially problematic in that they increase the risk of fire. Invasive plants also reduce the recruitment of native plants.
- Fire. A single large fire could severely limit food resources for the entire population.
- Predation. Predation by rats (*Rattus* spp.), feral cats (*Felis silvestris*), and the pueo (*Asio flammeus sandwichensis*) is an important factor limiting palila populations. In some areas, black rats (*R. rattus*) and feral cats may be responsible for up to 40 percent of nest failures, and feral cats have been documented preying on adults.
- Disease. Although palila are very susceptible to mosquito-borne diseases, mosquitoes do not occur at the elevations where palila currently occur. However, disease almost certainly contributed to the species’ decline and likely prevents palila from recolonizing low-elevation habitat.

- Non-native insects. Yellow jackets and Argentine ants threaten the native caterpillars that nestlings depend on for food.
- Severe weather. The species' current range exposes them to severe weather that results in mortality in some years. Freezing temperatures, heavy rains, droughts, and high winds all can result in egg and nestling mortality.
- Population size. Small populations are plagued by a variety of potentially irreversible problems that fall into three categories: demographic, stochastic, and genetic; the former are usually most problematic. Demographic factors include skewed sex ratios and stochastic factors include natural disasters. Habitat fragmentation exacerbates demographic and genetic problems.

CONSERVATION ACTIONS: Several conservation efforts have been conducted to protect the māmane woodlands for the protection and recovery of the palila. In the first half of the 20th century, 46,000 sheep as well as smaller numbers of feral cattle (*Bos tarus*), goats (*Capra hircus*), and pigs (*Sus scrofa*) were removed from māmane forests. This allowed the regeneration of the māmane trees on which the palila depend. Beginning in the late 1970s, control measures to reduce mouflon sheep numbers were initiated. Considerable research has been conducted on palila, including estimating population size and determining their geographic range, documenting home range size, dispersal behavior, reproductive parameters, limiting factors, and habitat characteristics. Ongoing research mostly focuses on quantifying limiting factors, especially predation, food availability, and genetics, as well as refining māmane restoration techniques. Beginning in 1993, translocations were conducted on an experimental basis to determine if new breeding populations could be established. In 2004, 32 birds were translocated and in 2005, 75 birds were moved. Although most birds return to their natal territory, approximately 25 percent remained at translocation sites. In 1996, a captive propagation program was initiated and palila have successfully bred in captivity. Between 2003 and 2004, 15 captive-raised birds were released into the wild. A second attempt to establish palila on the north slope was initiated in 2019, but it was unsuccessful. Initial attempts at outplanting māmane have been successful in areas where competing non-native vegetation is sparse. In addition to the above efforts, palila likely have benefited from management activities to conserve other endangered forest birds in Mauna Kea Forest Reserve and elsewhere on the island of Hawai'i, including fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies of disease and disease vectors. In addition to these efforts, future management specific to the palila will likely include the following:

- Stabilize and increase at least one of the small populations using translocation or establish a new self-sustaining population while continuing to intensely manage the primary population. These efforts must include the restoration and regeneration of māmane forest as well as control of mammalian predators and parasitoid wasps that threaten food sources.
- Evaluate sites throughout the species' historical range for potential māmane restoration and re-introduction. These sites should have a range of elevation or rainfall gradients to ensure year-round food availability.
- Establish cat or predator proof fencing around key palila breeding areas and maintain and expand small mammal trapping.
- Develop a comprehensive fire management plan.

- Increase public education and involvement in palila recovery. Volunteer opportunities exist in habitat restoration efforts, monitoring weeds and predators, and education.
- Remove feral sheep and mouflon from palila habitat.
- Continue protection and management of wildlife sanctuaries and refuges.
- Establish a rear and release program for young palila.
- Develop supplemental feeding for wild palila.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the palila include the following:

- Refine survey methods.
- Further refine techniques to facilitate the establishment of new populations.
- Develop methods to control and eradicate the most harmful non-native plants and non-native insects that threaten native insect food sources.

References:

- Banko PC, Johnson L, Lindsey GD, Fancy SG, Pratt TK, Jacobi JD, Banko WE. 2002. Palila (*Loxioides bailleui*). In The Birds of North America, No. 679 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Camp RJ, Brinck KW, Banko PC. 2014. Palila abundance estimates and trend. HCSU Technical Report HCSU-053. 18 pp.
- Hunt, N. J., C. K. Asing, L. Nietmann, P. C. Banko, and R. J. Camp. 2025. 2022–2024 Status and trend of the palila (*Loxioides bailleui*). Hawai‘i Cooperative Studies Unit Technical Report HCSU-115. University of Hawai‘i at Hilo. 22 pages.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service.



Photo: DOFAW

Forest Birds

Po'ouli

Melamprosops phaeosoma

SPECIES STATUS:

Federally Listed as Extinct

State Listed as Extinct

State Recognized as Endemic

NatureServe Heritage Rank: G1 – Presumed Extinct


IUCN Red List Ranking: Extinct

Revised Recovery Plan for Hawaiian Forest Birds - USFWS 2006

SPECIES INFORMATION: The po'ouli is a stocky Hawaiian honeycreeper (Family: Fringillidae) endemic to Maui that was not discovered until 1973. Po'ouli have short wings and tail, a finch-like bill, and distinctive plumage. Aptly named “black-faced” in Hawaiian, po'ouli have a large black face mask, white cheeks, throat, and underparts and brown wings and back; no other Hawaiian forest bird is similarly colored. Although not well-studied, males and females are similar, although females have a grayish throat and breast. Little is known of the species' life history. Po'ouli often join mixed species foraging flocks. Forages primarily in the subcanopy and understory on tree branches of native shrubs and trees where they search moss, lichens, and bark for snails and arthropods. Breeding biology is based on observations from a single nesting pair and may be biased because of extremely low population density. For example, territorial behavior has not been observed. Nests are similar to those of other Hawaiian honeycreepers and are placed in 'ōhi'a (*Metrosideros polymorpha*) trees. Only the female incubates eggs and broods nestlings; male feeds the female on and off the nest. Last seen in the wild in 2004. Declared extinct by U.S. Fish & Wildlife Service in 2023.

DISTRIBUTION: In 2004, it was restricted to a 1,300 hectare (3,200 acre) area between 1,440 and 2,100 meters (4,750 – 7,000 feet) elevation on the northeastern slope of Haleakalā on Maui. No historical data on range, although fossil evidence indicates that po‘ouli once occurred over a wider geographic range.

ABUNDANCE: The Hawai‘i Forest Bird Survey in 1980 estimated the population at 140 ± 280 (95% confidence interval) individuals. As of 2003 there were three known individuals. One captured and brought into captivity in 2004 died in November of that year and the remaining two birds have not been seen since January 2004. The species’ poor detectability has hampered attempts to locate additional individuals or estimate population size. The species has been declared extinct.



LOCATION AND CONDITION OF KEY HABITAT: Mixed shrub montane wet forest dominated by ‘ōhi‘a, with an understory dominated by a diversity of small trees and shrubs, many of which are used as foraging substrates (e.g., kanawao [*Broussaissea arguta*], kawa‘u [*Ilex anomala*]). Habitat conditions in areas occupied by po‘ouli are variable, but improving. All known individuals occur (red) in the Hanawā Natural Area Reserve which is managed by the State of Hawai‘i.

THREATS: Unknown. However, po‘ouli is likely susceptible to the same factors that threaten other native Hawaiian forest birds, including habitat loss and degradation, predation by introduced mammals, and disease. For po‘ouli, the following are likely of particular concern:

- Habitat degradation. The species appears to prefer areas with low levels of soil and vegetation disturbance and therefore may be particularly sensitive to understory and ground cover damage by feral pigs (*Sus scrofa*).
- Predation. Rats (*Rattus* spp.) are abundant in po‘ouli habitat and may depredate adults and nests.
- Competition. In addition to direct predation on adults and nests, rats also may compete with po‘ouli for food resources, especially snails. Also, the non-native garlic snail (*Oxychilus alliarius*) is abundant and preys on native snails.
- Disease: the last known individual that was brought in captivity and later died had chronic avian malaria. It is likely that disease spread from non-native mosquitoes contributed to their extinction.

CONSERVATION ACTIONS: Major efforts to conserve po‘ouli include the establishment of the 3,000 hectares (7,500 acres) Hanawā Natural Area Reserve in 1986 to protect the species’

entire known range, and fencing and pig removal has facilitated recovery of the understory. In 1995, the Maui Forest Bird Recovery Project was created to research and protect native Hawaiian forest birds, including the po‘ouli. Activities to date include small mammal control, research on optimizing rodent control methods, surveys for native land snails, and banding and collecting blood samples to monitor demography and disease prevalence. In 2002, the team successfully translocated a female po‘ouli to the home range of the male, however, she quickly returned to her own home range. In September 2004, one po‘ouli was captured in an attempt to establish a captive population but it died in November of the same year. The Hawai‘i Department of Land and Natural Resources and the U.S. Geological Survey continue to search Hanawī and adjacent habitats for additional birds. In addition, po‘ouli populations likely have benefited from management efforts to conserve other endangered forest birds on northeastern Haleakalā including fencing, ungulate and small mammal control, forest restoration, habitat monitoring, and studies on disease and disease vectors. In addition to these efforts, future management specific to the po‘ouli may include the following:

- Continue pig and small mammal control efforts.
- Continue efforts to locate and capture the two known individuals and additional birds.
- Conduct public outreach about the importance of pig and rodent control to the remaining forest bird populations on Maui.
- Continue protection and management of wildlife sanctuaries and refuges.
- Implement mosquito control techniques to reduce non-native disease in forest bird populations.

MONITORING: Continue forest bird surveys and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining the ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. In addition to the ongoing research on po‘ouli outlined above, additional priorities specific to po‘ouli include the following:

- Determine the distribution and abundance of the species’ prey base to determine if food resource is a limiting factor.

References:

- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Pratt TK, Kepler CB, Casey TLC. 1997. Po‘ouli (*Melamprosops phaesoma*). In *The Birds of North America*, No. 272 (Poole A, Gill F, editors). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service. 428 pp.



Forest Birds

Puaiohi

Myadestes palmeri

SPECIES STATUS:

Federally Listed as Endangered

State Listed as Endangered

State Recognized as Endemic


NatureServe Heritage Rank G1—Critically Imperiled

IUCN Red List Ranking—Critically Endangered


Revised Recovery Plan for Hawaiian Forest Birds—USFWS 2006

SPECIES INFORMATION: The puaiohi, or small Kauaʻi thrush, is the smaller of two solitaires (family: Turdidae) endemic to Kauaʻi, and was the last of the island’s avifauna to be discovered by western ornithologists. In the late 1800s, the puaiohi was considered exceedingly rare, but this assessment was likely due to its cryptic behavior and preference for remote, inaccessible ravines. Adult puaiohi are extremely sedentary and appear to have specific habitat requirements. Of the five Hawaiian solitaires, it is behaviorally, morphologically, and vocally the most divergent. Compared to the kāmāʻo, or large Kauaʻi thrush (*M. myadestinus*), the puaiohi has a short tail, and a white eye-ring; in addition, it has a relatively simple song. The species life history is relatively well-known because of recent, intensive studies. Fruit dominates the non-breeding season diet; insects are important during the breeding season. Nests are built by the female in tree cavities or on cliff ledges, and only females incubate eggs and brood young. Breeding peaks from April to June, and re-nesting occurs after failed and successful nest attempts. This behavior, plus a high rate of nest success and a long breeding season, can result in high annual productivity. However, female and juvenile survival is low (0.46 ± 0.12 and 0.23 ± 0.06 , respectively). Hatch-year and second-year birds (i.e., helpers) are known to assist in nest defense and feeding of related nestlings and fledglings at some sites. Young are very sedentary for two to four days after fledging.

DISTRIBUTION: Restricted to a < 20 square kilometer (7.6 square mile) area on the southern and central plateau of the Alakaʻi Wilderness Preserve. Currently occurs above 1,050 meters (3,450 feet), which is similar to its upper limit historically. It is more likely to occur in streams between 1,140 and 1,340 meters



with large numbers of tall, steep cliffs than in less incised streams. Prehistorically occurred at sea level.



All known streams above 1,050 meters on the Alakai Plateau, Kaua‘i, which represents the best estimate of the entire puaiohi range.

ABUNDANCE: The most recent surveys in 2011-2013 and subsequent modeling estimated the population at 494 (95% confidence interval: 414-580, Crampton et al. 2017) individuals. Occupancy during these surveys ranged from 0.12 to 0.81 on 12 streams across the range, and has been relatively consistent since 2012 on four streams that were resurveyed in 2024 (KFBRP, unpubl. data). Densities peaked at 16 breeding pairs per square kilometer (0.62 square miles), but varied across streams. This species has been considered rare since the late 1800s.

LOCATION AND CONDITION OF KEY HABITAT: Puaiohi seem somewhat tolerant of avian malaria, so it is not clear what is restricting them to higher elevations—perhaps loss and degradation of habitat (see below). Occurs in deeply dissected, steep-walled ravines supporting wet montane forest dominated by ‘ōhi‘a (*Metrosideros polymorpha*) and ‘ōlapa (*Cheirodendron* spp.). Their former range included mesic areas dominated by ‘ōhi‘a and koa (*Acacia koa*). These areas are now largely dominated by introduced plant species including fire tree (*Myrica faya*)

and strawberry guava (*Psidium cattleianum*). Densities are very low even in apparently suitable habitat. The entire known range of this species is within the Alaka'i Wilderness Preserve and Na Pali Kona Forest Reserve, which are and is managed by the State of Hawai'i. However, models predict that low numbers of puaiohi occur on private lands adjacent to these areas, which is supported by opportunistic observations when traveling near property boundaries.

THREATS:

- Disease. In 1995-1997, seven puaiohi were tested for disease; none carried malarial (*Plasmodium relictum*) antibodies. Since 2011, 35/181 puaiohi have tested positive for malarial antibodies (KFBRP, unpubl. data).. Survival of puaiohi that tested positive was similar to that of puaiohi that tested negative, indicating a possible resistance to or tolerance of malaria.
- Predation. Two years of field data indicate that rats (*Rattus* spp.) were responsible for 14 to 22 percent of nest failures, as well as mortality of three adult females. Subsequent studies documented at least ten other instances of predation on adults at the nest, and the lower survival of females (0.46 ± 0.12) compared to males (0.71 ± 0.09) is indicative of predation on females at the nest. In 2020, cat (*Felis cattus*) predation of a puaiohi nest in a cliff alcove was documented. Fledglings typically spend their first days out of the nest within 2 meters (6.5 feet) of the ground, which makes them vulnerable to feral cats and rats. Cats have become more common in the Alaka'i in the last 10 years, following the new fence lines.Competition. Several non-native birds, including the introduced Warbling white-eyes (*Zosterops japonicus*), melodious laughing thrush (*Copsychus malabaricus*), and white-rumped shama (*Copsychus malabaricus*), occupy the same habitat and may compete with the puaiohi for food and nest resources. Shama have been observed nesting in nest boxes erected to promote puaiohi nesting on at least two occasions. Preliminary dissection of fecal samples indicate that Japanese white-eyes eat primarily small non-native fruits, whereas puaiohi eat native fruits almost exclusively. Fecal samples of shama and laughing thrush have not been obtained.
- Habitat degradation. Feral pigs (*Sus scrofa*) and goats (*Capra hircus*) have facilitated the invasion of non-native plants into puaiohi habitat. The establishment of these plants has altered the structure of these forests, especially the ground and shrub layer, and especially in the western part of the puaiohi's range where it is now exceedingly rare. Hurricanes in 1982 and 1992 further altered and degraded the forests of Kaua'i, including those within the Alaka'i Wilderness Preserve.
- Non-native arthropods. Recently introduced non-native insects, especially yellow jackets (*Vespula pensylvanica*) and Argentine ants (*Linepithema humile*), may compete with the puaiohi's native arthropod prey or disrupt the pollination of the species' food plants. Introduced herbivorous insects also could reduce the abundance of food plants.
- Natural disasters. Hurricanes in 1982 and 1992 likely caused the death of an unknown number of individuals.

CONSERVATION ACTIONS: In 1995, an intensive field ecology and behavior study was initiated. The results from this study are the basis for the current management actions for the species. Using captive propagation and re-introduction techniques developed using the 'ōma'ō (*M. obscurus*), a captive breeding flock of puaiohi was established in 1996. Between 1999 and

2012, 222 captive-bred puaiohi were released into the wild on Kaua'i. Initially, the program was successful, with several captive-raised birds documented successfully breeding, but in later releases, survival and recruitment was lower. From 2005 to 2010, captive-bred birds released as hatch years had survival of 0.26 ± 0.21 , and captive-bred birds released as adults had even lower survival (0.05 ± 0.06). Overall, only 20 released birds were observed paired up, nesting, and raising at least 24 chicks to fledging. Given these results, and the apparent stability of the wild population, this program was discontinued, and the remaining captive puaiohi were released at Halepa'akai in 2016 and 2017. PVA results published in 2018 (Fantle-Lepczyk et al.) indicates rat control would be the most effective management strategy to increase puaiohi abundance, primarily by increasing female and juvenile survival.

In addition, puaiohi likely have benefited from management activities to conserve other endangered forest birds including the establishment of the Alaka'i Wilderness Preserve, regular surveys of forest bird populations, monitoring of habitat conditions, control of feral ungulates through public hunting and fencing, suppression of rats on two A24 trapping grids over approximately 150 ha (470 ac), studies of disease and disease vectors, and public education efforts featuring Kauai's endangered forest birds. In addition to these efforts, future management specific to the recovery of puaiohi will likely include the following:

- Systematically control rats using registered rodenticides or traps in puaiohi nesting habitat over a larger area than currently under protection.
- Aggressively control ungulates to improve habitat quality and facilitate recovery of degraded, but potential, habitat.
- Control non-native plants as part of forest restoration efforts.
- Suppress rats, feral cats and barn owls (*Tyto alba*) in the Alaka'i Wilderness Preserve and Kokee Region.
- Reduce breeding habitat for mosquitoes on the Alaka'i Plateau,
- Suppress mosquito populations through releases of incompatible or sterile mosquitoes.
- Investigate options to eradicate mosquitoes, including use of genetic manipulations.
- Prevent the spread of Rapid Ohia Death on the Alaka'i Plateau
- Prevent the introduction of the small Indian mongoose and other possible predators.
- Conduct public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.
- Translocate to other islands with appropriate habitat.

MONITORING: Continue puaiohi specific occupancy surveys, nest monitoring, and habitat monitoring.

RESEARCH PRIORITIES: Research priorities for most Hawaiian forest birds include improving methods for controlling rats and feral cats in native forests, determining ecological requirements of *Culex* mosquitoes at mid- and high-elevation forests, and developing methods to control mosquito populations. Research priorities specific to the puaiohi include the following:

- Investigate immediate impacts of acute disease (malaria, pox) on puaiohi survival and sub-lethal impacts on fecundity
- Conduct captive-breeding of disease-resistant birds to establish disease-resistant populations at lower elevations.

- Continue field studies to document survival and dispersal, habitat use, and causes of high female and juvenile mortality.
- Investigate efficacy of other methods of controlling invasive rodents (e.g., different methods of applying rodenticide, snap traps).
- Assess the impact of introduced rodents on food sources important to puaiohi, including fruits and invertebrates.

References:

- Atkinson, CT, RB Utzurrum, DA LaPointe, RJ Camp, LH Crampton, JT Foster, and TW Giambelluca. 2014. Changing Climate and the Altitudinal Range of Avian Malaria in the Hawaiian Islands - an Ongoing Conservation Crisis on the Island of Kaua'i. *Global Change Biology* 20, 2426–2436, doi: 10.1111/gcb.12535.
- Crampton LH, Brinck KW, Pias KE, Heindl BAP, Savre T, Diegmann JS, Paxton EH. 2017. Linking occupancy surveys with habitat characteristics to estimate abundance and distribution in an endangered cryptic bird. *Biodiversity and Conservation* 26: 1525-1539.
- Fantle-Lepczyk J, Taylor A, Duffy DC, Crampton LH, Conant S. 2018. Using population viability analysis to evaluate management activities for an endangered Hawaiian endemic, the Puaiohi (*Myadestes palmeri*) *Plos one* 13: e0198952.
- Foster JT, Tweed EJ, Camp RJ, Woodworth BL, Adler CD, Telfer T. 2004. Long-term population changes of native and introduced birds in the Alaka'i swamp, Kaua'i. *Conservation Biology* 18:716-725.
- IUCN Red List of Threatened Species. 2015. Version 2014.3. Available at: www.iucnredlist.org. (Accessed May 2015).
- Scott JM, Mountspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Switzer R., Lieberman A, Nelson J, Crampton LH. 2013. Augmentation of the Puaiohi population through captive propagation and release on the Alakai Plateau, Kauai, Hawaii, USA. Reintroduction Specialist Group Book, IUCN.
- U.S. Fish and Wildlife Service. 2006. Revised Recovery plan for Hawaiian forest birds. Portland, (OR): U.S. Fish and Wildlife Service..
- Vanderwerf, EA, LH Crampton, PK Roberts, JS Diegmann, and DL Leonard. 2014. Survival estimates of wild and captive-released Puaiohi, an endangered Hawaiian thrush. *Condor* 116: 609–618. DOI: 10.1650/CONDOR-14-18.1
- Wakelee KM, Fancy SG. 1999. 'Oma'o (*Myadestes obscurus*), kama'o (*Myadestes myadestinus*), oloma'o (*Myadestes lanaiensis*), and 'amaui (*Myadestes woahensis*). In *The Birds of North America*, No. 460 (Poole A, Gill F, editors.). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.



Raptors

ʻIo

Buteo solitarius

SPECIES STATUS:

State Listed as Endangered
State Recognized as Endemic NatureServe Heritage
Rank G2—Imperiled IUCN Red List Ranking—Near
Threatened, Hawaiian Hawk Recovery Plan—USFWS 1984
Bird of Conservation Concern - USFWS 2021

SPECIES INFORMATION: The ʻio, or Hawaiian hawk, is the only broad-winged hawk (Family: Accipitridae) in Hawaiʻi. ʻIo are considered ʻaumākua, or family gods, by Native Hawaiians. Similar to many birds of prey, females are larger than males, and often weigh approximately 25 percent more than males. Also similar to many *Buteos*, two color morphs, light and dark, occur in ʻio populations. Prior to the arrival of Polynesians, ʻio may have exclusively preyed on birds, including now extinct flightless ibis, and rails. Its diet now includes non-native insects, birds and rodents, as well as native insects and birds. ʻIo form monogamous long-term pair-bonds and defend territories year-round. Nest construction is protracted, beginning up to two months before the first egg is laid, and continuing into the nestling period. Egg-laying generally occurs from March to June, and fledging from July to September. Both sexes contribute to nest-building. Clutch size is nearly always one, although historically clutches of two and three were reported. Both sexes incubate, although females perform most of the brooding of nestlings; males provide most of the food to chicks and female. Both adults feed fledglings, which are dependent on adults for up nine months. The ʻio was delisted from the Federal Endangered Species Act in 2019 but remains classified as an endangered species under the State of Hawaii.

DISTRIBUTION: Occurs throughout the island of Hawaiʻi from 300 to 2,600 meters (1,000 to 8,530 feet). Based on fossil evidence, they once occurred on Kauaʻi, Molokaʻi, and Oʻahu.

ABUNDANCE: Based on an island-wide survey in 2007, the population is estimated at 3,000 birds with a stable population trend over the past 20 years.

LOCATION AND CONDITION OF KEY HABITAT: Lowland non-native forests, urban areas, agricultural lands, pasturelands, and high-elevation native forests. Most nesting occurs in native ʻōhiʻa trees, although also occurs in non-native trees, including eucalyptus, ironwood, mango, coconut palm, and macadamia. In winter, they have been reported in subalpine māmane-naio forest, suggesting some seasonal movements.

THREATS:

- Habitat loss and degradation: Habitat is negatively affected by urbanization, land

conversion to unsuitable foraging habitat (e.g., pasture and cane fields to eucalyptus forest), increase in fire frequency that may eliminate nesting and perching habitat, and invasion of understory plants which can conceal prey and reduce foraging success. ‘Io nests occur in a variety of trees, but over 80% of nests observed in a 2003 study occurred on the native ‘ōhi‘a tree (Klavitter et al. 2003). Threats to native forests on Hawai‘i Island are current and rapid ‘ōhi‘a death is changing the landscape level distribution of ‘ōhi‘a trees with mortality rates as high as 90% being found in some areas.

- Volcanic activity: Forested areas have been lost either through direct lava inundation or increased levels of toxic sulphur dioxide fumes.
- Disease: ‘Io does not appear to be susceptible to the avian malaria and avian pox that have devastated other Hawaiian endemic forest birds. However, West Nile virus could affect the species if the disease reaches Hawai‘i.
- Environmental contaminants and toxins: ‘Io are secondarily exposed to both first and second-generation anticoagulant rodenticides and likely face both lethal and sublethal effects from these toxicants. Increased use of urban, agricultural, and other human-altered environments puts ‘io at greater risk of encountering these environmental toxins.
- Shooting: Harassment of nesting birds and shooting was identified in the U.S. Fish and Wildlife Service’s Hawaiian Hawk Recovery Plan (USFWS 1984) as a significant factor affecting the species. In 2018, four cases of ‘Io deaths were attributed to firearms and another two were reported in 2013 on Hawai‘i Island.

CONSERVATION ACTIONS: ‘Io likely benefit from actions for other endangered birds, such as fencing, ungulate and small mammal control, and forest restoration. Actions specific to conservation of the ‘io should include the following:

- Protect and manage the species’ foraging and nesting habitat.
- Minimize use of rodenticides and other environmental toxins within known ‘Io occupancy areas
- Concerted outreach for ‘Io to reduce and prevent harassment and shooting.

MONITORING: Conduct regular island-wide surveys to monitor population dynamics and abundance, distribution, and disease.

RESEARCH PRIORITIES: Research priorities specific to ‘Io include:

- Evaluate potential effects of land management and changes in fire frequency and intensity on the species.
- Continue regular population monitoring as outlined in the 2019 U.S. Fish and Wildlife Post-delisting Monitoring Plan for the Hawaiian Hawk or ‘Io, to assess the impacts of delisting the species from the federal ESA.

References:

Clarkson KE, Laniawe LP. 2000. Hawaiian hawk (*Buteo solitarius*). In *The Birds of North America*, No. 523 (Poole A, Gill F, editors). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American

Ornithologists' Union.

Hawai'i Natural Heritage Program [Hawai'i Biodiversity and Mapping Program]. 2004. Natural diversity database. University of Hawai'i, Center for Conservation Research and Training. Honolulu, HI.

IUCN Red List of Threatened Species. 2015. Version 2014.3. www.iucnredlist.org. (Accessed May 2015).

NatureServe. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (Accessed May 2015).

Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.

Sierts, S.R, S. E Volker, C.G Payne, R. T. Sugihara, C.E. Swift, J.T. Nelson, A.B. Shiels, and W.C. Pitt. 2019. Screening of Hawaiian bird and bat carcasses for rodenticide residues. Unpublished final report (QA-2240). USDA, APHIS, WS, NWRC. Hilo, HI.

U.S. Fish and Wildlife Service. 1984. Hawaiian Hawk Recovery Plan. Portland, Oregon.

U.S. Fish and Wildlife Service. 2008. Endangered and threatened wildlife and plants; withdrawal of proposed reclassification of the Hawaiian hawk or io (*Buteo solitarius*) from endangered to threatened; proposed rule to remove the Hawaiian hawk from the federal list of endangered and threatened wildlife. Federal Register 73:45680–45689.

U.S. Fish and Wildlife Service. 2019. Post-delisting Monitoring Plan for the Hawaiian Hawk, or Io (*Buteo solitarius*). Endangered Species Division, Pacific Island Fish and Wildlife Office, Honolulu, HI.



Raptors

Pueo or Hawaiian Short-eared Owl

Asio flammeus sandwichensis

SPECIES STATUS:

State listed as
Endangered on O‘ahu State recognized
as Endemic at the subspecies level
NatureServe Heritage Rank G5/T2 –
Species secure/Subspecies imperiled
Bird of Conservation Concern
- USFWS 2021

SPECIES INFORMATION: The Pueo, or Hawaiian Short-eared Owl, is an endemic subspecies of the nearly globally distributed Short-eared Owl (*Asio flammeus*; Family: Strigidae). Pueo are ‘aumakua (deified ancestors) that have the ability to restore life or save people from imminent death. The species is thought to have colonized the Hawaiian Islands sometime after the arrival of Polynesians.

Unlike most owls, Pueo are occasionally active during the day, however, they are most active during crepuscular (morning and evening) and nocturnal times (Wilhite et al. 2024). They are commonly seen hovering or soaring over open areas but they also occupy forested and developed habitats (Wilhite et al. 2024). Like Short-eared Owls in continental environments, those in Hawai‘i primarily consume small mammals, but unlike their continental counterparts Pueo will also hunt birds (including native seabirds) and invertebrates. Their relatively recent establishment on Hawai‘i may have been tied to the Pacific rats (*Rattus exulans*) that Polynesians brought to the islands. Little is known about the breeding biology of Pueo, but nests or recently fledged young have been found between November and July. Males perform aerial displays known as a sky dancing display to prospective females.

Nests are constructed by females and are simple scrapes in the ground occasionally lined with grasses and down feathers. Clutch sizes generally range between 2-7 eggs. Females perform all incubating and brooding. Males feed females and both sexes will defend nests. Chicks hatch asynchronously and are fed by the female with food delivered to her by the male. Young fledge from the nest on foot before they are able to fly, beginning at approximately 10 days old, but are flightless and remain on the ground in the vicinity of the nest until approximately 28 days old. After fledging, they depend on their parents for another month.

DISTRIBUTION: Found on all the Main Hawaiian Islands from sea level to 2,450 meters (8,000 feet). Individuals are documented to move between islands, such as from Kaho‘olawe to Maui, or from O‘ahu to Moloka‘i, although the frequency at which inter-island movements occur is still unknown.

ABUNDANCE: Unknown. Because of relatively few detections, the Hawaiian Forest Bird Survey did not estimate the population size of the Pueo. Pueo were widespread at the end of the 19th century and are thought to be declining, but recent analysis of eBird data indicates that Pueo occupancy rates across the Main Hawaiian Islands have remained low but stable from 2010-2020. However, focused surveys for the species will still be necessary to determine population size. However, focused surveys for the species will still be necessary to determine population size.

LOCATION AND CONDITION OF KEY HABITAT: Pueo occupy a variety of habitats, including wet and dry forests, suburban and developed areas, and open habitats such as grasslands, shrublands, wetlands, and montane parklands, including those actively managed for conservation. Pueo are also documented to breed in wet and dry native forests, grasslands, shrublands, wetlands, and coastal scrubland. Because of a lack of historical population data and the species' current, broad habitat use, key habitat variables are difficult to determine. Pueo occur in many areas that are managed by the State of Hawai'i or Federal agencies.

THREATS: Pueo are likely susceptible to the same factors that threaten other native Hawaiian birds, including: loss and degradation of habitat, predation by introduced mammals, exposure to anthropogenic threats such as rodenticides and collisions with vehicles and structures, and disease. However, their persistence in lowland, non-native and rangeland habitats suggests that they may be less vulnerable to extinction than other native birds, especially because they may be resistant to avian malaria (*Plasmodium relictum*) and avian pox (*Poxvirus avium*).

Despite this, for Pueo populations, the following are of particular concern:

- **Predation:** Because Pueo nest on the ground, eggs, young, and even adult birds are vulnerable to predation by rats (*Rattus* spp.), feral cats (*Felis silvestris*), feral dogs (*Canis lupus familiaris*) and the small Indian mongoose (*Herpestes auropunctatus*).
- **Habitat loss:** Though Pueo are found in a variety of habitat types Pueo use of heavily altered landscapes such as agricultural and urban areas may increase their exposure to anthropogenic risks such as contaminants and human interactions (see below). May be particularly important for O'ahu Pueo populations.
- **Contaminants or toxins:** Because Pueo are top predators, they are likely secondarily exposed to both first and second-generation anticoagulant rodenticides and face both lethal and sublethal effects from these toxicants. Increased use of urban, agricultural, and other human-altered environments puts pueo at greater risk of encountering these environmental toxins.
- **Human interaction:** Hunting behavior and habitat use predispose Pueo to collisions with infrastructure such as powerlines and vehicles,.
- **Climate change:** Pueo appear to be shifting their activity temporally and in elevation seasonally, potentially due to higher temperatures at low elevations.

CONSERVATION ACTIONS: Pueo likely have benefited from management activities designed to conserve other endangered birds, such as fencing, ungulate control, and invasive predator removal. They also may benefit from game bird management; high densities of Pueo occur on lands where game birds also are common. In addition to these efforts, future management specific to the Pueo may include the following:

- Determine population trends and abundance estimates through systematic surveys. Public outreach and education.
- Continue protection and management of wildlife sanctuaries and refuges.

MONITORING: Regular island-wide population surveys are necessary to determine population trends for this species. This information is needed to assess the efficacy of habitat management efforts.

RESEARCH PRIORITIES: Research priorities specific to Pueo include the following:

- Analysis of population trends and changes in habitat occupancy, especially on O‘ahu.
- Analysis of habitat selection to better understand if certain habitat types are of greater value to Pueo and meet different life history needs (for example, increased breeding success in certain habitat types over others)
- Determine primary causes of mortality and its potential effect on populations. Quantify the number of infrastructure, powerline and vehicular collisions and determine the level of threat to populations.

References:

- Berger AJ. 1981. Hawaiian birdlife. Honolulu: University of Hawai‘i Press. 260 pp.
- Holt DW, Leasure SM. 1993. Short-eared owl (*Asio flammeus*). In The Birds of North America, No. 62 (Poole A, Gill F, editors). Philadelphia, (PA): The Academy of Natural Sciences; and Washington DC: The American Ornithologists' Union.
- Mostello, C.S. and Conant, S., 2018. Diets of native and introduced apex predators in Hawai‘i. Pacific Conservation Biology, 24(1), pp.25-34.
- Scott JM, Mountainspring S, Ramsey FL, Kepler CB. 1986. Forest bird communities of the Hawaiian islands: their dynamics, ecology and conservation. Lawrence, (KS): Cooper Ornithological Society.
- Sierts, S.R, S. E Volker, C.G Payne, R. T. Sugihara, C.E. Swift, J.T. Nelson, A.B. Shiels, and W.C. Pitt. 2019. Screening of Hawaiian bird and bat carcasses for rodenticide residues. Unpublished final report (QA-2240). USDA, APHIS, WS, NWRC. Hilo, HI.
- Stormcrow, K., Kealiikanakaoleohaililani, K., Puniwai, N. Price, M.R., Vaughan, M.B. Winter, K.B. In Prep. ‘Ike ku‘una (Indigenous Knowledge) of Pueo (Hawaiian Short-eared Owl; *Asio flammeus sandwichensis*). Chapter in Owls in Myth and Culture. Springer? Johnson, D.H. (Editor).
- Stormcrow, K., Winter, K.B., Puniwai, N., Risch, D., Price, M.R. In Prep. Factors influencing detectability of Pueo (Hawaiian Short-eared Owl; *Asio flammeus sandwichensis*).
- Wang O, Garcia-Heras M-S., Wilhite CJ., Naguwa WH., Price MR. In Prep. Breeding ecology and management of Pueo (Hawaiian Short-eared Owl; *Asio flammeus sandwichensis*). Available at: <https://scholarspace.manoa.hawaii.edu/items/9a442660-1f0e-4a0c-8cae-620e214af80d>
- Wilhite CJ, Cotin J, Wang O, Garcia-Heras M-S, Idle JL, Rípodas Melero AI, Price MR. 2024. Diurnal preference for wooded habitat hints at intraspecific niche expansion in the Pueo (Hawaiian Short-eared Owl; *Asio flammeus sandwichensis*). Avian Conservation and Ecology.